

Assessment of seawater pollution by heavy metals in the neighbourhood of Algiers: use of the sea urchin, Paracentrotus lividus, as a bioindicator

Dina Soualili, Philippe Dubois, Pol Gosselin, Philippe Pernet, Monique

Guillou

▶ To cite this version:

Dina Soualili, Philippe Dubois, Pol Gosselin, Philippe Pernet, Monique Guillou. Assessment of seawater pollution by heavy metals in the neighbourhood of Algiers: use of the sea urchin, Paracentrotus lividus, as a bioindicator. ICES Journal of Marine Science: Journal du Conseil, 2008, 65 (2), pp.132-139. 10.1093/icesjms/fsm183. hal-00460169

HAL Id: hal-00460169 https://hal.univ-brest.fr/hal-00460169

Submitted on 26 Feb 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. The use of the sea urchin *Paracentrotus lividus* (L.) as bioindicator of the sea water quality of the Algerian coastal environment

Soualili D. Dubois Ph, Gosselin P., Pernet Ph., Guillou M.

Abstract

This study assessed marine contamination in heavy metals in the vicinity of the Algiers metropolis by combining chemical and toxicological data using the sea urchin *Paracentrotus lividus*. Metals were analyzed in the sediment and in the sea urchin gonads. The sediment toxicity was assessed by bioassays using sea urchin larval development. This study discriminated a site near the Algiers town as highly polluted in Pb. The contamination levels appeared to be toxic for the *P. lividus* larval development. The occurrence of other metals (Fe, Cd, Cu) was poor compared with that reported in the Mediterranean Sea, with exception of Zn that showed high values in female gonads.

Keywords : Algeria, bioassays, gonads, heavy metals, sediment, Paracentrotus lividus.

Soualili Dina : Laboratoire de Biologie et d'Ecologie Marine, Faculté des Sciences de la Nature, Université des Sciences et Technologie Houari Boumedienne, Bab-Ezzouar, Algiers, Algeria. Dubois Philippe, Pernet Philiipe : Laboratoire de Biologie Marine, Université Libre de Bruxelles, CP 160/15, Av. F.D. Roosevelt, B-1050 Bruxelles, Belgium Gosselin Pol : Laboratoire de Biologie Marine, Université de Mons-Hainaut, Av. du Champs de Mars, Pentagone, 7000Mons, Belgium Guillou Monique : Laboratoire des Sciences de l'Environnement Marin, UMR CNRS 6539, Institut Universitaire Européen de la Mer, Place N. Copernic, 29280 Plouzané, France.

Corresponding author : Guillou Monique, tel : 33 2 98 49 86 34 ; Fax 33 2 98 49 86 45

e-mail : mguillou@univ-brest.fr

1 Introduction

2 The Atlantic-Mediterranean urchin (Lamarck) sea Paracentrotus lividus 3 (Echinodermata: Echinoidea) is distributed from Ireland to southern Morocco, 4 including throughout the Mediterranean Sea. This opportunistic species has an 5 important ecological role in different ecosystems especially in Mediterranean Sea where 6 its grazing activity can locally control the dynamics of seaweeds (see review Lozano et 7 al., 1995) and seagrasses (see review Tomas et al., 2004). However human activities in 8 coastal zones can alter *P. lividus* morphology, biology and demography (Harmelin et 9 al., 1981; Delmas et Régis, 1984, Pancucci et al., 1992) and over the long-term can 10 modify its ecological role. The sedentary habit of P. lividus and its sensitivity to 11 pollutants has led to use this species as biological-biochemical indicator of local 12 pollution (Warnau et al. 1998, Coteur et al. 2001; Bayed et al. 2005). The sensitivity of 13 embryos and larvae of P. lividus prompted its use in embryotoxicity tests (Hagström 14 and Lönning, 1973; Pagano, 1986; Warnau et al. 1996). The embryonic and larval 15 development of sea urchins is one of the toxicity assays used in monitoring and risk 16 assessment programmes (see review Beiras et al., 2003).

On the Algerian coasts, *P. lividus* is a dominant species of the shallow water ecosystems where it can reach 25 individuals.m⁻² (Semroud, 1993). It colonizes different biotop like *Posidonia oceanica* and *Cymodocea nodosa* meadows, rocky substrate with photophile algae and overgrazed rocky substrate (Semroud, 1993 ; Guettaf, 2000). It lives from the low-water limit to about 10 m depth (Dieuzeide, 1933 ; Semroud, 1993).

22 Around the Bay of Algiers, the populations are exposed to anthropogenic disturbance.

23 More than 35% of the national littoral population live in the Algiers metropolitan area

(between Chenoua and Cap Djinet) or more than 4.3 million persons (950 habitants. 1 2 km-²) in a 115 km long coastal zone, This area is highly industrialized, concentrating about 25 % of the Algerian factories with about 1000 factories in 2001 (Larid, 2003; 3 4 PAC, 2005) including metallurgical, chemical, pharmaceutical, building material and 5 hydrocarbon industries and in a lower proportion mechanical, electric and electronic engineering industries and food and paper factories. A multitude of small polluting 6 activities, official or not, spreads also along the littoral. Two rivers flow in the bay of 7 Algiers, El Hamiz, and principally El Harrach (970 km⁻²; 6 m³ h⁻¹) which drain the 8 9 domestic and industrial waste waters of the Algiers city. Among the wastewaters which flow in the bay (225 millions m³ y⁻¹) only 8% are treated (PAC, 2005). The marine area 10 11 under the influence of the Algiers metropolis catches the highest pollution flow of the Algerian coasts (PAC, 2003) with 100 thousand t y^{-1} of organic chemical compounds, 12 175 thousands t y^{-1} of suspended matter, 1 500 t y^{-1} of nitrogen and 4 thousands t y^{-1} of 13 14 phosphorus (Larid, 2003). In 2004, 46 beaches of the Bay were prohibited from swimming. The sediments of Algiers Bay are highly polluted in heavy metals, 15 16 hydrocarbons and organic matter. The main metal detected in the coastal water are Hg, 17 Pb, Cu and Zn (PAC, 2003). The biological effects of the pollution on marine 18 ecosystem are important and the recent program on the management of the coastal 19 environment (PAC) pointed out a biodiversity decrease of 14% in the species of major 20 ecological interest (PAC, 2005).

The purpose of the present study was to assess the marine pollution in this area by combining toxicological and chemical data using the sea urchin *P. lividus*. Embryolarval sediment toxicity bioassays and metal contaminant analysis were used. The metal were analyzed in the sediment. The toxicity of their bioavailable fraction was detected by abnormalities in the *P. lividus* development as this species is a valuable indicator of metal contamination and can accumulate metals as a function of the contamination level of the environment (Warnau, 1998). The results of bioassays were compared to the analysis of metal contamination in the sediments and in the gonads of adult sea urchin inhabiting the sediments previously tested. Three sites were chosen in the vicinity of the Bay of Algiers according to potential sources of pollution. The information obtained will allow the ordination of the sampling stations in term of environmental quality.

7

8 Material and Methods

9

10 Sampling sites

Two stations, Algiers Beach and Tamentfoust, were located in the Bay of Algiers and 11 12 another Sidi Fredj, in the North of the Bay of Bou-Ismail (east of the Bay of Algiers) 13 (Fig. 1). The site of Algiers Beach should be the most polluted as it is the nearest of the 14 Algiers town. This site is characterized by a habitat mixing rocks with photophile algae 15 and degraded meadows. The site of Tamentfoust, situated in a half closed creek, is 16 characterised by overgrazed rocks with a sparse Posidonia oceanica beds. The Bay of 17 Bou-Ismail where the site of Sidi Fredj is located has pollution levels much lower than 18 the levels of the Bay of Algiers (PAC, 2005). More distant from the very industrialized 19 area, it encloses a marine medical centre and shows a better health meadow. But 20 Guehioueche and Zelmat (in PAC, 2005) indicated a progressive meadow regression 21 which could be due to the mechanical and organic disturbance caused by the medical 22 centre.

23

24 Collection and preparation of samples

Forty sea urchins of 45-65 mm of diameter were sampled by scuba diving in March
 2002 from each sites between 1 and 5 m depth. This period correspond to the maturity
 period of the gonads (Semroud and Kada, 1987; Guettaf, 2000) before the main
 spawning which can partly eliminate metal load in the gonad (Warnau et al., 1998).

Just after collection, the sea urchins were measured (ambital diameter) in the laboratory and dissected. The sex was determined and the gonads were dried at 60°C for 48h and stored separately in hermetically-sealed polyethylene containers. All the manipulations were performed with stainless steel instruments.

9 Concomitantly to the sea urchin sampling, the upper 2 cm layer of the sediments 10 inhabited by sea urchins was collected by coring (5 cm diameter). The samples of 11 sediments were placed into sealed polyethylene bags, carried to the laboratory on ice 12 and immediately dried at 60°C for 48h until constant weight then stored in hermetically-13 sealed polyethylene containers at room temperature.

14

15 Metal analysis

16 The concentration of zinc (Zn), lead (Pb), cadmium (Cd), copper (Cu) and iron (Fe) 17 were measuring in the gonads of the urchins and in the sediment according to the 18 method described by Coteur et al. (2003). As the sediment was very heterogeneous with 19 big gravels, only the fraction with grain sizes < 1mm was retained and considered in 20 this study as the 'total fraction'. Metal concentrations were also measured on the 21 fraction with grain sizes $< 63 \mu m$. This fraction allows to estimate the organic matter 22 content of the sediment. Pb, Cd and Cu concentrations were determined by graphite-23 furnace electrothermic atomic absorption spectrometry using a Varian GTA100 24 SpectrAA640Z spectrometer. Concentration of Zn and Fe were measured by flame 25 atomic absorption spectrometry using a GBC 906AA spectrometer. Accuracy of the

method was tested using certified reference material (*Mytilus edulis* tissues, BCR nr
278R from the Community Bureau of Reference, Commission of the European Union,
Brussels, Belgium). Detection limits for Zn, Pb, Cd, Cu and Fe were 0.002, 0.014,
0.001, 0.002 and 0.004 µg of metal per mL of digested sample respectively.
The gonads of 10 males and 10 females per site were analysed. For the sediment, 3 to 6
replicates of total fraction and 0 to 3 replicates of the fraction < 63 µm were used.

8 Embryo-larval bioassays

9 The toxicity of the sediments from the previous three sites was assessed by embryo-10 larval bioassays in May 2003. Natural sea water used for these tests was collected in 11 front of the marine station of Luc-sur-Mer (Normandy, France), the reference site of the 12 Laboratory of Marine Biology of the U.L.B. (Free University of Brussels, Belgium) 13 where the metal analysis where performed.

Sea-water was previously allowed to decant for 48 h, filtered by a 22 µm membrane and maintained at 20 ± 1°C (FSW, filtered seawater). Adult *Paracentrotus lividus* genitors were collected intertidally from a reference population inhabiting the rocky basins of Morgat (Bay of Douarnenez, Brittany, France). Individuals were transferred in the cultivation system of the marine laboratory of the U.M.H. (University of Mons-Hainaut, Belgium). For the embryotoxicity test, reference sediment was sampled at Wimereux (Nord-Pas-de-Calais, France).

The embryo-larval bioassays were performed at the laboratory of Mons-Hainaut. Those bioassays were based on the method described by Coteur *et al.* (2003). Spawning was induced by injection of 20 µl per gram of KCL 0.5N through the peristomial membrane. Gametes from 3 females and 3 males were collected in FSW and their quality (*i.e.*, general eggs' shape and sperm's motility) was checked under Olympus TO41 inverted

light microscope. Eggs from each female were fertilised by the pooled sperm from the 3 1 2 males. Embryos at early gastrula stage (4-5 h after fertilization at $14^{\circ}C \pm 1^{\circ}C$) from the 3 females were mixed before to be used for the bioassays (this pooling leads to one 3 4 bioassay per site using the same mixed pool of zygotes). Experiments with embryos 5 were performed in six-well plates (Falcon, ref. 35-3046). One plate was dedicated to each sampled sediments (Alger Plage, Tamenfoust, Sidi-Fredi and Wimereux). Each 6 7 well of those plates were filled with 0.1 g of dried sediment and 10 ml of FWS was 8 added before the bioassay. One plate filled with FWS was used as control. At the 9 beginning of the bioassay, batches of 250-300 embryos were transferred in each well. 10 After 72 h at $14 \pm 1^{\circ}$ C, larvae were fixed by adding 1 ml of formalin (commercial 11 solution, 35%) in each well and the plates were maintained in an oven at 70°C during 2 12 hours. Plates with fixed larvae were stored at 4°C. The frequency of developmental 13 stages was scored in each well on a random sample of 100 larvae. Developmental stages 14 were scored using an inverted light microscope according to the morphological criteria 15 adapted from Warnau and Pagano (1994). Our larvae were classified in 4 categories: 16 normal plutei ("Normal", N), retarded plutei presenting a delayed development 17 ("Retarded", R), abnormal plutei with skeletal malformations and/or gut abnormalities 18 ("Pathologic 1", P1) and embryos whose development ended at the blastula stage or the 19 gastrula stage ("Pathologic 2", P2). On our figures, the rates of "Viable" larvae were 20 obtained by summing the rates of "Normal" and "Retarded" larvae from each batch.

21

22 Statistical analyses

After arcsin transformation, the developmental rates of normal and viable larvae werecompared using a one-way ANOVA followed by a Bonferroni's test (Zar 1996).

Dunnet's tests were used to compare the different rates measured to their corresponding 1 2 controls (Zar 1996). Significant differences were determined at the 95% level. 3 Contamination levels in gonads were compared by 2-way analysis of variance 4 (ANOVA) and subsequent Tukey HSD multiple mean comparison test (effects: sex and 5 sampling site). 6 The relationship between contamination in gonad and sediment and percentage of viable 7 larvae was studied by factor analysis using the principal-component method with an 8 extraction matrix based on Pearson correlation coefficients and the "varimax" method 9 of factor rotation. 10 11 Results 12 13 Contamination level in the sediment and in the sea urchin gonads 14 The fraction of sediment $< 63 \,\mu\text{m}$ was more or less abundant according to the site, 0.03, 15 0.26, 1.56 % of the total fraction for Sidi Fredj, Tamenfoust and Algiers Beach 16 respectively. The sediment of Algiers Beach had an obviously higher mud content. 17 Tables 1 and 2 present the heavy-metal concentrations in the sediment. Only the levels 18 in the total sediment have been statistically compared because of the low level of mud 19 in the site of Tamentfoust and Sidi Fredj leading to a too low number of replicates in 20 these both sites. Fe, Pb and Zn concentrations in the total fraction of the sediment 21 differed according to the sampled sites. The site of Algiers Beach was the most 22 contaminated in Pb but the less contaminated in Fe and Zn. Fe concentrations were 23 significantly higher in the sediment from Sidi Fredj. 24 Tables 3 and 4 present the heavy-metal accumulation in the sea urchin gonads. Metal

25 concentrations in the gonads were compared according to sex and sampling site by 2-

1 way ANOVA. Cd and Zn concentrations were significantly higher in female than in 2 male gonads while Cu was more concentrated in male gonads $(p<10^{-4})$. Pb and Fe 3 concentrations did not significantly differ according to sex. Pb, Cu, and Fe 4 concentrations in the gonads differed according to the sampled sites $(p=10^{-3})$. Pb and Cu 5 concentrations were significantly higher in sea urchins collected respectively in Algiers 6 Beach and Sidi Fredj (Table 3). Fe concentrations in sea urchins from Tamentfoust 7 significantly differed from those of Algiers Beach No interaction was significant.

8

9 Embryo-larval development

A significant reduction of the percentages of both normal and viable larvae was only observed when embryos were exposed to sediment samples from Alger Beach (Fig. 2 A, B). The low rate of normal larvae induced in response to the sediment from Alger Beach is related to an increase of the rates of abnormal plutei (P1) and retarded plutei (R) (see Table 5). The high rate of retarded plutei induced in response to the sediment from Alger Beach (Table 5) explains why the induced rate of viable plutei remains relatively high (Fig 2B.).

17

18 Integrated data analysis

The metal concentrations in sediment, in male gonads, in female gonads and percentage of viable larvae were used in the factor analysis. The variables were distributed along axes representing factors according to the correlation coefficient of the variable with the factors (Fig. 3). The first two factors accounted for 99.9 % of the total variance of the data. On the x-axis, the concentrations in Fe, Cu and Zn in the sediment were opposed to the concentrations in Cd and Pb in the sediment. The relationships between the metal level in the sediment and the metal level in the gonads was dependent on the metal. For Pb the link was strong between sediment and the both gonads, for Cd and Cu it was high between sediment and the male gonads. On the x-axis the percentage of viable larvae was closed to the concentrations in Fe, Cu and Zn in the sediment (e.i located on the other end of the x-axis compared to the Pb and Cd contaminations in sediment and gonads). This suggests a close relationship between Pb and Cd levels in sediment and in gonads and a strongly negative interaction with these latter parameters and the percentage of viable larvae.

8 The y-axis comprised, at the negative end, the levels of Zn in the male and female 9 gonads, and the level of Fe in the female gonads, and at the positive end, the levels of 10 Cd and Cu in the female gonads. The percentage of viable larvae was located on the 11 negative part of this axis suggesting that this parameter was negatively intercorrelated 12 with Cd and Cu levels in the female gonads.

13

14 Discussion

15 The main goal of the present study was to assess the marine pollution in the vicinity of 16 the Bay of Algiers by combining chemical and toxicological data using the sea urchin *P*. 17 *lividus* as bioindicator.

18 No site was distinguishable from others by a generalized higher concentration of metals 19 in the total sediments or in the sea urchin gonads. The sediments of Algiers Beach was 20 separated from the others by the highest concentration in Pb but the lowest 21 concentrations in Zn and Fe, the site of Sidi Fredj by the highest concentration in Fe. 22 Algiers Beach is the most muddy site but its present a significant increase in metal 23 contamination only for a metal, Pb, while usually the load of metal in the sediments 24 increase along with the percentage of the fine fraction. This fact can be due to the low 25 sediment contamination in the other metals (see below). In the gonads, Pb also isolated the population of Algiers Beach from the others. In this site, Fe was low but not
 different from Sidi Fredj. This latter site presented the highest concentrations in Cu. Cd
 did not discriminate any site.

In the sediment, Cd and Cu values can be considered in the background concentrations 4 of the Mediterranean range. Cd values were ranged between to 0.12 to 0.76 μ g g⁻¹ dry 5 6 wt in the total fraction of the sediment while the background concentrations of the Mediterranean were estimated between 0.05 and 1 μ g g⁻¹ dry wt (EEA, 1999). We note 7 8 however a higher concentration in the fraction with grain sizes $<63 \mu m$ of the site of Algiers Beach (1.22 $\mu g g^{-1}$ dry wt). The Cu values ranged from 4.1 to 6.4 $\mu g g^{-1}$ dry wt 9 in the total fraction of the Algerian sediment and to 20.9 μ g g⁻¹ in the fraction with grain 10 11 sizes <63 µm (Algiers Beach) and the background concentrations of the Mediterranean ranged between 5 to 30 μ g g⁻¹ (EEA, 1999). The Zn (between 0.02 and 0.08 μ g g⁻¹ dry 12 wt) and Fe concentrations (between 7.1 and 41.5 $\mu g g^{-1}$) were very low compared to 13 values usually observed in the Mediterranean Sea ranged from 35 to 150 $\,\mu g \; g^{\text{-1}}$ for Zn 14 (Saad et al. 1981, Hoogstraten and Nolting, 1991; Storelli et al. 2001) and $> 10^3 \ \mu g \ g^{-1}$ 15 for Fe (Saad et al. 1981, Storelli et al. 2001, Menchi et al., 2002). Only Pb values in the 16 total fraction of the sediment of Algiers Beach (39.6 μ g g⁻¹) and in the fraction with 17 grain sizes $<63 \mu m$ of Sidi Fredj (40.6 $\mu g g^{-1}$) exceeded the background concentrations 18 of the Mediterranean range detected between 5.2 to 23.2 $\mu g g^{-1}(EEA, 1999)$ or the 19 background (4-17 μ g g⁻¹) given by the reference tables of NOAA (Buchman, 1999). 20

In the gonads, comparisons with *P. lividus* data available from the literature (Table 6) and especially the study of Warnau et al (1998), confirms the conclusions on the level of metal contamination deduced from the sediment analysis except for Zn. The gonad concentrations in Cd, Cu and Fe from the three sampling locations averaged background concentrations in *P. lividus* gonads. Only the gonad contamination in Zn from the 3

sites and in Pb from Algiers Beach were obviously higher that the Zn and Pb 1 2 background concentrations for this species. The Zn concentrations, not significantly different among the sites, were high, comparable to the concentrations estimated in 3 female gonads of *Sphaerechinus granularis* from the bay of Bay of Brest (between 190 4 and 700 μ g g⁻¹) (Guillou et al. 2000). This bay is known to be contaminated by this 5 metal which covers the whole roof of the town (Troadec, 1995). But in the present study 6 7 the Zn gonad levels appeared disproportionate with the Zn levels in sediment indicating 8 that sediment possibly is not the most contaminated abiotic compartment. But as Zn was 9 dominant in the gonads of the females, the results about this essential element for 10 animal metabolism must be cautiously interpreted (Hambidge et al., 1986). On the 11 contrary, the high Pb levels in gonads reflected the high Pb levels in the sediment. 12 Among the compared sites, only Rabat (Morocco) presented higher concentrations due 13 to the untreated pottery activity of this town (Bayed et al., 2005).

14 However when metal discriminates the site in term of contamination, the response given 15 by the sediment can differ from that given by the gonad. The geographical Pb gradient 16 observed in the sediment matches closely the gradient reported by the gonad 17 accumulation, pointing out the site of Algiers Beach as the most contaminated in this 18 metal. But Fe and Zn bioaccumulation did not discriminate the sites in the same way as 19 Fe and Zn sediment concentrations. Three explanations can explain the differential 20 metal pattern: i) the bioavailability of the metals is different ii) essential elements as Fe 21 and Zn have different accumulation pattern that the other metals iii) Fe, Zn and Cu 22 concentrations were too low in the sediment to be suitably expressed in the biological 23 compartment. Although the two first hypothesis are plausible, the third one is clear (see 24 above).

1 The high contamination level in Pb in the site of Algiers Beach expressed by the 2 sediment and gonad analysis was confirmed by a recent study in the water and stream 3 sediment of Oued El Harrach which flows in the Algiers Bay. Levels of 21 to 41 μ g g⁻¹ 4 dry wt, close to the levels detected in the total fraction of the sediment of Algiers Beach 5 (39.6 μ g g⁻¹) were detected in the sediments in the mouth of the Oued. This pollution 6 was probably caused by the discharge of an Algiers untreated industrial wastewater 7 (Yoshida et al., 2005).

8 The results of the bioassays pointed out Algiers Beach as the only site showing toxicity on sea-urchins embryos with 24.3 % of abnormal larvae (vs 7.9 \pm 2.8 % for the 3 other 9 10 sites) and 47,8 % of retarded larvae (vs 22.56 \pm 4.4 % for the 3 other sites). No 11 difference was detected between the two other Algerian sites and the reference site and 12 the control. According to the Kobayashi criteria (1991), the level of normal plutei of 13 Algiers Beach (20.5%) expressed a strong inhibition of the sea urchin development in 14 relation with a high disturbance. The results of these assays appeared negatively related 15 with the Pb concentrations in the male and female gonads which were positively 16 correlated with the Pb concentration in the total fraction of the sediment. The integrated 17 data analysis confirms these conclusions. The high positive relationship between the 18 rate of viable larvae and the Cu and Fe concentrations in the sediment and male gonad 19 and the Zn concentration in the sediment would not express a positive effect of these 20 metals on the larval development but more precisely a lack of negative effect du to the 21 low levels of these metals. Nevertheless the position of the viable larvae suggests a 22 negatively correlation with Cd and Cu levels in the female gonads. Although these 23 contamination levels were low, a possible higher influence of the female go ad on the 24 larval viability would be considered when these gonads were contaminated by metals so 25 toxic as Cd and Cu.

The Pb pollution was not spread out as the sediment and the sea urchins of the site of Tamenfoust, a little farther from the Oued El Harrach than the site of Algiers Beach, did not present high Pb accumulation or developmental anomalies. In a semi- closed creek, this former site could be protected from the sea water and sediment flows coming from the Oued. The site of Sidi Fredj considered as less contaminated because more distant from the very industrialized area influenced by Algiers metropolis cannot be distinguished from the site of Tamenfoust according to the criteria used in this study.

8

9 In conclusion, this study confirms the link between developmental abnormalities, metal 10 accumulations in the gonads and metal contamination in the sediment when the metal 11 concentration in sediment is sufficiently high. It accredits the indicator value of *P*. 12 *lividus*. In the present study the analysis discriminates a site and a metal, Pb, but the 13 method could be improved by increasing the number of toxicants analysed and by 14 extending the metal analysis to other sea urchin storage organs as the gut which can 15 better reflect a heavy metal accumulation (Warnau et al, 1998; Guillou et al, 2000).

16

17 Acknowledgments.

18 This paper is a part of the thesis of D. Soualili; this study was carried out in the 19 « Laboratoire de Biologie Marine de l'Université Libre de Bruxelles, ULB », in the 20 « Laboratoire de Biologie Marine de l'Université de Mons-Hainaut, UMH » (Belgium), 21 and ended in the «laboratoire des Sciences de l'Environnement Marin de l'Université 22 de Bretagne Occidentale, UBO » (France). The study in Belgium was funded by a grant 23 of the "Institut des Sciences Agronomiques et Vétérinaires de l'Université de Blida 24 (Algeria)", the stay in France by a grant of the Algerian Department of Higher 25 Education and Research. D. Soualili gratefully acknowledges Professeur M. Jangoux

References

Bayed, A., Quiniou, F., Benrha, A., Guillou, M. 2005. The *Paracentrotus lividus* populations from the Northern Moroccan Atlantic coast: growth, reproduction and health condition. Journal of Marine Biological Association of the United Kingdom 85, 999-1007.

Beiras, R., Fernandez, N., Bellas, J., Besada, V., Gonzalez-Quijano, A., Nunes, T. 2003. Integrative assessment of marine pollution in Galician estuaries using sediment chemistry, mussel bioaccumulation, and embryo-larval toxicity bioassays. Chemosphere 52(7), 1209-1224.

Buchman, M.F. 1999. NOAA Screening Quick Reference Tables, NOAA HAZMAT Report 9-1, Seattle WA, Coastal Protection and Restoration Division, National Oceanic and Atmospheric Administration, 12pp.

Coteur, G., Danis, B., Fowler, S.W., Teysiié, J.L., Dubois, Ph., Warnau, M. 2001. Effects of PCB's on reactive oxygen species (ROS) production by the immune cells of *Paracentrotus lividus* (echinodermata). Marine Pollution Bulletin 42, 667-672.

Coteur, G., Gosselin, P., Wantier, P., Chambost-Manciet, Y., Danis, B., Pernet, Ph., Warnau, M., Dubois, Ph. 2003. Echinoderms as bioindicators, bioassays and impact assessment tools of sediment-associated metals and PCBs in the North Sea. Archives of Environmental Contamination and Toxicology 45, 190-202.

Dieuzeide, R. 1933. Les Echinoides réguliers de la baie de Castiglione. Bulletin des Travaux Scientifiques de la Station Aquacole et de Pêche de Castiglione, 1-9.

European Environment Agency (EEA), 1999. Environmental State and Threats. In: State and pressures of the marine and coastal Mediterranean environment. UNEP, 6-104.

Guettaf, M., Gustavo, A., San Martin, G.H., Francour, P. 2000. Interpopulation variability of the reproductive cycle of *Paracentrotus lividus* (Echinodermata: Echinoidea) in the south-western Mediterranean. Journal of Marine Biological Association of the United Kingdom 80, 899-907.

Guillou, M., Quiniou, F., Huart, B., Pagano, G. 2000. Comparison of embryonic development and metal contamination in several populations of the sea urchin *Sphaerechinus granularis* (Larmarck) exposed to anthropogenic pollution. Archives of Environmental Contamination and Toxicology 39, 337-344.

Hagström, B.E., Lönning, S. 1973. The sea urchin egg as testing object in toxicology. Acta Pharmacologica and Toxicologica 13 : 7-49.

Hambrige, K.M., Casey, C.E., Krebs, N.F. 1986. Zinc. In: Mertz, W. (Ed), Trace elements in human and animal nutrition. Vol 2, 5th edn., Academic Press, Orlando, FL.

Hoogstraten van, R.J., Nolting, R.F. 1991. Trace and major elements in sediments and inporewaters from the North Western basin of the Mediterranean Sea. Netherlands Institute for Sea Research (NIOZ), Rapport 1991 10, 1-72.

Kobayashi, N. 1991. Marine pollution bioassay by using sea urchin eggs in the Tanabe Bay, Wakayama Perfecture, Japan, 1970-1987. Marine Pollution Bulletin 23, 709-713.

Larid, M. 2003. Analyse de la durabilité dans le cadre du PAC "Zone côtière algéroise" (Algérie). Rapport de la première étape 36 pp.

Lozano, J., Galera, J., Lopez, S., Turon, X., Palacin, C., Morera, G. 1995. Biological cycles and recruitment of *Paracentrotus lividus* (Echinodermata: Echinoidea) in two contrasting habitats. *Marine Ecology Progress Series* 122, 179-191.

Menchi, V., Balocchi, C., Pozo, K., Perra, G., Graziozi, M., Focardi, S. 2002. Distribution of PCBs, PAHs and heavy metals in bottom sediments of the Eastern Mediterranean Sea. In : Challenges in Environmental risk Assessment and Modelling, Proceedings of the SETAC, Europe 12 th Annual Meeting. Vienna, Austria p.147.

Pagano, G., Cipollaro, M., Corsale, G., Esposito, E., Ragucci, E., Giordano G.G., Trieff, N.M. 1986. The sea urchin: Bioassay for the assessment of damage from environmental contaminants. In : Cairns, J. JR. (ed), Community toxicity testing. American Society for testing and materials, Philadelphia, PA, USA, 66-92.

PAC (Rapport d'Aménagement Côtier) 2003. Rapport sur l'état et l'avenir de l'environnement. Ministère de l'Aménagement du Territoire et de l'Environnement. République Algérienne Démocratique et Populaire. 463pp.

PAC (Rapport d'Aménagement Côtier) 2005. "Zone côtière algéroise". Protection des sites sensibles naturels marins du secteur Cap Djinet au Mont Chenoua. Impact des activités anthropiques. Ministère de l'Aménagement du Territoire et de l'Environnement. République Algérienne Démocratique et Populaire. 88 pp.

Pancucci, M.A., Panayotidis, P., Zenetos, A. 1993. Morphological changes in sea urchin populations as a response to environmental stress. In Aldrich, J.C. (ed), Quantified phenotypic responses in morphology and physiology. JAPAGA, Ashford. pp. 247-257.

Saad, M.A., El-Rayis, O.A., El-Nady, F.E. 1981. Occurrence of organic matter and heavy metals in sediments from the Mediterranean. In : Stuckey, D., Hamza, A. (eds), Management of Industrial Wastewater in Developing Nations. Pergamon Press, Oxford. pp.127–139.

Semroud, R. 1993. Contribution à la connaissance de l'écosystème à *Posidonia oceanica* (L) Delile dans la région d'Alger (Algérie): étude de quelques compartiments. PhD thesis, Institut des Sciences de la Nature, Université des Sciences et de la Technologie Houari Boumediene, Algeria.

Semroud, R., Kada, H. 1987. Contribution à l'étude de l'oursin *Paracentrotus lividus* (Larmarck) dans la région d'Alger (Algérie): indice de réplétion et indice gonadique. In:

Boudouresque, C.F. (ed), Colloque international sur *Paracentrotus lividus* et les oursins comestibles. GIS Posidonie Publications, Marseille, France, pp 117-124.

Storelli, M.M., Storelli, A., Marcotrigiano, G.O. 2001. Heavy metals in the aquatic environment of the Southern Adriatic Sea, Italy Macroalgae, sediments and benthic species. *Environment International* 26, 505-509.

Tomas, F., Romero, J., Turon, X. 2004. Settlement and recruitment of the sea urchin *Paracentrotus lividus* in two contrasting habitats in the Mediterranean. Marine Ecology Progress Series 282, 173-184.

Troadec, P. 1995. La qualité du milieu marin: la rade de Brest. Communauté urbaine de Brest, rapport interne, 165 pp.

Warnau, M., Pagano, G. 1994. Developmental toxicity of PbC12 in the echinoid *Paracentrotus lividus* (Echinodermata). Bulletin of Environmental Contamination and Toxicology 53, 434-441.

Warnau, M., Iaccarino, M., De Biase, A., Temara, A., Jangoux, M., Dubois, Ph. 1998. Spermiotoxicity and embryotoxicity of heavy metals in the echinoid *Paracentrotus lividus*. Environmental Toxicology and Chemistry 15, 1931-1936.

Warnau, M., Biondo, R., Temara, A., Bouquegneau, J.M., Jangoux, M., Dubois, Ph. 1998. Distribution of heavy metal in the echinoid *Paracentrotus lividus* (Lmk) from the Mediterranean *Posidonia oceanica* ecosystem: seasonal and geographical variations. Journal of Sea Research 39, 267-280.

Yoshida, M., Moali, M., Houas, O., Lakhdari, M., Nechaoui, L., Guerrida, D., Chatal, A., Oussalem, S., Makour, F., Khelifi, F., Laleg, A. 2005. Environmental Pollution in Oued El Harrach area, Alger A Preliminary Report on Mercury and Heavy Metals Contaminations. Compte-Rendu du Séminaire "Pollution et Protection de l'Environnement en Algérie". ONEDD, JICA, Alger, pp 19-37.

Zar, J.H. 1996. Biostatistical analysis. Prentice-Hall Inc., Upple Saddle River, New Jersey.

Legends of figures



Figure 1 : Location of the three sampling sites: Algiers Beach, Tamenfoust and Sidi Fredj.



Figure 2. Percentages (mean \pm S.D.) of normal (**A**) and viable (**B**) larvae of *Paracentrotus lividus* after exposition to dried sediments and control throughout embryogenesis (72 h). 6 replicates by exposure, 100 larvae scored by replicate. For each species, there were no significant differences between the series designed by the same letter (Bonferroni, p = 0.05).



Fig. 3. Factor analysis (principal-component method) showing the relationships between the metal concentrations in the sediment (metal and suffix SED), in the male gonads (metal and suffix MG), in the female gonads (metal and suffix FG) and the percentage of viable larvae (VIABLARV). The first (x-axis) and the second (y-axis) factors accounts 70% and 29.9%. of the total variance respectively.

Legends of tables

Table 1. Metal concentration (mean \pm SD ; $\mu g g^1$ dw) in the total fraction (n =3 to 6) and in the <63 μ m grain-size fraction (n=0 to 3) of the sediment collected in the three Algerian sites (AB: Algiers Beach; TM: Tamenfoust; SF: Sidi Fredj).

Metals in sediments		Zn	Pb	Cu	Cd	Fe
Total fraction						
AB	Means	0.023	39.63	4.08	0.76	7.11
	SD	0.005	7.93	0.62	1.05	2.18
TM	Means	0.045	14.59	5.73	0.15	17.85
	SD	0.010	1.55	1.65	0.06	3.71
SF	Means	0.050	12.37	6.43	0.12	31.32
	SD	0.006	4.07	1.32	0.01	6.49
<63 µm fraction						
AB	Means	0.081	23.76	20.89	1.22	19.70
	SD	0.008	0.75	0.75	0.04	2.08
SF	Means	0.08	40.63	10.48	0.25	41.50
	SD					

Metal in sediments	p ANOVA	Level of contamination ^b		
Total fraction		+		
Zn	0.03	<u>SF TM</u> AB		
Pb	> 10 ⁻²	AB <u>TM SF</u>		
Cu	NS ^a	TM SF AB		
Cd	NS			
Ea	× 10 ^{−2}	<u>AB TM SF</u>		
ГС	> 10	SF TM AB		

Table 2. Comparison of the metal concentrations in the total fraction of the sediments measured in the three Algerian sites (AB: Algiers Beach; TM: Tamenfoust; SF:Sidi Fredj) .

^a NS, no significant difference ^b stations which do not differ in metal concentration are underlined (p >0.05, Tukey HSD test)

Metals in gonads		Zn	Pb	Cu	Cd	Fe
AB						
Females	Means	385.5*	6.14	2.84	0.14	73.8
	SD	344.1	3.46	0.97	0.08	35.5
Males	Means	32.9	7.78	3.19	0.08	19.3
	SD	13.5	8.77	0.83	0.04	19.7
TM						
Females	Means	538.2*	1.5	2.49*	0.12	113
	SD	324.3	1.72	0.47	0.08	37.6
Males	Means	76.1	0.88	3.88	0.05	112.6
	SD	172.2	0.44	0.84	0.01	66
SF						
Females	Means	366.9	0.68	3.42	0.14*	71.1
	SD	178.3	0.12	0.85	0.09	54.8
Males	Means	52.9	0.90	4.42	0.05	92.7
	SD	73.2	0.41	0.56	0.03	78.8

Table 3. Metal concentrations (mean \pm SD; μ g g⁻¹ dw; n=10) in the gonads of *Paracentrotus lividus* collected in the three Algerian sites (AB: Algiers Beach; TM: Tamenfoust; SF: Sidi Fredj).

* indicates a significant difference between the sexes

Table 4 Comparison of metal concentrations in the gonads of *Paracentrotus lividus* measured in the three Algerian sites (2-factor ANOVA: sex and site) (AB: Algiers Beach; TM: Tamenfoust; SF:Sidi Fredj).

Metal in gonads	Sex	p ANOVA Site	Interaction	n	Level of contamination ^b
Zn	<10-4	0.26 ^a	0.53	60	TM SF AB
Pb	0.69 ^a	<10-4	0.66	60	AB TM SF
Cu	$< 10^{-4}$	<10 ⁻²	0.11	60	
Cd	<10 ⁻⁴	0.45 ^a	0.80	60	SF TM AB
Fe	0.40 ^a	<10 ⁻³	0.06	60	SF AB TM
					TM SF AB

^a no significant difference

^b stations which do not differ in metal concentration are underlined (p > 0.05, Tukey HSD test)

	1		1 0	50			
	Ν	R	P1	P2	Du1	\mathbf{V}	Du2
Control	59.2 ± 9.7	23.7 ± 5.9	8.7 ± 4.7	8.5 ± 3.2		82.8 ± 6.3	
					= 0.6		= 0.05
Wimereux	61.7 ± 5.8	27.7 ± 6.2	5.0 ± 2.3	5.7 ± 2.2		89.3 ± 2.4	
					= 0.6		= 0.4
Sidi Fredj	60.7 ± 8.1	19.8 ± 2.6	10.7 ± 3.1	8.8 ± 4.2		80.5 ± 6.8	
					= 0.6		= 0.5
Tamentfou							
st	62.2 ± 6.6	20.2 ± 3.9	8.2 ± 3.3	9.5 ± 2.6		82.3 ± 4.8	
					< 0.0005		< 0.0005
Algiers	$20.5 \pm$	$47.8 \pm$					
Beach	17.4	13.5	24.3 ± 7.9	7.3 ± 3.3		68.3 ± 6.4	

Table 5. Frequencies (means \pm standard errors) of developmental stages in *Paracentrotus lividus* larvae exposed to sediments samples throughout embryogenesis.

N: normal plutei; R: retarded plutei; P1: abnormal plutei; P2; Blastula; V: viable plutei; Du1: result of the Dunnett test comparing the rate of normal plutei to the control; Du2: result of the Dunnett test comparing the rate of viable plutei to the control.

A .1	a:	7	DI	0	01	-		
Authors	Sites	Zn	Pb	Cu	Cd	Fe		
(sampling date)								
Present study	Algiers Beach							
March 2002	Females	385.5	6.14	2.84	0.14	73.8		
	Males	32.9	7.78	3.19	0.08	19.3		
	Tamentfoust							
	Females	538.2	1.5	2.49	0.12	113		
	Males	76.1	0.88	3.88	0.05	112.6		
	Sidi Fredj							
	Females	366.9	0.68	3.42	0.14	71.1		
	Males	52.9	0.90	4.42	0.05	92.7		
Bayed et al 2005	Atlantic Morocco							
March 2000	Rabat							
	Females		<u>35.32</u>	3.34	25.15			
	Males		12.41	0.52	2.58			
	Bouznika							
	Females		11.3	<u>5.56</u>	2.21			
	Males		7.18	1.52	1.51			
	Mohammedia							
	Females		5.22	2.51	2.24			
	Males		6.14	1.18	2.32			
Warnau et al	Calvi	124	2.25	0.15	3.47	51		
1998	Ischia	140	3.02	0.41	3.41	90		
	Marseille	109	3.68	0.19	3.51	39		
Storelli et al 2001	Adriatic Sea	157.1	0.86	0.24	5.19	18.37		
April 1998								

Table 6. Comparison of the mean concentrations ($\mu g g^{-1} dw$) of metals in the gonads of *Paracentrotus lividus*.

The higher values are in bold and underlined