

**Molecular identification of differentially regulated genes
in the hydrothermal-vent species *Bathymodiolus
thermophilus* and *Paralvinella pandorae* in response to
temperature**

Isabelle Boutet, Didier Jollivet, Bruce Shillito, Dario Moraga, Arnaud Tanguy

► **To cite this version:**

Isabelle Boutet, Didier Jollivet, Bruce Shillito, Dario Moraga, Arnaud Tanguy. Molecular identification of differentially regulated genes in the hydrothermal-vent species *Bathymodiolus thermophilus* and *Paralvinella pandorae* in response to temperature. *BMC Genomics*, BioMed Central, 2009, 10, pp.222-. <10.1186/1471-2164-10-222>. <hal-00645893>

HAL Id: hal-00645893

<http://hal.univ-brest.fr/hal-00645893>

Submitted on 28 Nov 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Research article

Open Access

Molecular identification of differentially regulated genes in the hydrothermal-vent species *Bathymodiolus thermophilus* and *Paralvinella pandorae* in response to temperature

Isabelle Boutet^{1,2}, Didier Jollivet^{1,2}, Bruce Shillito³, Dario Moraga⁴ and Arnaud Tanguy*^{1,2}

Address: ¹CNRS, UMR 7144, Adaptation et Diversité en Milieu Marin, Station Biologique, de Roscoff, 29682 Roscoff, France, ²UPMC Univ Paris 06, UMR 7144, Equipe Génétique et Adaptation en Milieu Extrême, Station Biologique de Roscoff, 29682 Roscoff, France, ³UPMC Université Paris 6, UMR 7138, Systématique, Adaptation et Evolution, 75005 Paris, France and ⁴UMR CNRS 6539, Laboratoire des Sciences de l'Environnement Marin, Institut Universitaire Européen de la Mer, Université de Bretagne Occidentale, Place Nicolas Copernic, 29280 Plouzané, France

Email: Isabelle Boutet - boutet@sb-roscoff.fr; Didier Jollivet - jollivet@sb-roscoff.fr; Bruce Shillito - Bruce.Shillito@snv-jussieu.fr; Dario Moraga - dario.moraga@univ-brest.fr; Arnaud Tanguy* - atanguy@sb-roscoff.fr

* Corresponding author

Published: 13 May 2009

Received: 24 July 2008

BMC Genomics 2009, 10:222 doi:10.1186/1471-2164-10-222

Accepted: 13 May 2009

This article is available from: <http://www.biomedcentral.com/1471-2164/10/222>

© 2009 Boutet et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Hydrothermal vents and cold seeps represent oases of life in the deep-sea environment, but are also characterized by challenging physical and chemical conditions. The effect of temperature fluctuations on vent organisms in their habitat has not been well explored, in particular at a molecular level, most gene expression studies being conducted on coastal marine species. In order to better understand the response of hydrothermal organisms to different temperature regimes, differentially expressed genes (obtained by a subtractive suppression hybridization approach) were identified in the mussel *Bathymodiolus thermophilus* and the annelid *Paralvinella pandorae irlandei* to characterize the physiological processes involved when animals are subjected to long term exposure (2 days) at two contrasting temperatures (10° versus 20°C), while maintained at *in situ* pressures. To avoid a potential effect of pressure, the experimental animals were initially thermally acclimated for 24 hours in a pressurized vessel.

Results: For each species, we produced two subtractive cDNA libraries (forward and reverse) from sets of deep-sea mussels and annelids exposed together to a thermal challenge under pressure. RNA extracted from the gills, adductor muscle, mantle and foot tissue were used for *B. thermophilus*. For the annelid model, whole animals (small individuals) were used. For each of the four libraries, we sequenced 200 clones, resulting in 78 and 83 unique sequences in mussels and annelids (about 20% of the sequencing effort), respectively, with only half of them corresponding to known genes. Real-time PCR was used to validate differentially expressed genes identified in the corresponding libraries. Strong expression variations have been observed for some specific genes such as the intracellular hemoglobin, the nidogen protein, and Rab7 in *P. pandorae*, and the SPARC protein, cyclophilin, foot protein and adhesive plaque protein in *B. thermophilus*.

Conclusion: Our results indicate that mussels and worms are not responding in the same way to temperature variations. While the results obtained for the mussel *B. thermophilus* seem to indicate a metabolic depression (strong decrease in the level of mRNA expression of numerous genes)

when temperature increased, the annelid *P. pandorae* mainly displayed a strong regulation of the mRNA encoding subunits and linkers of respiratory pigments and some proteins involved in membrane structure. In both cases, these regulations seem to be partly due to a possible cellular oxidative stress induced by the simulated thermal environment (10°C to 20°C). This work will serve as a starting point for studying the transcriptomic response of hydrothermal mussels and annelids in future experiments in response to thermal stress at various conditions of duration and temperature challenge.

Background

The distribution of terrestrial, as well as marine, organisms is strongly influenced by environmental factors (variation, gradient, intensity), and temperature was identified as one of the most important. Temperature is known to affect the spatial distribution of species according to their thermal tolerance [1]. Environmental temperature challenge has a direct impact on ectothermic marine animals, and influences biological functions at all levels, from molecules to whole organisms [2,3]. Variation and/or gradients of temperature are considered as strong selective factors [4,5]. Numerous studies dealing with the effect of temperature have been conducted on coastal marine species at the molecular, physiological and protein levels [6,7]. An elevation of temperature can increase reaction rates and affect reaction equilibrium through higher kinetic energy. In addition, high temperatures induce protein denaturation, resulting in complete and often irreversible loss of function [2].

Hydrothermal vents and cold seeps represent oases of life in the deep-sea environment but are also characterized by challenging environmental conditions, when compared to the surrounding deep-sea. The hydrothermal fluid is the result of chemical modifications of the deep-sea water by interaction with the hot rocks (near the magma chamber) during a long percolating period through the oceanic crust. The resulting fluid is a hot water (up to 400°C), often anoxic, acidic (pH 2), and containing high concentrations of methane, carbon dioxide, sulfide, heavy metals, and arsenic-containing compounds [8-10]. As it comes out of the sea-floor, the hydrothermal fluid is chaotically mixed with the cold, more oxygenated deep-sea water. Abiotic factors such as changes in fluid flow, temperature and chemical composition affect species distributions at vents [11-13]. The endemic hydrothermal vent organisms live in the mixing zone according to their tolerance to – and requirements for– various factors (temperature, pH, O₂ concentration...). However, the effect of temperature variations on hydrothermal vent organisms has not been well studied, particularly at the molecular level. Available data on the vent mussel of genus *Bathymodiolus* are mainly focused on the effect of heavy metals and oxidative stress on enzymatic activities [14] and specific gene expression (e.g. metallothioneins) [15]. A few

immune system-relevant genes have also been described [16], and a cDNA library from *Bathymodiolus azoricus* has been partially sequenced [17]. Numerous studies also described these mussels' symbiosis with chemoautotrophic bacteria, often at both the phylogenetic and the biochemical levels [18-20]. HSP70, a typical thermal stress marker, has also been studied in hydrothermal vent species in response to temperature and other parameters. The HSP70 gene has been characterized in the vent shrimps *Mirocaris fortunata* and *Rimicaris exoculata* [21], in the vent annelid *Paralvinella grasslei* [22], and variations in HSP70 protein content were correlated to variations of environmental parameters for the vent mussel *B. azoricus* [23]. In Alvinellidae, a polychaete family endemic of hydrothermal vents that includes the genera *Alvinella* and *Paralvinella*, hemoglobins were largely studied at both the molecular and the functional levels [24]. Recent studies demonstrated the temperature preference of vent worm species by incubating them along a thermal gradient ranging from 15 to 60°C [25,26]. The existence of thermally sensitive alleles was shown for some proteins in *P. pandorae* compared to other species living at higher temperatures, such as *A. pompejana*, *P. sulfincola* or *P. hessleri* [27]. These data could be partly related to the species distribution in their respective habitats that mostly differ by their temperature range.

The vent mussel *B. thermophilus* is found under a wide range of environmental conditions on the East Pacific Rise. It may occur with tubeworms in areas of active hydrothermal flow where temperature and hydrogen sulfide concentrations are high, as well as in areas of diffuse flow, where the temperature and the concentration of hydrogen sulfide are much lower [28]. Studies have shown that *B. thermophilus* can quickly adapt to a wide range of environmental conditions (especially chemical changes) but usually lives between 4 and 14°C [29]. No studies report a thorough characterization of *P. pandorae irlanidei*'s habitat but this species is mainly found in the cracks at the base of the tubes of the vestimentiferan tubeworm *Tevnia jerichonana* [30] where it encounters temperatures ranging from 5 to 22°C [31]. Because temperature highly fluctuates in time, a thermal regime of 20°C therefore represents an upper thermal condition for both species whereas an average temperature of 10°C could be

more related to their thermal preference. In this work, we used a suppression subtractive hybridization (SSH) approach to (1) verify if such a temperature challenge induces a transcriptomic response, (2) characterize genes and the physiological processes involved in response to temperature variations, and (3) compare common and specific responses attributable to these two highly-divergent species.

Results

Mussel SSH libraries sequencing results

The sequencing of 200 clones from the forward (individuals incubated at 10°C versus individuals exposed at 20°C) SSH library, as well as from the reverse library (individuals incubated at 20°C versus individuals exposed at 10°C), allowed the identification of 78 unique sequences for *B. thermophilus* (19% of the whole sequencing effort: other sequences are redundant). Table 1 shows the unique sequences obtained in both SSH libraries with the best E-values for the sequences when identified after Blast analysis. The identified sequences indicate that temperature regulates genes involved in various cell functions: (1) cell cycle regulation, DNA repair, protein regulation and transcription (8.5%), (2) mitochondrial respiratory chain (4%), (3) metabolism (6.4%), (4) stress response and detoxification (0.5%), (5) cell communication, membrane receptors and immune system (10.2%), (6) cytoskeleton production and maintenance (7.5%), (7) ribosomal proteins (10%), (8) proteins of unknown function (0.5%) and (9) unknown sequences (52%). A higher proportion (60%) of the genes identified in our libraries were expressed in mussel exposed to 10°C and were mainly involved in metabolism, stress response, transcription and cytoskeleton.

Annelid SSH libraries sequencing results

The sequencing of 200 clones from the forward (individuals incubated at 10°C versus individuals exposed at 20°C) SSH library, as well as from the reverse library (individuals incubated at 20°C versus individuals exposed at 10°C) allowed the identification of 83 different gene sequences for *P. pandorae* (20% of the whole sequencing effort: the other sequences are redundant). Table 2 shows the unique sequences obtained in both SSH libraries with the best E-values for the sequences when identified after Blast analysis. The identified sequences indicate that temperature regulates the expression of genes involved in various cell functions: (1) respiratory chain (9.6%), (2) sulfur and oxygen transport (8.4%), (3) metabolism (2.4%), (4) cell communication, membrane receptors and immune system (4.8%), (5) cytoskeleton production and maintenance (3.6%), (6) ribosomal proteins (8.4%), (7) proteins of unknown function (2.4%) and (8) unknown sequences (60.4%). Interestingly, most transcripts identified in the reverse subtractive library cor-

respond to genes involved in the respiratory chain and in oxygen transport. A high proportion of unknown sequences were also obtained limiting the identification of new temperature-regulated putative candidate gene.

Relative expression level of some target genes

Levels of expression of 20 transcripts obtained in the *B. thermophilus* library and 14 transcripts identified in the *P. pandorae* library were quantified from pools of individuals of each thermal condition by using real-time PCR. For all the genes, we showed that the differential expression observed for the genes corresponds to the library they came from. In addition, stronger differential expressions were observed in *B. thermophilus* for metabolic genes such as S-adenosylhomocysteinase-hydrolase, arginine kinase or $\Delta 5$ -desaturase, and some genes such as secreted protein acidic rich in cystein (SPARC, a basal membrane component), elongation factor beta, actin and proteins related to mussel mobility (foot protein and adhesive plaque matrix protein) (Table 3). While it was identified in the reverse library, the gene encoding the kaliculidine did not display a clear differential expression between the two experimental conditions. In *P. pandorae*, differences between levels of expression of the two sets of experimented individuals were higher than those observed in mussels. Some genes also showed a very high level of expression in samples exposed to 10°C as compared to those exposed at 20°C, especially for the secreted nidogen domain protein, Rab7, the intracellular hemoglobin and the ribosomal protein S16 that were strongly inhibited in annelids exposed to 20°C. Almost all genes encoding subunits of hemoglobins (except linker L2) displayed a coherent expression pattern of up-regulation in annelids exposed to 20°C (Table 4).

Discussion

In the present paper, we described and analyzed gene expression in two hydrothermal species in response to two temperatures, one included in the range of temperature encountered by both species (10°C), and the other near (for worms) or above (for mussels) their thermal limit range (20°C). The following parts of the discussion deal with the respective responses of these two species in terms of differences and common features typifying their ability to adapt different thermal regimes.

Unidentified sequences

Over 50% of the sequences for each species could not be identified based on homologies. This may be due to the limited amount of data available for invertebrates, or to the SSH protocol itself that requires the use of a restriction enzyme, possibly leaving only the UTRs for cloning. This may explain that we did not obtain typical heat stress proteins such as the inducible heat shock protein 70 (HSP70). This could also be due to the long term acclima-

Table 1: Regulated genes identified in the SSH libraries of thermally challenged *Bathymodiolus thermophilus* with significant database matches (only sequences with E-value above 0.005 are shown).

Homolog (protein); Blastx value	Homolog species	Insert size (bp)	Accession number	SSH library
Cell cycle, DNA repair, protein regulation and transcription				
Elongation factor I alpha; 1e-14	<i>Mytilus galloprovincialis</i>	428	GHI96568	forward
Elongation factor beta; 2e-05	<i>Strongylocentrotus purpuratus</i>	153	GHI96577	forward
Elongation factor-2; 9e-89	<i>Hutchinsoniella macracantha</i>	577	GHI96570	forward
S-cyclophilin; 2e-14	<i>Gallus gallus</i>	384	GHI96575	forward
H3 histone; 4e-43	<i>Mus musculus</i>	538	GHI96576	reverse
Myc homolog; 7e-10	<i>Crassostrea virginica</i>	704	GHI96563	reverse
Respiratory chain				
Cytochrome c oxidase subunit II; 5e-32	<i>Lampsilis ornata</i>	424	GHI96573	reverse
Cytochrome c oxidase subunit III; 1e-20	<i>Arbacia lixula</i>	260	GHI96584	reverse
ATP synthase; 3e-21	<i>Bos taurus</i>	415	GHI96588	reverse
Metabolism				
S-adenosylhomocysteine hydrolase 2e-60	<i>Branchiostoma belcheri tsingtaunense</i>	505	GHI96574	forward
Cytosolic Lactate/malate dehydrogenase; 2e-14	<i>Caenorhabditis elegans</i>	421	GHI96590	forward
Arginine kinase;	<i>Octopus vulgaris</i>	135	GHI96579	forward
Anhydrase carbonic 2; 2e-02	<i>Oncorhynchus mykiss</i>	114	GHI96567	forward
Delta-5-desaturase; 4e-25	<i>Strongylocentrotus purpuratus</i>	195	GHI96582	forward
Stress response and detoxication				
Glutathione peroxidase; 0.001	<i>Bombyx mori</i>	101	GHI96562	forward
Heat Shock Protein 90; 4e-21	<i>Tetraodon nigroviridis</i>	156	GHI96561	forward
Cell communication, membrane receptors, immune system				
Kallicludine I; 3e-16	<i>Anemonia sulcata</i>	406	GHI96560	reverse
Techylectin-5A; 8e-29	<i>Tachypleus tridentatus</i>	337	GHI96558	reverse
Electron-transfer-flavoprotein; 4e-57	<i>Strongylocentrotus purpuratus</i>	535	GHI96557	reverse
ItmI protein, 4e-27	<i>Xenopus laevis</i>	179	GHI96556	reverse
Clq-like adipose specific protein; 2e-07	<i>Salvelinus fontinalis</i>	570	GHI96594	reverse

Table 1: Regulated genes identified in the SSH libraries of thermally challenged *Bathymodiolus thermophilus* with significant database matches (only sequences with E-value above 0.005 are shown). (Continued)

β -1,3-N-acetylglucosaminyltransferase 6; 5e-48	<i>Homo sapiens</i>	586	GHI96571	reverse
Secreted protein, acidic, rich in cysteine SPARC; 2e-20	<i>Artemia franciscana</i>	523	GHI96580	forward
Defensin; 1e-04	<i>Haliotis discus hannai</i>	105	GHI96559	forward
Cytoskeleton production and maintenance				
α -2-tubulin; 4e-27	<i>Gecarcinus lateralis</i>	348	GHI96585	forward
Actin; 1e-35	<i>Mytilus galloprovincialis</i>	541	GHI96578	forward
Adhesive plaque matrix protein; 6e-06	<i>Mytilus galloprovincialis</i>	291	GHI96589	forward
Foot protein 2; 1e-6	<i>Mytilus edulis</i>	108	GHI96581	forward
Pedal retractor muscle myosin; 6e-50	<i>Mytilus galloprovincialis</i>	426	GHI96564	forward
Hemicentin; fibulin 6; 2e-14	<i>Rattus norvegicus</i>	304	GHI96587	reverse
Ribosomal proteins				
Ribosomal protein L3; 2e-55	<i>Argopecten irradians</i>	495	GHI96591	forward
Ribosomal protein L4; 4e-80	<i>Latimeria chalumnae</i>	926	GHI96586	forward
Ribosomal protein S14; 3e-04	<i>Rattus norvegicus</i>	256	GHI96555	forward
Ribosomal protein S15; 1e-64	<i>Argopecten irradians</i>	465	GHI96566	forward
Ribosomal protein S19; 2e-09	<i>Chlamys farreri</i>	362	GHI96569	forward
Ribosomal protein S25; 1e-28	<i>Crassostrea gigas</i>	379	GHI96583	forward
Ribosomal protein L15; 7e-16	<i>Siniperca kneri</i>	214	GHI96565	reverse
QM protein; 7e-11	<i>Apis mellifera</i>	622	GHI96572	reverse
Unknown function				
CG33171-PC, isoform C; 4e-07	<i>Drosophila melanogaster</i>	615	GHI96592	reverse
Repeat organellar protein-related; 1e-05	<i>Plasmodium yoelii yoelii</i>	765	GHI96595	reverse
Unidentified sequences				
39 sequences			GHI96596 to GHI96632	

Forward: genes over-expressed at 10°C versus 20°C;
Reverse: genes over-expressed at 20°C versus 10°C.

tion (about 2 days) offered to the animals leading to an attenuation of the stress machinery with time. None of the SSH libraries (mussel and annelid) indeed contained mRNA coding for any HSP70s, that are involved in the protection of other proteins from denaturation caused by a variety of stressors [32,33]. Available data dealing with

HSP70 in hydrothermal species were only obtained from polychaetes, mussels and shrimps exposed to brief heat shocks [21,22,34]. In previous studies, a positive correlation between the levels of DNA strand breakage and HSP70 expression in response to decompression stress were also found by Pruski and Dixon [35]. In the shrimp

Table 2: Regulated genes identified in the SSH libraries of thermal exposed *Paralvinella pandorae irlandei* with significant database matches (E-value above 0.005).

Homolog (protein); Blastx value	Homolog Species	Insert size (bp)	Accession number	SSH library
Respiratory chain				
Cytochrome c oxidase subunit I; 1e-73	<i>Littorina saxatilis</i>	772	GHI96478	reverse
Cytochrome oxidase subunit II; 1e-46	<i>Lumbricus terrestris</i>	715	GHI96499	reverse
Cytochrome c oxidase subunit III; 2e-34	<i>Albula glossodonta</i>	457	GHI96480	reverse
Cytochrome c oxidase polypeptide Va; 1e-20	<i>Strongylocentrotus purpuratus</i>	149	GHI96490	reverse
Ubiquinol-cytochrome c reductase complex; 4e-23	<i>Mus musculus</i>	537	GHI96502	reverse
Cytochrome b; 1e-23	<i>Urechis caupo</i>	258	GHI96495	reverse
NADH dehydrogenase subunit I; 4e-48	<i>Urechis caupo</i>	571	GHI96493	reverse
NADH dehydrogenase subunit 6; 7e-05	<i>Clymenella torquata</i>	190	GHI96501	reverse
Sulfur and oxygen transport				
Hemoglobin A2c chain; 3e-31	<i>Arenicola marina</i>	569	GHI96487	reverse
Hemoglobin B1 chain precursor; 4e-21	<i>Arenicola marina</i>	326	GHI96479	reverse
Hemoglobin B2 chain; 1e-12	<i>Arenicola marina</i>	516	GHI96491	reverse
Hemoglobin linker LY precursor; 5e-86	<i>Riftia pachyptila</i>	529	GHI96505	reverse
Extracellular globin linker L1 precursor; 0.002	<i>Alvinella pompejana</i>	335	GHI96504	reverse
Hemoglobin linker L2 precursor; 0.005	<i>Alvinella pompejana</i>	727	GHI96483	forward
Intracellular hemoglobin; 5e-41	<i>Alvinella pompejana</i>	547	GHI96492	forward
Metabolism				
Trypsin; 6e-60	<i>Litopenaeus vannamei</i>	661	GHI96494	reverse
S-adenosylhomocysteine hydrolase; 2e-66	<i>Branchiostoma v</i>	529	GHI96476	forward
Cell communication, membrane receptors, immune system				
Lipid binding protein 9; 5e-05	<i>Caenorhabditis elegans</i>	463	GHI96481	forward
GTP-binding protein (rab7); 7e-89	<i>Canis familiaris</i>	641	GHI96489	forward
Xylan endohydrolase isoenzyme, 6e-22	<i>Arabidopsis thaliana</i>	455	GHI96472	reverse
Cyclophilin B; 4e-14	<i>Gallus gallus</i>	384	GHI96477	reverse
Cytoskeleton production and maintenance				
Chymotrysin; 4e-11	<i>Lumbricus rubellus</i>	230	GHI96484	forward

Table 2: Regulated genes identified in the SSH libraries of thermal exposed *Paralvinella pandorae irlandei* with significant database matches (E-value above 0.005). (Continued)

Secreted nidogen domain protein; 4e-08	<i>Strongylocentrotus purpuratus</i>	325	GHI96473	forward
Actin A1; 2e-35	<i>Haliotis iris</i>	541	GHI96475	forward
Ribosomal proteins				
Ribosomal protein L5; 3e-58	<i>Rattus norvegicus</i>	423	GHI96497	reverse
Ribosomal protein L28; 6e-07	<i>Haliotis asinina</i>	262	GHI96486	reverse
Ribosomal protein L24; 1e-38	<i>Danio rerio</i>	439	GHI96482	forward
Ribosomal protein S3; 3e-30	<i>Crassostrea gigas</i>	205	GHI96474	forward
Ribosomal protein S16; 2e-41	<i>Gallus gallus</i>	337	GHI96500	forward
Ribosomal protein SA; 8e-43	<i>Xenopus tropicalis</i>	514	GHI96496	forward
Ribosomal protein P1; 2e-11	<i>Drosophila yakuba</i>	444	GHI96498	forward
Unknown function				
CG14235-PA, isoform A; 1e-27	<i>Tribolium castaneum</i>	483	GHI96488	forward
CBG19860; 2e-08	<i>Caenorhabditis briggsae</i>	266	GHI96485	reverse
Unidentified sequences				
50 sequences			GHI96507 to GHI96554	

Forward: genes over-expressed at 10°C versus 20°C; Reverse: genes over-expressed at 20°C versus 10°C.

Rimicaris exoculata, regulation of HSP70 in response to temperature was detected at the protein level [21], but no data related to HSP70 gene regulation is available so far.

Is P. pandorae better adapted to higher temperature than B. thermophilus?

Metabolic adjustments in response to thermal challenges are essential for aquatic ectotherms, whose body temperature fluctuates over the full range of temperature in their habitat [36]. Temperature can also influence metabolic regulation, eliciting transition to anaerobiosis even in oxygenated waters [37]. A high thermal sensitivity of metabolism over the environmental range is associated to increased long-term metabolic costs and with a lower tolerance to extreme temperatures. Classically, lowering metabolic rate and, thus energy saving is also considered as one of the most important adaptations for hypoxia endurance [38,39]. Adaptation to these conditions has resulted in reduced growth rates, as well as reduced development and metabolism [40]. Previous studies conducted on the marine gastropod *Littorina saxatilis* showed that an acute long-term temperature increase could disturb metabolism, leading to progressive metabolic depression and adverse changes in the cellular energy status due to its

transition to partial anaerobiosis [41]. Antarctic marine species are also much less capable to survive elevated temperatures [42] and calculated temperature envelopes for these organisms were 2–4 times smaller than those for temperate species [43]. Studies on the Mediterranean mussel *Mytilus galloprovincialis* showed that a long acclimation of up to 30 days at high temperatures (18 to 30°C) leads to behavioral (increase of duration of valve closure), metabolic (metabolic depression with a shift from aerobic to anaerobic metabolism) and molecular (increase in HSPs protein levels) responses [44]. Interestingly, the deep-sea mussel *B. thermophilus* seems to share some general features with organisms living in polar oceans that are characterized by very stable low temperatures, below 5°C. We first hypothesized that annelids were able to cope with a larger range of temperature compared to the mussels, because they were described as early colonizers of new chimneys at hydrothermal vents [45], and thus able to sustain highest temperatures, at least over short periods of time. While few genes seem to be regulated similarly between the two species (actin, mitochondrial cytochrome oxidase, S-adenosylhomocysteine hydrolase), most cell functions are very dissimilar, suggesting a different response of these organisms to temper-

Table 3: mRNA expression of genes in *B. thermophilus* presented as a fold-change value with animals exposed at 10°C as a calibrator.

Genes	Fold-change (calibration to 10°C)
Down-regulated at 20°C	
Defensin	1192.69
Elongation Factor beta	4.59
Ribosomal protein L3	6.23
Foot protein	3.36
Pedal retractor muscle myosin	5.17
Adhesive plaque matrix protein	2.83
Actin	4.03
Secreted Protein, Acidic, Rich in Cystein (SPARC)	7.21
S-adenosylhomocysteine hydrolase	5.24
Cyclofilin S	4.35
Δ5-desaturase	8.28
Carbonic anhydrase 2	11.39
Adenylate kinase	6.59
Glutathione peroxidase	6.28
Cytosolic malate dehydrogenase	8.75
HSP90	6.23
BthermEST1	2.57
Up-regulated at 20°C	
Kalcludine	1.06
Myc homolog	12.30
Techlectin 5A	6.63

ature. The genes that were identified in the mussel libraries could indicate that this species tends to react as a stenoeicious species rather than an euryeicious species (as could be expected for organisms living in a highly fluctuating environment). A general depression is indeed observed in expression of *Bathymodiolus* genes involved in transcription/translation, mobility, energetic metabolism, and oxidative stress in response to temperature increase. Conversely, genes involved in cell disorder and immune system (ie myc-homolog, kalcludin...) are up-regulated at

the highest temperature. No similar pattern is observed in *P. pandorae*, leading to the hypothesis that this species could better adapt to high temperatures.

A long exposure at a temperature of 20°C (43 hours) clearly appears to be a thermal physiological limit for *B. thermophilus* that lives in colder habitats. As mussels encounter short pulses of hot water under *in situ* conditions, it seems that the duration of the heat exposure is critical and probably more important than the tempera-

Table 4: mRNA expression of genes in *P. pandorae* presented as a fold-change value with animals exposed at 10°C as a calibrator.

Genes	Fold-change (calibration to 10°C)
Up-regulated at 20°C	
Hemoglobin A2c	250
Hemoglobin B2	31.25
Linker L1	142.86
PpandEST2	13.33
Xylan endohydrolase	5.13
Down-regulated at 20°C	
Intracellular Hemoglobin	1226.22
Linker L2	4.03
Secreted Nidogen domain protein	1278.29
S-adenosylhomocysteine hydrolase	35.51
Chymotrypsinogen	20.46
PpandEST1	1.53
Ribosomal protein S16	1640.59
Lipid binding protein	1.80
Rab 7	2469.49

ture value itself. According to the experimental conditions used for this study, it remains difficult to evaluate how long *B. thermophilus* is really able to withstand a temperature of 20°C without severe physiological damage. Complementary experiments such as the determination of differential mortality kinetic in longer-term exposures and at different ranges of temperatures have to be performed in further studies to better understand the thermal resistance/response/adaptation of this species. As all individuals survived for about 2 days (43-hour experiment), *B. thermophilus* can deal with thermal stress for at least few days but it is not clear whether they can then recover from such stress.

The hypothesized limited adaptation of *B. thermophilus* to high temperature is also supported by a decrease of expression of nearly all genes identified in the mussels exposed to 20°C, a pattern indicative of global metabolic depression. Among these genes, there were many ribosomal proteins and some elongation factors, indicating that the protein synthesis pathway was clearly involved in response to temperature. However, different ribosomal proteins are identified in both reverse and forward libraries showing a complex regulatory process (or the absence of regulation) in intra-molecular interactions in ribosomes. This result is commonly observed in transcriptomic studies in mollusks in response to various environmental parameters [46-49]. A higher number of ribosomal proteins were however identified in the mussel forward libraries suggesting a possible metabolic depression in samples exposed to 20°C when compared to those exposed at 10°C. In *P. pandorae* libraries, similar pattern of down regulation at 20°C of 5 ribosomal proteins among the 7 identified is observed.

More specifically, it is noteworthy that several genes of the mussel energetic pathways were down-regulated. Among them, arginine kinase (ArgK) and cytosolic malate dehydrogenase (cMDH) are found to be down regulated in mussels exposed to 20°C. ArgK catalyzes the transfer of phosphate between ATP and arginine (arginine phosphate + MgADP + H⁺ ↔ arginine + MgATP²⁻), and plays a critical role in cellular energy metabolism in invertebrates [50]. It also serves as an energy reserve because it can readily transfer phosphor-arginine to ATP when energy is needed [51,52]. However, it was never found associated with thermal stress. To our knowledge, no data on the thermal regulation of mRNA expression of ArgK has been reported to date. Its regulation has mostly been studied at a protein level and this is the only phosphagen kinase known in crustaceans and mollusks. ArgK is indeed regulated in crustaceans and mollusks under hypoxia [53,54]. In the crustacean *Marsupenaeus japonicus*, the up-regulation of ArgK under hypoxia may represent a provision for oxygen recovery after a short period of hypoxia [54]. The

second metabolic enzyme is the cytosolic malate dehydrogenase, which catalyzes the dehydrogenation of malate (malate + NADP⁺ ↔ oxaloacetate + NADPH + H⁺). It plays a major role in a number of metabolic pathways, including the malate-aspartate (or NADH) shuttle and the acetate shuttle active in lipogenesis, amino acid synthesis and gluconeogenesis. cMDH is an interesting candidate gene to study adaptation to temperature as this enzyme showed differences in the effects of temperature on kinetic properties in shallow water species [7,55,56]. Even though no cMDH mRNA expression has been reported in these studies, differences in protein properties strongly suggest a clear involvement of this key gene in response to temperature. The down-regulation of both ArgK and cMDH in *B. thermophilus* could lead to a decrease of mitochondrial respiration, leading to a lower ATP production, and resulting in the establishment of a global metabolic depression in response to temperature.

The cDNA coding for a HSP90 was found in the mussel SSH libraries. However, HSP90 displayed a down-regulation at 20°C when compared to 10°C suggesting that the process of protein re-naturation was probably over. HSP90 proteins have key roles in signal transduction, protein folding, protein degradation, and morphological evolution [57-59]. HSP90 is up-regulated in response to heat stress in *Drosophila subobscura* [60], the whitefly *Bemisia argentifoli* [61], and the flesh fly, *Sarcophaga crassipalpis* [62]. It is induced by thermal stress in the Goby fish but could also decrease in expression to a normal level during the acclimatization process [63]. In *M. galloprovincialis*, both HSP70 and HSP90 protein expression were shown to increase in response to long-term thermal challenge [44].

Does temperature generate a stronger oxidative stress in mussels than in annelids?

We identified several genes that are classically expressed in response to oxidative stress in the mussel libraries but not in the annelid libraries, suggesting a differential behavior of both species. Two main hypotheses can explain the presence of an oxidative stress: (1) a direct effect of temperature changes on lipid composition or (2) variations of the oxygen concentration during experiments. In the first hypothesis, temperature directly affects cells by modifying membrane composition through replacement of unsaturated fatty acids at low temperatures towards saturated fatty acids at high temperatures [64], and secondly by inducing apoptosis *via* activation of the sphingomyelin pathway that leads to the process of lipid peroxidation [65]. Many biological structures, such as enzymes and lipid bilayer membranes, depend on a particular degree of molecular instability or fluidity, which is directly affected by temperature. In the particular case of our experimented hydrothermal species, lipids of cell membrane bilayers must be both fluid and structurally coherent to form a

functional membrane, a characteristic very sensitive to temperature change [2]. Lipid peroxyl radicals (LPO) are the result of a reaction between lipid and oxygen and are known to damage cells by changing the fluidity and permeability of the membrane and/or by directly damaging DNA and other intracellular molecules, such as proteins [66]. As a consequence of the lipid peroxidation process, superoxide anion radicals can be produced. Lipid peroxidation has been studied in hydrothermal vent mussels and high levels of LPO were detected in *B. azoricus* in response to a strong effect of environmental heavy metal concentrations [67]. Recently, heavy metal stresses, such as copper exposure, or changes in hydrostatic pressure were also shown to produce LPO in *B. azoricus* [68]. Fatty acid desaturases are very important during the process of fatty acid metabolism that contributes to the structural and functional maintenance of biological membranes in living organisms. The down-regulation of $\Delta 5$ -desaturase mRNA expression of *B. thermophilus* exposed to 20°C is consistent with a modification of membrane lipid content. We also identified a gene encoding SPARC, which is more expressed in mussels incubated at 10°C when compared to those exposed to 20°C. SPARC is classically known to modulate cellular interaction with the extracellular matrix through interactions with proteins such as laminins and collagen [69,70]. SPARC was also shown to be up-regulated in response to heat-shock and other stresses [71,72]. SPARC also possesses a chaperone-like activity *in vitro* suggesting its involvement in stress response [73]. Its down-regulation at 20°C is coherent with results observed by previous authors and could reflect a strong disorder in membrane composition due to the high temperature.

In the second hypothesis, the generation of reactive oxygen species (ROS) as side products of electron transfer during aerobic metabolism [74] can explain the regulation of genes encoding protective proteins in mussels. Here, oxidative stress can be due to the experimental conditions used where sea-water was at a low-oxygen concentration (below 120 μ M) associated with a consumption by animals and the effect of temperature. In the presence of low oxygen concentration or anoxic conditions, organisms use anaerobic metabolism and annelids and mollusks are able to use more efficient mitochondrial pathways of fermentation [75-77]. Under normal physiological conditions, anaerobic metabolism produces free radicals, and cells tend to maintain a balance between generation and neutralization of ROS. When organisms are subjected to xenobiotics, temperature increase or anoxia events, the generation of ROS can exceed the scavenging capacity [78]. All organisms possess their own cellular antioxidant defense system, composed of both enzymatic (superoxide dismutase, catalase and glutathione peroxidases) and non-enzymatic (glutathione, vita-

mins...) components. Glutathione peroxidases (GPx), that have protective roles against oxidative stress, have been identified in *B. thermophilus* libraries suggesting an oxidative stress as a direct or indirect result of temperature challenge. Surprisingly, GPx expression is lower in mussels exposed to 20°C than those exposed to 10°C despite the fact that oxidative stress is supposed to be stronger at 20°C, supporting the idea that *B. thermophilus* is no longer able to regulate expression of oxidative stress related genes. We also identified a gene encoding a myc homolog which is up-regulated at 20°C compared to 10°C-exposed mussels. This protein belongs to a transcription factor family and is involved in the cell division control. Myc and its binding partners regulate the expression of a large number of genes that regulate diverse functions, including protein synthesis, apoptosis, and DNA and energy metabolism [79-81]. Generally speaking, over-expression of a myc-homolog enhances apoptosis by acting as a transcription repressor [82,83]. In bivalves, c-myc has previously been shown to be up-regulated by hypoxia [49] and hydrocarbon stresses [46]. Identification of c-myc in mussel exposed to 20°C seems to be indicative of the very poor biological condition of the 20°C-exposed individuals and thus in accordance with the hypothesis of a low tolerance of *B. thermophilus* to extended exposure to high temperature.

A gene that is involved in adenosine metabolism, and that has previously been shown to be regulated in response to hypoxia, has also been found in both mussel and annelid libraries. This enzyme called S-adenosylhomocysteinase hydrolase (SAHH, EC 3.3.1.1) catalyses the reversible hydrolysis of S-adenosylhomocysteine to form homocysteine and adenosine [84]. AdenosineMethionine/AdenosineHomocysteine turnover is believed to play a critical role in methionine metabolism and the regulation of biological methylation processes. Tissue hypoxia induces a variety of functional changes, including enhanced transcriptional activity associated with high transmethylation activity (e.g. mRNA cap methylation) in the nucleus. Disturbance in DNA methylation pattern has previously been observed in response to various stressors, such as heavy metals, as a consequence of toxicity [85,86]. In both our species, the mRNA expression of this gene is lower in animals exposed to 20°C than to 10°C. This regulation of SAHH mRNA expression supports the hypothesis of a response to a direct or indirect oxidative stress. Presence of SAHH in response to temperature also illustrates the importance of methylation processes as a response to temperature increase. Generally speaking, a strong DNA methylation leads to a decrease or an inactivation of gene expression. Methylation processes regulation could be an interesting type of response to temperature in hydrothermal species.

Specific responses to temperature in mussels and annelids

In the mussel libraries, we interestingly identified three down-regulated genes at 20°C that are involved in foot activity (foot protein and pedal retractor muscle myosin) and byssus activity (adhesive plaque matrix protein). These results are in sharp contrast with previous studies performed on the brackish-water mussel, *Mytilopsis leucophaeata* that showed an increase in foot activity index and byssus thread production in response to thermal challenge [87]. Authors demonstrated that both foot activity and byssus production were higher when temperature increased from 4 to 20°C, remained stable between 20 and 28°C, and then strongly decreased beyond 28°C. This species commonly lives in a range of temperatures comprised between 4°C (winter) and 20°C (summer). Mussels of the genus *Bathymodiolus* are able to change location when the conditions are not adequate [88]. This can be viewed as an escape response in the presence of stress factors. They also probably use this mobility to optimize their position in the hydrothermal fluid in order to acquire the sulfide (and/or methane for some species) they need to feed their symbionts. The decrease of the expression of genes encoding proteins related to mobility in mussels exposed at 20°C, again reflects the poor physiological condition of these individuals, since mussels usually live in colder waters (4 to 14°C).

In the annelid SSH libraries, we identified several genes encoding various extracellular globin chains (B1, A2 and B2), one intracellular globin and also three linkers called linker L1, linker L2 and linker LY. These results illustrate a strong involvement of respiratory pigment in general and in particular of the hexagonal bilayer hemoglobin (HBL-Hb) in response to temperature in this species. In Alvinellidae, respiratory gas transport is performed by the blood and the coelomic fluid, and three main types of globins are present: non-circulating in the cytoplasm, circulating and intracellular in the coelom, and extracellular in the vascular system [24]. Earlier work reported the temperature effect on both the function and the stability of Hbs in the annelid *Alvinella pompejana*, under atmospheric pressure and for temperatures ranging from 10°C to 40°C. These Hbs are able to maintain a capacity to reversibly bind oxygen *in vitro* over this range. At 50°C, the Hbs are oxidized and aggregated during the de-oxygenation and the re-oxygenation [89]. These results are in agreement with the hypothesis that annelids, even if they are able to withstand a strong thermal stress [25], are nonetheless unable to sustain high temperature for a long time [24]. Because *P. pandorae* lives in a relatively cold environment compared to other *Paralvinella* species, such as *P. sulficola*, and probably do not experience very high temperatures, the involvement of Hbs in temperature response could be the result of several processes and not only driven by Hb thermostability properties. We observed a strong increase of mRNA encoding extracellular Hb sub-

nits in individuals exposed to 20°C and conversely, a decrease of intracellular subunit mRNA expression. It has been suggested that extracellular Hbs were preferentially involved in oxygen uptake and transport, while intracellular Hbs acted as an oxygen reserve for the worm and potentially returned oxygen to the extracellular Hb [90]. Because oxygen availability decreases with temperature, *Paralvinella* increased their extracellular Hb production to optimize the oxygen uptake and transport. At the same time the intracellular Hb was down-regulated, possibly to avoid the release of O₂ to the tissue since the worms experienced hot temperature but not hypoxia). We also observed an opposite regulation in the expression of linkers L1 and L2 in *P. pandorae* in response to thermal stress suggesting a possible rearrangement of the linker composition of the HBL-Hb molecule under temperature-induced oxidative stress. Very few studies have dealt with the linker function and regulation at a transcriptional level. In the Earthworm *Lumbricus terrestris*, linkers have been shown to exhibit a superoxide dismutase activity conferring a protection against superoxide ions for HBL-Hb molecules [91]. Linkers of the thermally-stressed alvinellids may therefore have been mobilized as an active defense against newly-produced ROS.

In the *P. pandorae* library, we also identified one gene encoding a secreted nidogen domain protein that showed a strong down-regulation at 20°C. Secreted nidogen domain protein, also known as entactin, belongs to basement membrane proteins. These membranes are made of type IV collagens and laminins, both of which exist as various isoforms in animals [92,93]. These proteins are cell-adhesive and form networks that confer mechanical stability to the basement membranes. Other ubiquitous basement membrane components are the proteoglycan perlecan and nidogen/entactin. Previous *in vitro* experiments showed that recombinant nidogen-1 interacted through different binding sites with the three main basement membrane components (laminin, collagen IV, and perlecan), and mediated the formation of ternary complexes between laminin and collagen IV [94]. These results therefore suggest that secreted nidogen domain protein is a key component of alvinellid basement membranes assembly, connecting the laminin and collagen networks, and integrating other basement membrane components as previously reported by Timpl and Brown [93]. We suggest that temperature (and/or pressure) above normal could induce strong changes in the membrane composition of the worms and therefore increase interactions between secreted nidogen domain protein, collagen and other membrane protein in order to readjust porosity/permeability. Other proteins that are thought to partially play a role in membrane component modeling have also been characterized. We identified a Ras-associated binding 7 (Rab 7) protein belonging to the Rab family. These are small GTPases of the Ras superfamily that con-

tinuously cycle between the cytosol and different membranes. The Rab family appears to be essential for the regulation of intracellular membrane traffic in mammalian cells. Rab proteins are anchored to the cytoplasmic surface of specific intracellular membrane compartments via the geranyl-geranyl group that is post-translationally added to the C-terminal cysteines and is important for their function [95]. Each Rab protein regulates one (or more) specific step of intracellular membrane traffic in eukaryotic cells, probably by assembling the general tethering/docking/fusion machinery [96]. Moreover, several lines of evidence suggest an involvement of Rab proteins in actin- and microtubule- based processes [97]. Rab7, a member of the Rab family small G proteins, has been shown to regulate intracellular vesicle traffic to late endo-

some/lysosome and lysosome biogenesis, but the exact roles of Rab7 are still undetermined [98,99]. Accumulating evidence suggests that each Rab protein has multiple target proteins that function in the exocytic/endocytic pathway. Because no studies showing how temperature could affect Rab 7 expression, its down regulation observed in *P. pandorae* by temperature remains difficult to explain but could be associated with results observed for nidogen protein.

Conclusion

Our results indicate that the mussels and the worms did not cope with temperature in the same way. While the mussel *B. thermophilus* seems to show a general metabolic depression (strong decrease of mRNA expression for

Table 5: Primer sequences used in real-time PCR expression analysis for *B. thermophilus* SSH libraries validation

Genes	Primer sequences
Elongation factor beta	For: 5' GATCTTAAAAGTAAAGCTGGTCAGCAAGC 3' Rev: 5' AACAAATCAAATCATCATCACCGCC 3'
Ribosomal protein L3	For: 5' AGATATATCGTATTGGAGAGGGATACACACC 3' Rev: 5' GCTGTTGCATCCTTCTTCAATGGACCCAT 3'
Foot protein	For: 5' AATAATGGTAAATGTGTTGCTAATGGCTA 3' Rev: 5' CCGTATCCCCTTCTACAACATCTACCGCC 3'
Pedal retractor myosin	For: 5' AGAACCGACGAATTGGAAGAGGCCAAGAG 3' Rev: 5' AACAAATCAGCAGAGTAACTGCGGGCCTC 3'
Adhesive plaque matrix	For: 5' AAAAGATGTGAAGTAAACAGATGCAGCCCA 3' Rev: 5' CCGTATCCCCTTCTACAACATCTGCCGCC 3'
Actin	For: 5' ACGCGGGTCAGGGTCGGACGTAGCCACGC 3' Rev: 5' ATGGAGATCAGACGGAGATGGTCCTCCTC 3'
Adenosylhomocysteinase	For: 5' GTAAATCTTGTTGTGCTCATGGTCATCC 3' Rev: 5' GATTTGAATGGTCCTTCTTTAGGTAGACC 3'
Cyclophilin S	For: 5' TTGAATAAAGCCAGATGGATGGATGGAAA 3' Rev: 5' AAATCTTCATCGCTAGCTGCTTGTGCTTC 3'
Arginine kinase	For: 5' ATGGGTGAAGTAGCAGAATTGTGGGCTAA 3' Rev: 5' TCACATGCATACAATCCGACTCCGCT 3'
Glutathione peroxidase	For: 5' ATGGGCATAAACTTGGGAGACATTTT 3' Rev: 5' CCTAATTCTGTTGTACAAACAGGGGTATA 3'
BthermEST1	For: 5' AGTGACTTCACAACCTGCCCGTATGTGGAA 3' Rev: 5' TTGGTGCACATCATCAAGAAGGAGAGTAT 3'
Kallicludine	For: 5' CCATGCAATGAAGATTGTCTTTTGCCAAA 3' Rev: 5' TTTGGCAAAAAGACAATCTTCATTGCATGG 3'
Defensin	For: 5' ATGTTCAAAGTAACTTTGTTCTTCGTTGG 3' Rev: 5' TACGATCTGCAGTGACACACGTTATG 3'
SPARC	For: 5' AACGCAGACGACCACCGTACAGACGC 3' Rev: 5' TATGCATCACACTTGTCTGTAATGTCAACC 3'
$\Delta 5$ -desaturase	For: 5' AACGACTGGTTTACAGGGCATCTAAA 3' Rev: 5' TTTAGATGCCCTGTAAACCAGTCGTT 3'
Carbonic anhydrase-2	For: 5' GATGACAAGGAAGGATCTGAGCACACTCT 3' Rev: 5' AGAGTGTGCTCAGATCCTTCTTGTGTCATC 3'
HSP90	For: 5' ATGCCTGAACCTGAAACAACATATGGATGA 3' Rev: 5' GAATACATCTGGGAATCTGCAGCTGGTGG 3'
Myc homolog	For: 5' TCTGTTTATGATGCCTGGGTCACTCC 3' Rev: 5' GGAGTGACCCAGGCATCATAAACAGA 3'
Cytosolic malate dehydrogenase	For: 5' ATGGCAGTTCCTTCAGATGGATCTTA 3' Rev: 5' CAATACACAAAAACAGACACTGTATACAT 3'
Techylectin 5A	For: 5' GGATATCAGGGTAATGCAGGAGATGC 3' Rev: 5' GCATCTCCTGCATTACCCTGATATCC 3'

numerous genes), possibly due to maladaptation and cell disorders when temperature increased, the annelid *P. pandorae* mainly displayed a strong regulation of the mRNA encoding subunits and linkers of respiratory pigments and some proteins involved in membrane constitution. In both cases, these regulations seem to be partly due to a possible cellular oxidative stress induced by temperature increase (10°C to 20°C). The large number of unknown sequences makes definitive conclusions difficult. The data collected may contain a number of candidate genes regulated by temperature that require annotation and functional characterization before meaningful interpretation of temperature adaptation of both *P. pandorae* and *B. thermophilus* is possible. New tools, such as microarrays will help evaluate the expression and characterization of these genes. This work will serve as a starting point for studying the transcriptomic response of hydrothermal mussels and annelids in future experiments in response to thermal stress at various conditions of exposure duration and thermal level.

Methods

Animal sampling and treatment

Specimens of both the hydrothermal vent mussel *Bathymodiolus thermophilus* (67.8 ± 30 mm in length) and the

annelid worm *Paralvinella pandorae irlandei* (≤ 10 mm in length) were collected together from the Oasis site (17° 25.42S, 113° 12.28W) on the East Pacific Rise during the BIOSPEEDO cruise [100] by using the telemanipulated arm of the submersible Nautille. The individuals were collected in a cold zone associated with weak diffuse flow (< 10°C). Once on board the ship, animals were immediately transferred from the insulated collection basket to the pressurized aquaria IPOCAMP™ (Incubateur Pressurisé pour l'Observation et la Culture d'Animaux Marins Profonds [101]) under an *in situ* pressure of 260 bars. Indeed, several studies showed that experiments conducted on mussels at atmospheric pressure lead to a global increase of stress parameters (lipid peroxidation, anti-oxidant enzymes activity, DNA damage) and to a limitation of stress response capacity [14,23,102]. One group of 5 mussels and 20 worms was placed at 10°C, and another group at 20°C, inside two different pressure vessels for 43 hours. The pressure vessels were operated in a flow-through mode (20 L/hour) for the first 3 hours, and then isolated. Every 15 hours, seawater was re-circulated for 3 hours, until the end of the incubation. When in flow-through configuration, the pressure aquaria were fed with sea-water aerated with a low-oxygen (8%) air mixture. Therefore, oxygen levels during the experiments were at a

Table 6: Primer sequences used in real-time PCR expression analysis for *P. pandorae irlandei* SSH libraries validation.

Genes	Primer sequences
Intracellular hemoglobin	For: 5' CTTGCCGATAACATTACTGCTGTTTCGAGG 3' Rev: 5' TCGGCATCACCCGCTTCTCCGCTACGTC 3'
Hemoglobin A2c	For: 5' GTTCTGATCATAATCGCTGTCTGTCTGG 3' Rev: 5' TGTAGAAGCTGGGCATTCAGCGTGTCCGGG 3'
Linker L1	For: 5' ATCATGGCAGGCCTGGTGGCACTCGCCAT 3' Rev: 5' TCTGATCCGTCATGACAGTCGTTAGCACC 3'
Linker L2	For: 5' ATGGTTGACGATGAGATGGACTTGATGGA 3' Rev: 5' ATCTTATAGCTGTCTATAGTAACCCG 3'
Hemoglobin B2	For: 5' CTGGATCATCTCGGCCGTCAGCATGTTGT 3' Rev: 5' GTGTGGTCGAGACGCGTTGCTCGGTCCGC 3'
PpandEST 2	For: 5' CAACCATGTGCCTACTTTACCTTGTTAG 3' Rev: 5' AAGAACAGCTGCTGCAGAATATCTCCAC 3'
Xylan endohydrolase	For: 5' GAGATGGAAGAGAGAAAATCCAGATTGGCT 3' Rev: 5' AGTTTCGCCTTCGAGTACCGCTGTAGGC 3'
Rab 7	For: 5' CTTATTCAAGCTAGCCACGAGATCCAGA 3' Rev: 5' CAACTCTCCGCTGAAGTCTTCGCCCGGTC 3'
Adenosylhomocysteinase	For: 5' ATTGTGTGCAATATTGGACATTTTGACTGTGA 3' Rev: 5' AGACCTAAATAGCCAGCCTGGTCATCTGA 3'
Chymotrypsin	For: 5' ACAGAGGTGCAATACGAGGTGATGACAAT 3' Rev: 5' ATGCCACCTGAGCTAAGAGTTCCCATCC 3'
PpandEST 1	For: 5' GAAGCTGACCTAGCTTACGCCGGTCTGAA 3' Rev: 5' TAGGGCTCGAGCGGCCGCCCGGGCAG 3'
Ribosomal protein S16	For: 5' GCTGTTGCTCACTGCAAACAGGGCAAAGGT 3' Rev: 5' TTGATCTCTTTCTTTGATGCTTCGTCGTC 3'
Nidogen secreted domain protein	For: 5' CAATGCAAGTATTGGCCATGGCATGGTAG 3' Rev: 5' ACCCAGCGTCTGCTTTGGCGACGTTACT 3'
Lipid binding protein	For: 5' TTCAACATGTCTCAATTGAATGGGAAATGGAA 3' Rev: 5' TCCAGGCCCGTTTCGTTACCTCGATACG 3'

maximum value of about 120 μM during circulation periods, and obviously decreased when the vessels were isolated, due to oxygen consumption by experimented animals. No mortality was observed at the end of the experiments, and samples of mussels (pool of gills, foot, mantle, adductor muscle) and worms (whole organisms) were collected and immediately frozen in liquid nitrogen until use.

RNA extraction

Total RNA was extracted from the gill, adductor muscle, mantle and foot of 5 thermally-challenged mussels, and 6 whole worms with the Trizol Reagent according to the manufacturer's instructions. Total RNAs extracted from the different tissues were pooled and poly(A⁺) mRNA was isolated using the PolyATtract[®]mRNA Isolation System (Promega, Madison, WI, USA) according to the manufacturer's instructions.

Suppression subtractive hybridization

Both forward (individuals incubated at 10°C versus individuals incubated at 20°C) and reverse (individuals incubated at 20°C versus individuals incubated at 10°C) subtracted libraries were produced from 2 μg of mRNA extracted from experimented mussels and worms. First and second strand cDNA synthesis, *RsaI* endonuclease enzyme digestion, adapter ligation, hybridization, and PCR amplification were performed as described in the PCR-select cDNA subtraction kit manual (Clontech, Palo Alto, CA, USA).

Cloning and sequencing

The differentially expressed PCR products were ligated into a pGEM-T vector (Promega, Madison, WI, USA) and 200 white colonies per library were cultured in LB medium supplemented with 100 mg/L ampicillin. Plasmids were then extracted using an alkaline lysis plasmid minipreparation, and sequenced using the Big Dye Terminator V3.1 Kit (Perkins-Elmer) and run on an AB3100 sequencer (Applied Biosystems Perkins-Elmer).

Sequence analysis and homology search

Chromatograms obtained after sequencing were treated with the Seqclean software (TGIR, the Institute for Genomic Research, Rockville, MD, USA) to remove vector and adaptor sequences. Cluster and contigs were then formed on each library sequence set. BLAST analyses of the sequences were performed on the NCBI server. The sequences were analyzed for homology with known sequences in databases using the BlastX and BlastN programs <http://www.ncbi.nlm.nih.gov/BLAST/>.

Validation of differential expression by real-time PCR

A validation step of the differentially expressed genes identified in *B. thermophilus* and *P. pandorae* SSH libraries

was carried out using real-time PCR and total RNA samples used for the SSH construction. For each sample, 5 μg of total RNA were submitted to reverse transcription using oligo-dT anchor primer (5'-GAC CAC GCG TAT CGA TGT CGA CT₍₁₆₎V-3') and M-MLV reverse transcriptase (Promega, Madison, WI, USA). Amplification of 20 genes isolated in the SSH libraries was carried out on the cDNA from both 10 and 20°C exposed mussel samples and 14 genes were amplified on pool of cDNA from samples of annelid exposed to both 10 and 20°C. The real-time PCR assay was performed in triplicate with 4 μL cDNA (1/20 dilution) in a final volume of 10 μL using the Chromo 4[™] System (BioRad). The concentrations of the reaction components were as follows: 1 \times Absolute[™] QPCR SYBR[®] Green mix (ABgene, UK) and 70 nM of each primer (Tables 5 and 6). The 18S ribosomal DNA was amplified as an endogenous PCR control, under the same amplification conditions using sense (5'-AAG GGC AGG AAA AGA AAC TAA C-3') and antisense (5'-GTT TCC CTC TAA GTG GTT TCA C-3') primers. The amplification was carried out as follows: initial enzyme activation at 94°C for 15 min, then 45 cycles of 94°C for 15 sec and 60°C for 1 min. A dissociation curve was generated and PCR efficiency (E) was estimated for each primer pair by using a serial dilution of reverse transcription products. Standard curves were generated for each primer pair and E was calculated using the formula $E = 10^{(-1/\text{slope})} - 1$. All primer pairs tested generated a single peak in the dissociation curve with a PCR efficiency estimated between 95 and 100%. Relative quantification (RQ) of each gene expression was calculated according to comparative CT method using the formula: $RQ = \text{exponential}(2^{-\Delta\Delta CT})$ with $\Delta\Delta CT = \Delta CT(10^\circ\text{C}) - \Delta CT(20^\circ\text{C})$ and $\Delta CT(10^\circ\text{C}) = CT(\text{gene X in the } 10^\circ\text{C sample}) - CT(18\text{S})$ and $\Delta CT(20^\circ\text{C}) = CT(\text{gene X in the } 20^\circ\text{C sample}) - CT(18\text{S})$; 18S ribosomal DNA being used as the endogenous control.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

IB carried out gene amplification and library screening, the sequence analysis and drafted the manuscript. AT produced cDNA libraries and helped to draft the manuscript. DJ and BS sampled the experimental animals, designed and performed the experimental work on board of the ship N/O L'Atalante using the pressurized vessels IPOC-AMP and helped to draft the manuscript. DM helped to draft the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We wish to thank the captain and crew of the NO L'Atalante, the crew of the Nautille submersible, and the chief scientist of the BioSpeedo cruise. We also thank the personnel of the Ouest-Géopôle sequencing platform in Roscoff, particularly Morgan Perennou. We would like to thank Stéphane

Hourdez for English editing. We thank the Marine Genomics Europe EDD Node. This work is part of the ANR project "AdapAlvinSTres" (# ANR-05-BLAN-0407-02) and was also supported by a grant from the Région Bretagne (PRIRE "Amethyst").

References

- Somero GN: **Proteins and temperature.** *Annu Rev Physiol* 1995, **57**:43-68.
- Hochachka PW, Somero GN: *Biochemical adaptation* Princeton: Princeton University Press; 1984.
- Hochachka PW, Somero GN: *Biochemical adaptation* New York: Oxford University Press; 2002.
- Eanes WF: **Analysis of selection on enzyme polymorphisms.** *Ann Rev Ecol Syst* 1999, **30**:301-326.
- Watt WB, Dean AM: **Molecular-functional studies of adaptive genetic variation in prokaryotes and eukaryotes.** *Annu Rev Genet* 2000, **34**:593-622.
- Somero GN: **Adaptation of enzymes to temperature: searching for basic "strategies".** *Comp Biochem Physiol B Biochem Mol Biol* 2004, **139**:321-333.
- Fields PA, Rudomin EL, Somero GN: **Temperature sensitivities of cytosolic malate dehydrogenases from native and invasive species of marine mussels (genus *Mytilus*): sequence-function linkages and correlations with biogeographic distribution.** *J Exp Biol* 2006, **209**:656-667.
- Von Damm KL: **Seafloor hydrothermal activity: black smoker chemistry and chimneys.** *Ann Rev Earth Planet Sci* 1990, **18**:173-204.
- Chevaldonné P, Desbruyères D, Le Hétre M: **Time-series of temperature from three deep-sea hydrothermal vent sites.** *Deep Sea Res* 1991, **38**:1417-1430.
- Sarrazin J, Juniper SK, Massoth GJ, Legendre P: **Physical and chemical factors influencing species distributions on hydrothermal sulfide edifices of the Juan de Fuca Ridge, northeast Pacific.** *Mar Ecol Prog Ser* 1999, **190**:89-112.
- Shank TM, Fornari DJ, Von Damm KL, Lilley MD, Haymon RM, Lutz RA: **Temporal and spatial patterns of biological community development at nascent deep-sea hydrothermal vents (9°50'N, East Pacific Rise).** *Deep Sea Res II* 1998, **45**:465-515.
- Sarrazin J, Juniper SK: **Biological characteristics of a hydrothermal edifice mosaic community.** *Mar Ecol Prog Ser* 1999, **185**:1-19.
- Luther GW, Rozan TF, Martial T, Nuzzio DB, Di Meo C, Shank TM, Lutz RA, Cary SC: **Chemical speciation drives hydrothermal vent ecology.** *Nature* 2001, **410**:813-816.
- Company R, Serafim A, Cosson RP, Fiala-Médioni A, Camus L, Colaço A, Serrão-Santos R, Bebianno MJ: **Antioxidant biochemical responses to long-term copper exposure in *Bathymodiolus azoricus* from Menez-Gwen hydrothermal vent.** *Sci Total Environ* 2008, **389**:407-417.
- Hardivillier Y, Denis F, Demattei MV, Bustamante P, Laulier M, Cosson R: **Metal influence on metallothionein synthesis in the hydrothermal vent mussel *Bathymodiolus thermophilus*.** *Comp Biochem Physiol C Toxicol Pharmacol* 2006, **143G**:321-332.
- Bettencourt R, Roch P, Stefanni S, Rosa D, Colaço A, Santos RS: **Deep sea immunity: unveiling immune constituents from the hydrothermal vent mussel *Bathymodiolus azoricus*.** *Mar Environ Res* 2007, **64**:108-127.
- Tanguy A, Bierne N, Saavedra C, Pina B, Bachere E, Bonhomme F, Boudry P, Boulo V, Boutet I, Dossat C, Favrel P, Huvet A, Jollivet D, Klages S, Lapegue S, Moal J, Moraga D, Reinhard R, Samain JF, Zouros E, Canario A: **Increasing genomic information in Bivalves by new EST collections in four species.** *Gene* 2008, **408**:27-36.
- Craddock C, Hoeh WR, Gustafson RG, Lutz RA, Hashimoto J, Vrijenhoek RJ: **Evolutionary relationships among deep-sea mytilids (*Bivalvia*: Mytilidae) from hydrothermal vents and cold-water methane/sulphide.** *Mar Biol* 1995, **121**:477-485.
- Fiala-Médioni A, McKiness Z, Dando P, Boulegue J, Mariotti A, Alayse-Danet A, Robinson J, Cavanaugh C: **Ultrastructural, biochemical, and immunological characterization of two populations of a new species of mytilid mussel, *Bathymodiolus azoricus*, from the Mid-Atlantic Ridge: evidence for a dual symbiosis.** *Mar Biol* 2002, **141**:1035-1043.
- Duperron S, Bergin C, Zielinski F, Blazejak A, Pernthaler A, McKiness ZP, DeChaine E, Cavanaugh CM, Dubilier N: **A dual symbiosis shared by two mussel species, *Bathymodiolus azoricus* and *Bathymodiolus puteoserpentis* (*Bivalvia*: Mytilidae), from hydrothermal vents along the northern Mid-Atlantic Ridge.** *Environ Microbiol* 2006, **8**:1441-1447.
- Ravaux J, Gaill F, Le Bris N, Sarradin PM, Jollivet D, Shillito B: **Heat-shock response and temperature resistance in the deep-sea vent shrimp *Rimicaris exoculata*.** *J Exp Biol* 2007, **206**:2345-2354.
- Cottin D, Ravaux J, Léger N, Halary S, Toullec JY, Sarradin PM, Gaill F, Shillito B: **Thermal biology of the deep-sea vent annelid *Paralvinella grasslei*: in vivo studies.** *J Exp Biol* 2008, **211**:2196-2204.
- Pruski AM, Dixon DR: **Toxic vents and DNA damage: first evidence from a naturally contaminated deep-sea environment.** *Aquat Toxicol* 2003, **64**:1-13.
- Hourdez S, Weber RE: **Molecular and functional adaptations in deep-sea hemoglobins.** *J Inorg Biochem* 2005, **99**:130-141.
- Lee RW: **Thermal tolerances of deep-sea hydrothermal vent animals from the northeast Pacific.** *Biol Bull* 2003, **205**:98-101.
- Girguis PR, Lee RW: **Thermal preference and tolerance of Alvinellids.** *Science* 2006, **312**:231.
- Jollivet D, Desbruyères D, Ladrat C, Laubier L: **Evidence for differences in the allozyme thermostability of deep-sea hydrothermal vent polychaetes (*Alvinellidae*): a possible selection by habitats.** *Mar Ecol Prog Ser* 1995, **123**:125-136.
- Van Dover CL: **Community structure of mussel beds at deep-sea hydrothermal vents.** *Mar Ecol Prog Ser* 2000, **230**:137-156.
- Fisher CR, Childress JJ, Arp AJ, Brooks JM, Distel D, Favuzzi JA, Felbeck H, Hessler R, Johnson KS, Kennicutt MC II, Macko SA, Newton A, Powell MA, Somero GN, Soto T: **Microhabitat variation in the hydrothermal vent mussel, *Bathymodiolus thermophilus*, at the Rose Garden vent on the Galapagos Rift.** *Deep Sea Res* 1988, **35**:1769-1789.
- Jollivet D: **Specific and genetic diversity at deep-sea hydrothermal vents: an overview.** *Biodiv Conserv* 1996, **5**:1619-1653.
- Jollivet D: **Distribution et évolution de la faune associée aux sources hydrothermales profondes à 13°N sur la dorsale du Pacifique oriental: le cas particulier des polychètes Alvinellidae.** Thèse de Doctorat NR, Université de Bretagne Occidentale, France; 1993.
- Sanders BM: **Stress proteins in aquatic organisms: an environmental perspective.** *Crit Rev Toxicol* 1993, **23**:49-75.
- Feder ME, Hofmann GE: **Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology.** *Annu Rev Physiol* 1999, **61**:243-282.
- Berger MS, Young CM: **Physiological response of the cold-seep mussel *Bathymodiolus childressi* to acutely elevated temperature.** *Mar Biol* 2006, **149**:1397-1402.
- Pruski AM, Dixon DR: **Heat shock protein expression pattern (HSP70) in the hydrothermal vent mussel *Bathymodiolus azoricus*.** *Mar Environ Res* 2007, **64**:209-224.
- Clarke A: **Temperature and energetics: an introduction to cold ocean physiology.** In *Cold Ocean Physiology* Edited by: Pörtner HO, Playle RC. Cambridge: Cambridge University Press; 1998:3-33.
- Pörtner HO: **Climate change and temperature-dependent biogeography: oxygen limitation of thermal tolerance in animals.** *Naturwissenschaften* 2001, **88**:137-146.
- Storey KB: **Molecular mechanisms of metabolic arrest in molluscs.** In *Surviving Hypoxia: Mechanisms of Control and Adaptation* Edited by: Hochachka PW, Lutz PL, Sick TJ, Rosenthal M, Thillart G. Boca Raton: CRC Press; 1993:253-271.
- Larade K, Storey KB: **A profile of the metabolic responses to anoxia in marine invertebrates.** In *Cell and Molecular Responses to Stress Volume 3*. Edited by: Storey KB, Storey JM. Amsterdam: Elsevier Press; 2002.
- Peck LS: **Ecophysiology of Antarctic marine ectotherms: limits to life.** *Polar Biol* 2002, **25**:31-40.
- Sokolova IM, Pörtner HO: **Metabolic plasticity and critical temperatures for aerobic scope in a eurythermal marine invertebrate (*Littorina saxatilis*, Gastropoda: Littorinidae) from different latitudes.** *J Exp Biol* 2003, **206**:195-207.
- Peck LS, Conway LZ: **The myth of metabolic coldadaptation: oxygen consumption in stenothermal Antarctic bivalves.** In *The evolutionary biology of the bivalvia Volume 177*. Edited by: Harper EM, Taylor JD, Crame JA. Geol Soc Spec Publ London; 2000:441-445.
- Peck LS: **Prospects for survival in the Southern ocean: extreme temperature sensitivity of benthic species.** *R Soc Spec Issue Antarct Sci* 2005, **17**:497-507.

44. Anestis A, Lazou A, Pörtner HO, Michaelidis B: **Behavioral, metabolic, and molecular stress responses of marine bivalve *Mytilus galloprovincialis* during long-term acclimation at increasing ambient temperature.** *Am J Physiol Regul Integr Comp Physiol* 2007, **293**:R911-R921.
45. Desbruyères D, Segonzac M, Bright M: *Handbook of the deep-sea hydrothermal vent fauna* 2nd edition. Linz, Austria: Denisia 18; 2006.
46. Boutet I, Tanguy A, Moraga D: **Response of the Pacific oyster *Crassostrea gigas* to hydrocarbon contamination under experimental conditions.** *Gene* 2004, **329**:147-157.
47. Tanguy A, Guo X, Ford SE: **Searching for genes involved in host defense against *Perkinsus marinus* and *Haplosporidium nelsoni* infections in the eastern oysters.** *Gene* 2004, **329**:121-131.
48. Tanguy A, Boutet I, Laroche J, Moraga D: **Molecular identification and expression study of differentially regulated genes in the Pacific oyster *Crassostrea gigas* in response to pesticide exposure.** *FEBS J* 2005, **272**:390-403.
49. David E, Tanguy A, Pichavand K, Moraga D: **Response of the Pacific oyster *Crassostrea gigas* to hypoxia exposure under experimental conditions.** *FEBS J* 2005, **272**:5635-5652.
50. Strong SJ, Ellington WR: **Isolation and sequence analysis of the gene for arginine kinase from the chelicerate arthropod, *Limulus polyphemus*: insights into catalytically important residues.** *Biochim Biophys Acta* 1995, **1246**:197-200.
51. Wyss M, Smeitink J, Wevers RA, Wallimann T: **Mitochondrial creatine kinase: a key enzyme of aerobic energy metabolism.** *Biochim Biophys Acta* 1992, **1102**:119-166.
52. Ellington WR: **Evolution and physiological roles of phosphagen systems.** *Annu Rev Physiol* 2001, **63**:289-325.
53. Greenway SC, Storey KB: **The effect of prolonged anoxia on enzyme activities in oysters (*Crassostrea virginica*) at different seasons.** *J Exp Mar Biol Ecol* 1999, **242**:259-272.
54. Abe H, Hirai S, Okada S: **Metabolic responses and arginine kinase expression under hypoxic stress of the kuruma prawn *Marsupenaeus japonicus*.** *Comp Biochem Physiol A* 2007, **146**:40-46.
55. Dahlhoff E, Somero GN: **Pressure and temperature adaptation of cytosolic malate dehydrogenases of shallow- and deep-living marine invertebrates: Evidence for high body temperatures in hydrothermal vent animals.** *J Exp Biol* 1991, **159**:473-487.
56. Dahlhoff E, Somero GN: **Kinetic and structural adaptations of cytoplasmic malate dehydrogenases of eastern Pacific abalone (genus *Haliotis*) from different thermal habitats: Biochemical correlates of biogeographical patterning.** *J Exp Biol* 1993, **185**:137-150.
57. Nadeau K, Das A, Walsh CT: **Hsp90 chaperonins possess ATPase activity and bind heat-shock transcription factors and peptidylprolyl isomerases.** *J Biol Chem* 1993, **268**:1479-1487.
58. Jakob U, Buchner J: **Assisting spontaneity – The role of Hsp90 and small HSPs as molecular chaperones.** *Trends Biochem Sci* 1994, **19**:205-211.
59. Rutherford S, Lindquist S: **Hsp90 as a capacitor for morphological evolution.** *Nature* 1998, **396**:336-342.
60. Arbona M, Defrutos R, Tanguay RM: **Transcriptional and translational study of the *Drosophila subobscura* hsp83 gene in normal and heat-shock conditions.** *Genome* 1993, **36**:694-700.
61. Salvucci ME, Stecher DS, Henneberry TJ: **Heat shock proteins in whiteflies, an insect that accumulates sorbitol in response to heat stress.** *J Therm Biol* 2000, **25**:363-371.
62. Rinehart JP, Denlinger DL: **Heat-shock protein 90 is down-regulated during pupal diapause in the flesh fly, *Sarcophaga crassipalpis*, but remains responsive to thermal stress.** *Insect Mol Biol* 2000, **9**:641-645.
63. Dietz TJ, Somero GN: **Acclimatization in Eurythermal Goby Fishes (Genus *Gillichthys*) The Threshold Induction Temperature of the 90-kDa Heat Shock Protein is Subject.** *Proc Natl Acad Sci USA* 1992, **89**:3389-3393.
64. Sinensky M: **Homeoviscous adaptation-A homeostatic process that regulates the viscosity of membrane lipids in *Escherichia coli*.** *Proc Natl Acad Sci USA* 1974, **71**:522-525.
65. Mathias S, Pena LA, Kolesnick RN: **Signal transduction of stress via ceramide. Ceramide also plays a pivotal role in stress-induced apoptosis.** *Biochem J* 1998, **335**:465-480.
66. Mattie MD, Freedman JH: **Protective effects of aspirin and vitamin E (α -tocopherol) against copper- and cadmium-induced toxicity.** *Biochem Biophys Res Com* 2001, **285**:921-925.
67. Bebianno MJ, Company R, Serafima A, Camusb L, Cosson RP, Fiala-Medoni A: **Antioxidant systems and lipid peroxidation in *Bathymodiulus azoricus* from Mid-Atlantic Ridge hydrothermal vent fields.** *Aquat Toxicol* 2005, **75**:354-373.
68. Company R, Serafim A, Bebianno MJ, Cosson R, Shillito B, Fiala-Medioni A: **Effect of cadmium, copper and mercury on antioxidant enzyme activities and lipid peroxidation in the gills of the hydrothermal vent mussel *Bathymodiulus azoricus*.** *Mar Environ Res* 2004, **58**:377-381.
69. Sasaki T, Hohenester E, Gohring W, Timpl R: **Crystal structure and mapping by site directed mutagenesis of the collagen binding epitope of an activated form of BM-40/SPARC/osteonectin.** *EMBO J* 1998, **17**:1625-1634.
70. Sweetwyne MT, Brekken RA, Workman G, Bradshaw AD, Carbon J, Siadak AW, Murri C, Sage EH: **Functional analysis of the matricellular protein SPARC with novel monoclonal antibodies.** *J Histochem Cytochem* 2004, **52**:723-733.
71. Kudo H, Hirayoshi K, Kitagawa Y, Imamura S, Nagata K: **Two collagen-binding proteins, osteonectin and HSP47, are coordinately induced in transformed keratinocytes by heat and other stresses.** *Exp Cell Res* 1994, **212**:219-224.
72. Neri M, Descali-Cancedda F, Cancedda R: **Heat shock response in cultured chick embryo chondrocytes. Osteonectin is a secreted heat-shock protein.** *Eur J Biochem* 1992, **205**:569-574.
73. Emerson RO, Sage EH, Ghosh JG, Clark JI: **Chaperone-like activity revealed in the matricellular protein SPARC.** *J Cell Biochem* 2006, **98**:701-705.
74. Fridovich I: **Oxygen toxicity: a radical explanation.** *J Exp Biol* 1998, **201**:1203-1209.
75. Bryant C: *Metazoan Life without Oxygen* Suffolk: Chapman and Hall; 1991.
76. Fenchel T, Finlay BJ: *Ecology and Evolution in Anoxic Worlds* New York: Oxford University Press; 1995.
77. Tielens AGM, Van Hellemond JJ: **The electron transport chain in anaerobically functioning eukaryotes.** *Biochim Biophys Acta Bioenerg* 1998, **1365**:71-78.
78. Halliwell B, Gutteridge JMC: *Free Radicals in Biology and Medicine* Oxford: Clarendon Press; 1999.
79. Boon K, Caron HN, van Asperen R, Valentijn L, Hermus MC, van Sluis P, Roobeek I, Weis I, Voăute PA, Schwab M, et al.: **N-myc enhances the expression of a large set of genes functioning in ribosome biogenesis and protein synthesis.** *EMBO J* 2001, **20**:1383-1393.
80. Schuldiner O, Benvenisty N: **A DNA microarray screen for genes involved in c-MYC and N-MYC oncogenesis in human tumors.** *Oncogene* 2001, **20**:4984-4994.
81. Watson JD, Oster SK, Shago M, Khosravi F, Penn LZ: **Identifying genes regulated in a Myc-dependent manner.** *J Biol Chem* 2002, **277**:36921-36930.
82. Matsumura I, Tanaka H, Kanakura Y: **E2F1 and c-Myc in cell growth and death.** *Cell Cycle* 2003, **2**:333-338.
83. To KW, Koshiji M, Hammer S, Huang LE: **Genetic instability: the dark side of the hypoxic response.** *Cell Cycle* 2005, **4**:881-882.
84. Dela Haba G, Cantoni GL: **The enzymatic synthesis of S-adenosyl-L-homocysteine from adenosine and homocysteine.** *J Biol Chem* 1959, **234**:603-608.
85. Liu J, Benbrahim-Tallaa L, Qian X, Yu L, Xie Y, Boos J, Qu W, Waalkes MP: **Further studies on aberrant gene expression associated with arsenic-induced malignant transformation in rat liver TRL1215 cells.** *Toxicol Appl Pharmacol* 2006, **216**:407-415.
86. Bagnyukova TV, Luzhna LI, Pogribny IP, Lushchak VI: **Oxidative stress and antioxidant defenses in goldfish liver in response to short-term exposure to arsenite.** *Environ Mol Mutagen* 2007, **48**:658-665.
87. Rajagopal S, Gaag M Van der, Velde G Van der, Jenner HA: **Upper temperature tolerances of exotic brackish-water mussel, *Mytilopsis leucophaea* (Conrad): An experimental study.** *Mar Environ Res* 2005, **60**:512-530.
88. Govener B, Freeman M, Bergquist DC, Johnson GA, Fisher CR: **Composition of a One-Year-Old *Riftia pachyptila* Community Following a Clearance Experiment: Insight to Succession Patterns at Deep-Sea Hydrothermal Vents.** *Biol Bull* 2004, **207**:177-182.
89. Toulmond A, Slitine F, Frescheville J, Jouin C: **Extracellular hemoglobins of hydrothermal vent annelids: structural and functional characteristics in three alvinellid species.** *Biol Bull* 1990, **179**:366-373.

90. Hourdez S, Lallier FH, De Cian M, Green BN, Weber RE, Toulmond A: **Gas Transfer System in *Alvinella pompejana* (Annelida Polychaeta, Terebellida): Functional Properties of Intracellular and Extracellular Hemoglobins.** *Physiol Biochem Zool* 2000, **73**:365-373.
91. Liochev S, Kuchumov S, Vonogradov N, Fridovich I: **Superoxide dismutase activity in the giant haemoglobin of the earthworm, *Lumbricus terrestris*.** *Arch Biochem Biophys* 1996, **330**:281-284.
92. Yurchenco PD, O'Rear JJ: **Basal lamina assembly.** *Curr Opin Cell Biol* 1994, **6**:674-681.
93. Timpl R, Brown JC: **Supramolecular assembly of basement membranes.** *Bioassay* 1996, **18**:123-132.
94. Fox JW, Mayer U, Nischt R, Aumailley M, Reinhardt D, Wiedemann H, Mann K, Timpl R, Krieg T, Engel J, Chu ML: **Recombinant nidogen consists of three globular domains and mediates binding of laminin to collagen type IV.** *EMBO J* 1991, **10**:3137-3146.
95. Rando RR: **Chemical biology of protein isoprenylation/methylation.** *Biochim Biophys Acta* 1996, **1300**:5-16.
96. Waters MG, Pfeffer SR: **Membrane tethering in intracellular transport.** *Curr Opin Cell Biol* 1996, **11**:453-459.
97. Nielsen E, Severin F, Backer J, Hyman A, Zerial M: **Rab5 regulates motility of early endosomes on microtubules.** *Nat Cell Biol* 1995, **1**:376-382.
98. Meresse S, Gorvel JP, Chavrier P: **The rab7 GTPase resides on a vesicular compartment connected to lysosomes.** *J Cell Sci* 1995, **108**:3349-3358.
99. Bucci C, Thomsen P, Nicoziani P, McCarthy J, van Deurs B: **Rab 7, a key to lysosome biogenesis.** *Mol Biol Cell* 2000, **11**:467-480.
100. Jollivet D, Lallier FH, Barnay AS, Bienvenu N, Bonnivard E, Briand P, Cambon-Bonavita MA, Comtet T, Cosson R, Daguin C, Donval JP, Faure B, Gaillard M, Glippa V, Guillou L, Hourdez S, Le Bris N, Morand S, Pradillon F, Rees JF, Segonzac M, Shillito B, Thiebaut D, Viard F: **The BIOSPEEDO Cruise: a new survey of hydrothermal vents along the South East Pacific Rise from 7°24'S to 21°33'S.** *InterRidge* 2004, **13**:20-26.
101. Shillito B, Jollivet D, Sarradin PM, Rodier P, Lallier F, Desbruyères D, Gaill F: **Temperature resistance of *Hesiolyra bergi*, a polychaetous annelid living on deep-sea vent smoker walls.** *Mar Ecol Prog Ser* 2001, **216**:141-149.
102. Dixon DR, Pruski AM, Dixon LR: **The effects of hydrostatic pressure change on DNA integrity in the hydrothermal-vent mussel *Bathymodiulus azoricus*: implications for future deep-sea mutagenicity studies.** *Mutat Res* 2004, **552**:235-246.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

