

# Tephrochronological dating of Holocene moraines at Icelandic glaciers, and climatic interpretations

Martin P. Kirkbride

### ▶ To cite this version:

Martin P. Kirkbride. Tephrochronological dating of Holocene moraines at Icelandic glaciers, and climatic interpretations. Iceland in the Central Northern Atlantic: hotspot, sea currents and climate change, May 2010, Plouzané, France. hal-00482052

## HAL Id: hal-00482052 https://hal.univ-brest.fr/hal-00482052v1

Submitted on 7 May 2010  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## TEPHROCHRONOLOGICAL DATING OF HOLOCENE MORAINES AT ICELANDIC GLACIERS, AND CLIMATIC INTERPRETATIONS.

Martin Kirkbride School of Social and Environmental Sciences University of Dundee Dundee DD1 4HN

#### Abstract

Fluctuations of Icelandic glaciers reveal the impact of regional climate change on the cryosphere, filtered by the different response characteristics of individual glaciers. Frequent tephra deposition upon steadily aggrading aeolian soils provides a useful dating environment, in which basal tephras often provide close minimum ages on underlying tills and outwash deposits in areas where the local tephrostratigraphy is well constrained. We have dated moraines at glaciers across Iceland to improve the Holocene glacial chronology in terms of its temporal extent and resolution. Tephrochronology also provides a test of lichenometric dating, an area for further research. At least five groups of regionallysynchronous advances occurred between c. AD 1700 and 1930 during the "Little Ice Age". The maximum extent of "Little Ice Age" glaciers varies by up to 200 years across Iceland, due more to the response chraracteristics of individual glaciers than to regional climatic variation. At Gígjökull, two glacier advances occurred before the 3rd century AD, with others in the 9th and 12th centuries AD bracketing the Medieval Warm Period. In central and northern Iceland, earlier glacier advances are dated to c. 4.5-5.0, c.3.0-3.5 ka BP, c. 2.0-2.5 ka BP. This classic "Neoglacial" sequence is comparable to other parts of Europe and Scandinavia, but is discernible only at smaller mountain glaciers. In contrast, the 19<sup>th</sup>-Century advance of large ice caps censored evidence of earlier fluctuations from the moraine record, and preservation potential is preconditioned by glacier type. In general, the forefields of steep, fast-responding glaciers contain more complete archives of Holocene climatic changes than do the margins of the large icefields. Glacier advances appear to be favoured by a weakening of zonal circulation (the negative mode of the North Atlantic Oscillation) associated with cooler, drier winters and cooler, wetter summers.

#### References

Kirkbride, M.P. and Dugmore, A.J. (2008) Two millennia of glacier advances in southern Iceland dated by tephrochronology. *Quaternary Research* 70, 398-411.

Kirkbride, M.P. and Dugmore, A.J. (2006). Responses of mountain ice caps in central Iceland to Holocene climate change. *Quaternary Science Reviews* 25, 1692-1707.

Kirkbride, M.P. (2002). Icelandic climate and glacier fluctuations through the terminus of the "Little Ice Age". *Polar Geography* **26**, 116-133.

Kirkbride, M.P. and Dugmore, A.J. (2001a) Can lichenometry be used to date the "Little Ice Age" glacial maximum in Iceland? *Climatic Change* **48**, 151-167.

Kirkbride, M.P. and Dugmore, A.J. (2001b) Timing and significance of mid-Holocene glacier advances in northern and central Iceland. *Journal of Quaternary Science* **16**, 145-153.