

The food poisoning power of *Bacillus cereus* Group strains varies 1 according to phylogenetic affiliation (groups I-VII), not to species affiliation

Marie-Hélène Guinebretière, Philippe Velge, Olivier Couvert, Frédéric Carlin, Marie-Laure Debuyser, Christophe Nguyen-The

► To cite this version:

Marie-Hélène Guinebretière, Philippe Velge, Olivier Couvert, Frédéric Carlin, Marie-Laure Debuyser, et al.. The food poisoning power of *Bacillus cereus* Group strains varies 1 according to phylogenetic affiliation (groups I-VII), not to species affiliation. *Journal of Clinical Microbiology*, American Society for Microbiology, 2010, 48 (9), pp.3388-3391. 10.1128/JCM.00921-10 . hal-00560894

HAL Id: hal-00560894

<https://hal.univ-brest.fr/hal-00560894>

Submitted on 31 Jan 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **The food poisoning power of *Bacillus cereus* Group strains varies according to**
2 **phylogenetic affiliation (groups I-VII), not to species affiliation**

3
4
5 **Marie-Hélène Guinebretière,^{1,2*} Philippe Velge,³ Olivier Couvert,⁴ Frédéric Carlin,^{1,2} Marie-**
6 **Laure Debuysier,⁵ Christophe Nguyen-The,^{1,2}**

7
8 ¹INRA, UMR408 Sécurité et Qualité des Produits d'Origine Végétale, F-84000 Avignon, France

9 ²Université d'Avignon et des Pays du Vaucluse, UMR408 Sécurité et Qualité des Produits d'Origine Végétale, F-84000
10 Avignon, France

11 ³INRA, UR1282, Infectiologie animale et santé publique, F-37380 Nouzilly, France

12 ⁴Université Européenne de Bretagne, Université de Brest, LUBEM EA 3882, UMT Physiopt 08.3, 6 rue de l'Université,
13 29334 Quimper cedex, France

14 ⁵AFSSA, Laboratoire d'études et de recherches sur la qualité des aliments et sur les procédés agroalimentaires,
15 23 avenue du Général-Leclerc, F-94706 Maisons-Alfort Cedex, France

16
17

18 **For: Journal of Clinical Microbiology**

19

20 **Running title:** *Food poisoning power and B. cereus phylogeny*

21

22 **Keywords:** food poisoning power, cytotoxicity, Toxins, *Bacillus cereus* Group, phylogenetic
23 groups, food safety

24

25 ***Correspondence.** Marie-Hélène Guinebretière, INRA, UMR A408, INRA – Université
26 d'Avignon, Domaine Saint-Paul, Site Agroparc, F-84914 Avignon Cedex 9, France. Tel.: (33) 4 32
27 72 25 24; Fax: (33) 4 32 72 24 92; e-mail: guinebre@avignon.inra.fr.

28

1 **ABSTRACT**

2

3 Cytotoxic activity levels of culture filtrates and toxin distribution varied according to the phylogenetic
4 group (I-VII) inside the *B. cereus* Group, suggesting that these groups are of different clinical significance
5 and are more suitable than species affiliation for determining food poisoning risk. A first-line, simple online
6 tool (<https://www.tools.symprevius.org/Bcereus/>) to assign strains to the different phylogenetic groups is
7 presented.

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

1 The *B. cereus* Group (*B. cereus sensu lato*) can induce diarrhea by producing heat-labile enterotoxins
2 during growth in the small intestine (8) and emesis by producing a single heat-stable peptide toxin called
3 cereulide (1). A suitable test for the detection of emetic toxin producers is available (7). For diarrhoeal
4 syndrome, three enterotoxins have been described: the cytotoxin CytK (17), the nonhaemolytic enterotoxin
5 Nhe (9) and the haemolysin HBL (2). Also, other factors may have a role in certain isolates. A way of
6 evaluating the enterotoxicity produced by bacterial strains is to test cytotoxicity of culture filtrates on cell
7 lines such as Vero cells (18) or Caco2 cells (4) which encompass all soluble factors.

8 Assessing the food poisoning power of *B. cereus sensu stricto* and other species of the Group
9 (*B. thuringiensis*, *B. mycoides*, *B. pseudomycoides*, *B. weihenstephanensis* and *B. anthracis*) may not be
10 feasible because they are not separate genomic species, some being distributed between few distinct
11 phylogenetic groups (10), some other both belonging to a same phylogenetic groups (Table 1A). In contrast,
12 the seven phylogenetic groups described in Table 1A covers the entire *B. cereus* Group (*sensu lato*), are
13 based on genetic delimitation (10) and represent a more homogeneous basis to assess the food poisoning
14 power. The aim of the present study was to determine food poisoning toxins and global cytotoxicity associated
15 with each of the seven phylogenetic groups to evaluate their food poisoning power.

16
17 A total of 391 independent strains (10) belonging to the *B. cereus* Group were used to determine presence
18 of known toxin genes. In a first step total DNA was extracted for each strain as previously described (13).
19 Detection of *hbl* genes and *nhe* genes was performed by PCR and Southern blots (to check false negative
20 responses) as previously described (11). New primer pairs were also designated to detect *nhe* genes in the
21 two most distant groups (I and VII), taking into account the sequence polymorphism observed between the
22 sequence of genome NVH 391-98 (group VII) and that of other sequenced genomes
23 (ATCC 14579, E33L, Ames, ATCC 10987 and KBAB4): ^{5'}GTAAATGCTGCVGATAGYCAAAC^{3'} / ^{5'}GGCATV
24 ATRTTYCCTGCTGC^{3'} and ^{5'}GGTTCRAAYGCTTTAGTAATGG^{3'} / ^{5'}ATTCCWGCRTCTTGACTAGC^{3'}, targeting
25 *nheAB* genes and *nheB* gene respectively. Strains carrying the *cytK* gene and its two polymorphic forms
26 *cytK-1* and *cytK-2* were detected using the duplex PCR (validated as without false positive or false negative
27 response) described previously (12). Cereulide producers were detected using *ces* gene-specific PCR (7).

28 A subset of 97 strains, belonging to the different phylogenetic groups (II to VII), was characterized for
29 cytotoxicity activity of culture filtrates. Cytotoxic activity was measured on Caco2 cells as described

1 previously (4). The Caco2 cell viability (V) was measured by DO_{620nm} , in % of the total viability (100%
2 viability was DO_{620nm} obtained for Caco2 cells treated with non-inoculated BHI instead of culture filtrates),
3 for serial 2-fold diluted culture filtrates ($D = 0$ to $1/32$). The non-linear regression ($V = f(D)$) was obtained
4 from duplicate values of the six tested dilutions. This curve and the D value for which V was 50% (SC50)
5 were calculated using the GraphPad Prism software (GraphPad software incorporated, San Diego, USA).
6 When no cytotoxic activity was recorded, SC50 was approximated to 1 to allow calculations. The
7 cytotoxicity activity (CA) of culture filtrates was expressed as $CA = 1 - SC50$. All the strains were tested in
8 the course of two independent experiments.

9 Fisher's test was used to determine whether bacterial populations were significantly distant for the tested
10 character, using the XLSTAT 2008 software package.

11

12 As two emetic strains (20) have been recently associated with other groups than the two usually known
13 lineages (6) (21), we examined the distribution of the cereulide-producing strains over all seven phylogenetic
14 groups. In our study, cereulide producers (7), were found only in group III, in subgroup III-2 containing the
15 emetic strain F4810 and subgroup III-3 (Table 1B), which correspond to the known emetic lineages. None of
16 the 137 tested strains in group VI (*B. thuringiensis* VI, *B. weihenstephanensis* and *B. mycooides*) carried the
17 *ces* gene, which suggests that emetic *B. weihenstephanensis* strains, first observed by Thorsen *et al.* (20), are
18 probably extremely rare.

19 *Nhe* genes were carried by 100% of the tested strains in each group and subgroup (Table 1B), in
20 accordance with data from available sequenced genomes affiliated to the *B. cereus* Group (82 sequenced
21 genomes). We can therefore now definitively assert that *nhe* genes are a constant part of the *B. cereus* Group
22 strains. In contrast, *hbl* genes frequency varied from 40% to 97% between the phylogenetic groups I, II, IV,
23 V and VI, and this operon was seldom carried by strains of the phylogenetic group III, particularly in
24 subgroups III-2 and III-3 containing emetic strains and III-4 containing the *B. anthracis* strains (Table 1B).
25 Thus rarity of *hbl* genes is not a specificity of the 'emetic' lineages as previously published (6) but also
26 concern other sub-groups in the phylogenetic group III (III-4). As for *hbl* genes, the distribution of *cytK* gene
27 was disrupted. However, it showed some interesting specificities (Table 1B) : *CytK-1* form was specific of
28 the most distant group VII '*B. cytotoxicus*' (versus the 387 strains tested from the other groups); *cytK-2* form
29 was particularly frequent in mesophilic groups III and IV (except for the 12 tested *B. anthracis* strains that

1 were negative in III-4 sub-group) whereas it was rare or absent in the psychrotolerant or moderately
2 psychrotolerant groups (VI, II and V). *CytK* gene was also absent in group I strains.

3 The disrupted distribution of *hbl* and *cytK* genes is in accordance with what is observed from available
4 sequenced genomes. It opens the way to new hypothetical mechanisms for spread history of enterotoxins,
5 such as horizontal gene transfers (HGT) in the *B. cereus* Group. This is coherent with traces of HGT
6 previously observed on *B. cereus* genome around *hbl* genes (14) or with existence of plasmids carrying
7 enterotoxin-like genes (15, 16). On this assumption, some phylogenetic groups might have failed to acquire
8 *cytK* or *hbl* (for example group VI for *cytK* gene or sub-group III-2 for *hbl*) because they occupy very
9 specific thermal niches unfavorable to cohabitation and exchange of genetic material with other phylogenetic
10 groups.

11 Low or no cytotoxic activity of culture filtrates was recorded for all the strains of group VI and subgroup
12 III-3 (Fig. 1). By contrast, higher cytotoxic activity ($p < 0.001$) was recorded for strains of the other groups
13 (Fig. 1a), reaching very high values for a great number of strains in group III. Cytotoxic activity was
14 heterogeneous in group III (Fig. 1b), with high cytotoxic activity in groups III-1 and III-2 and very low
15 cytotoxic activity in subgroup III-3 ($p < 0.0001$). In addition, the cytotoxicity recorded for strains of
16 subgroup IV-2 (Fig. 1b) was higher than for subgroups IV-1 and IV-3 ($p < 0.05$) in group IV. Finally, strain
17 391-98, representing group VII ('*B. cytotoxicus*'), exhibited a high level of cytotoxic activity (mean = 0.75).

18
19 The enterotoxic potential conferred by each phylogenetic group, independently of species affiliation, has
20 implications for health risk. Based on the frequency of highly enterotoxic strains (generally associated with
21 diarrheal syndrome) recorded in each group, food poisoning risk should be the highest for group III
22 (containing *B. cereus* III, *B. thuringiensis* III and *B. anthracis* like strains), particularly subgroups III-1, III-2
23 and III-4. Sub-groups III-2 and III-3 may be also dangerous for their emetic potential. This risk remains high
24 for groups VII, IV and II, decreases with group V and seems very low for group VI. Strains belonging to this
25 last group should be the safest with regard to diarrheal syndrome. This is consistent with the absence of
26 strains isolated from foodborne disease in this group (10). Concerning emetic syndrome, the possible
27 production of emetic toxin by some group VI strains, although presumably very rare, needs to be clarified.
28 As *B. weihenstephanensis* and *B. mycoides* strains are delimited into group VI, we can also consider these
29 two species as representing a low level of risk, on condition that they are reliably identified. For *B. cereus*

1 *sensu stricto* and *B. thuringiensis* species, the level of risk is impossible to determine without taking into
2 account phylogenetic affiliation of strains (I to VII), as they can belong to the phylogenetic groups II to VI
3 and thus reflect several different levels of food poisoning risk.

4 In conclusion, affiliation to a phylogenetic group provides a more accurate indication of the risk than
5 affiliation to a current species of the *B. cereus* Group. Associating the phylogenetic number (I-VII) to the
6 species name when identifying new isolates offers a first useful indicator of risk. The challenge is thus to
7 propose reliable methods for identifying the seven phylogenetic groups.

8 As shown in this study, the validated duplex-PCR method (12) permits to identify *cytK-1*-carrying strains
9 specific of the rare and hazardous group VII (*B. cytotoxicus*). A rapid method to assign bacterial isolates to
10 groups I-VII is to compare *panC* gene sequence from new isolates with those of reference strains published
11 in Guinebretiere *et al.* (10). For this purpose, we propose the online tool at the following link:
12 <https://www.tools.symprevius.org/Bcereus/english.php>. As *panC* gene sequence was previously shown to
13 reliably delimit the phylogenetic groups and sub-groups (10), this tool allows a rapid and reliable
14 identification (100% exact responses) at the phylogenetic group level (I-VII). It compares a query sequence
15 with those in a database (85 *panC* sequences for which the phylogenetic affiliation is known). Protocols and
16 homology search algorithm are available from the site.

17 This phylogenetic approach takes a first step towards evaluating the potential risk associated with the
18 *B. cereus* Group strains. It can in the future be extended by the presence or production of pathogenic
19 markers. To date, the *ces* gene is the most relevant marker for the emetic syndrome. The toxins Nhe and
20 HlyII have been directly or indirectly linked to the diarrheal syndrome (3, 19), but they do not yet explain the
21 pathogenicity of all strains. Some new associated-virulence factors (5) have not been evaluated as pathogenic
22 makers yet, and some other are doubtlessly still unknown. Further knowledge of virulence-associated factors
23 secreted by the *B. cereus* Group strains, is thus currently needed for a more accurate detection of food
24 poisoning strains.

25

26 **ACKNOWLEDGMENTS**

27 We thank the “Direction Generale de l’Alimentation, Ministère de l’Agriculture”, INRA and AFSSA, for
28 financial support under the projects “Aliment Qualité Sécurité 2000-R0013” and “Tiacer”.

29 We are grateful to Caroline Delseny and Sylvain Courtillier for their skilful contributions.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

REFERENCES

1. **Agata, N., M. Ohta, M. Mori, and M. Isobe.** 1995. A novel dodecadepsipeptide, cereulide, is an emetic toxin of *Bacillus cereus*. *FEMS Microbiology Letters* **129**:17-20.
2. **Beecher, D. J., J. L. Schoeni, and A. C. Wong.** 1995. Enterotoxigenic activity of hemolysin BL from *Bacillus cereus*. *Infection & Immunity* **63**:4423-8.
3. **Cadot, C., S. Tran, M. L. Vignaud, M. L. Debuyser, A. B. Kolsto, A. Brisabois, C. Nguyen-The, D. Lereclus, M. H. Guinebretière, and N. Ramarao.** 2010. InhA1, NprA, and HlyII as candidates for markers to differentiate pathogenic from nonpathogenic *Bacillus cereus* strains. *Journal of Clinical Microbiology* **48**:1358-1365.
4. **Choma, C., M. H. Guinebretiere, F. Carlin, P. Schmitt, P. Velge, P. E. Granum, and C. Nguyen-The.** 2000. Prevalence, characterization and growth of *Bacillus cereus* in commercial cooked chilled foods containing vegetables. *Journal of Applied Microbiology* **88**:617-625.
5. **Clair, G., S. Roussi, J. Armengaud, and C. Duport.** 2010. Expanding the known repertoire of virulence factors produced by *Bacillus cereus* through early secretome profiling in three redox conditions. *Molecular and Cellular Proteomics* **9**:1486–1498.
6. **Ehling-Schulz, M., B. Svensson, M. H. Guinebretiere, T. Lindback, M. Andersson, A. Schulz, M. Fricker, A. Christiansson, P. E. Granum, E. Martlbauer, C. Nguyen-The, M. Salkinoja-Salonen, and S. Scherer.** 2005. Emetic toxin formation of *Bacillus cereus* is restricted to a single evolutionary lineage of closely related strains. *Microbiology* **151**:183-197.
7. **Ehling-Schulz, M., N. Vukov, A. Schulz, R. Shaheen, M. Andersson, E. Märtilbauer, and S. Scherer.** 2005. Identification and partial characterization of the nonribosomal peptide synthetase gene responsible for cereulide production in emetic *Bacillus cereus*. *Applied and Environmental Microbiology* **71**:105-113.
8. **Granum, P. E.** 1994. *Bacillus cereus* and its toxins. *Journal of Applied Bacteriology Supplementary* **23**:61S-66S.
9. **Granum, P. E., K. O'Sullivan, and T. Lund.** 1999. The sequence of the non-haemolytic enterotoxin operon from *Bacillus cereus*. *FEMS Microbiology Letters* **177**:225-9.
10. **Guinebretière, M.-H., F. L. Thompson, A. Sorokin, P. Normand, P. Dawyndt, M. Ehling-Schulz, B. Svensson, V. Sanchis, C. Nguyen-The, M. Heyndrickx, and P. De Vos.** 2008. Ecological diversification in the *Bacillus cereus* Group. *Environmental Microbiology* **10**:851-865.

- 1 11. **Guinebretiere, M. H., V. Broussolle, and C. Nguyen-The.** 2002. Enterotoxigenic profiles
2 of food-poisoning and food-borne *Bacillus cereus* strains. *Journal of Clinical Microbiology*
3 **40**:3053-3056.
- 4 12. **Guinebretiere, M. H., A. Fagerlund, P. E. Granum, and C. Nguyen-the.** 2006. Rapid
5 discrimination of *cytK-1* and *cytK-2* genes in *Bacillus cereus* strains by a novel duplex PCR
6 system. *FEMS Microbiology Letters* **259**:74-80.
- 7 13. **Guinebretiere, M. H., and C. Nguyen-The.** 2003. Sources of *Bacillus cereus*
8 contamination in a pasteurized zucchini puree processing line, differentiated by two PCR-
9 based methods. *FEMS Microbiology Ecology* **43**:207-215.
- 10 14. **Han, C. S., G. Xie, J. F. Challacombe, M. R. Altherr, S. S. Bhotika, D. Bruce, C. S.**
11 **Campbell, M. L. Campbell, J. Chen, O. Chertkov, C. Cleland, M. Dimitrijevic, N. A.**
12 **Doggett, J. J. Fawcett, T. Glavina, L. A. Goodwin, K. K. Hill, P. Hitchcock, P. J.**
13 **Jackson, P. Keim, A. R. Kewalramani, J. Longmire, S. Lucas, S. Malfatti, K.**
14 **McMurry, L. J. Meincke, M. Misra, B. L. Moseman, M. Mundt, A. C. Munk, R. T.**
15 **Okinaka, B. Parson-Quintana, L. P. Reilly, P. Richardson, D. L. Robinson, E. Rubin,**
16 **E. Saunders, R. Tapia, J. G. Tesmer, N. Thayer, L. S. Thompson, H. Tice, L. O.**
17 **Ticknor, P. L. Wills, T. S. Brettin, and P. Gilna.** 2006. Pathogenomic sequence analysis
18 of *Bacillus cereus* and *Bacillus thuringiensis* isolates closely related to *Bacillus anthracis*.
19 *Journal of Bacteriology* **188**:3382-3390.
- 20 15. **Kamat, A. S., and C. K. K. Nair.** 1992. Evidence for plasmid-mediated toxin production in
21 *Bacillus cereus* BIS-59. *World Journal of Microbiology and Biotechnology* **8**:210-211.
- 22 16. **Lapidus, A., E. Goltsman, S. Auger, N. Galleron, B. Ségurens, C. Dossat, M. L. Land,**
23 **V. Broussolle, J. Brillard, M. H. Guinebretiere, V. Sanchis, C. Nguyen-the, D. Lereclus,**
24 **P. Richardson, P. Winker, J. Weissenbach, S. D. Ehrlich, and A. Sorokin.** 2008.
25 Extending the *Bacillus cereus* group genomics to putative food-borne pathogens of different
26 toxicity. *Chemico-Biological Interactions* **171**:236-249.
- 27 17. **Lund, T., M. L. De Buyser, and P. E. Granum.** 2000. A new cytotoxin from *Bacillus*
28 *cereus* that may cause necrotic enteritis. *Molecular Microbiology* **38**:254-261.
- 29 18. **Lund, T., and P. E. Granum.** 1996. Characterisation of a non-haemolytic enterotoxin
30 complex from *Bacillus cereus* isolated after a foodborne outbreak. *FEMS Microbiology*
31 *Letters* **141**:151-156.
- 32 19. **Moravek, M., R. Dietrich, C. Buerk, V. Broussolle, M. H. Guinebretiere, P. E. Granum,**
33 **C. Nguyen-the, and E. Märtilbauer.** 2006. Determination of toxic potential of *Bacillus*
34 *cereus* isolates by quantitative enterotoxin analyses. *FEMS Microbiology Letters* **257**:293-
35 298.

- 1 20. **Thorsen, L., B. M. Hansen, K. F. Nielsen, N. B. Hendriksen, R. K. Phipps, and B. B.**
2 **Budde.** 2006. Charaterization of emetic *Bacillus weihenstephanensis*, a new cereulide-
3 producing bacterium. Applied and Environmental Microbiology **72**:5118-5121.
- 4 21. **Vassileva, M., K. Torii, M. Oshimoto, A. Okamoto, N. Agata, K. Yamada, T.**
5 **Hasegawa, and M. Ohta.** 2007. A new phylogenetic cluster of cereulide-producing
6 *Bacillus cereus* strains. Journal of Clinical Microbiology **45**:1274-1277.
- 7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

1 **Table 1.** Characteristics of the studied strains, including (A) the phylogenetic groups to which they belong in the *B. cereus* Group, previously defined (10)
 2 and (B) the distribution of the known food poisoning toxin genes among these groups (this study).
 3

Phylogenetic group	A				B					
	Species	subgroup	Other designation	Range of growth T°C	No. of tested strains	<i>hbl</i>	% of strains carrying :			<i>ces</i>
						<i>hbl</i>	<i>cytK-2</i>	<i>cytK-1</i>	<i>nhe</i>	
I	<i>B. pseudomycooides</i>	BC10	I-1	10-43	17	41	0	0	100 ^a	0
		BC13	I-2		7	86	0			
II	<i>B. cereus</i> II, <i>B. thuringiensis</i> II	BC06	II	7-40	31	61	13	0	100	0
III	<i>B. cereus</i> III <i>B. thuringiensis</i> III,	BC12	III-1	15-45	15	67	73			0
		BC05	III-2		26	12	31			31
		BC09	III-3		14	14	57	0	100	7
		BC08	III-4		28	14	39			0
IV	<i>B. cereus</i> IV, <i>B. thuringiensis</i> IV	BC04	IV-1	10-45	34	97	79			
		BC03	IV-2		32	97	97	0	100	0
		BC07	IV-3		29	86	79			
V	<i>B. cereus</i> V, <i>B. thuringiensis</i> V	BC11	V	8-40	17	88	6	0	100	0
VI	<i>B. weihenstephanensis</i> <i>B. mycooides</i> , <i>B. thuringiensis</i> VI	BC01	VI-1	5-37	93	83	0	0	100	0
		BC02	VI-2		43	60	0		100	
VII	' <i>B. Cytotoxicus</i> ' sp. nov.	BC14	VII	20-50	5	0	0	100	100	0

4 ^a In group I, nine strains were tested for *nhe* genes instead of 24 strains




Figure 1. Cytotoxic activity of *B. cereus* culture filtrates against Caco2 cells.

1a, Data distribution of the global population (*B. cereus* Group) is represented by the first Box plot ('All data') whereas that into each sub-population (phylogenetic group II to VII) by the next box plots ('II' to 'VII'). 1b, Detailed results for subgroups in the phylogenetic groups III and IV.