



**HAL**  
open science

## Culture-independent methods for identifying microbial communities in cheese

Jean-Luc Jany, Georges Barbier

► **To cite this version:**

Jean-Luc Jany, Georges Barbier. Culture-independent methods for identifying microbial communities in cheese. *Food Microbiology*, 2008, 25 (7), pp.839-848. 10.1016/j.fm.2008.06.003 . hal-00561679

**HAL Id: hal-00561679**

**<https://hal.univ-brest.fr/hal-00561679v1>**

Submitted on 1 Feb 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

---

1 **Culture-Independent Methods for Identifying Microbial**  
2 **Communities in Cheese**

---

3  
4 **Jean-Luc Jany\* and Georges Barbier**

5  
6 Laboratoire Universitaire de Biodiversité et Ecologie Microbienne

7 Université Européenne de Bretagne - ESMISAB

8 Parvis Blaise Pascal

9 Technopôle de Brest Iroise

10 Plouzané, France

11

12

13

14

15

16 \* corresponding author :

17 Jean-Luc Jany

18 phone: + 33 2 98 05 61 26

19 e-mail: [jean-luc.jany@univ-brest.fr](mailto:jean-luc.jany@univ-brest.fr)

20

21

22

23

24

1 **Abstract**

2

3 This review focuses on the culture-independent methods available for the description of both  
4 bacterial and fungal communities in cheese. Important steps of the culture-independent  
5 strategy, which relies on bulk DNA extraction from cheese and polymerase chain reaction  
6 (PCR) amplification of selected sequences, are discussed. We critically evaluate the  
7 identification techniques already used for monitoring microbial communities in cheese,  
8 including PCR-denaturing gradient gel electrophoresis (PCR-DGGE), PCR- temporal  
9 temperature gradient gel electrophoresis (PCR-TTGE) or single-strand conformation  
10 polymorphism-PCR (SSCP-PCR) as well as some other techniques that remain to be adapted  
11 to the study of cheese communities. Further, our analysis draws attention to the lack of data  
12 available on suitable DNA sequences for identifying fungal communities in cheese and  
13 proposes some potential DNA targets.

14

## 1 **1. Microbial communities in cheese**

2

3 Cheese is produced throughout the world and more than 1000 varieties (Sandine and Elliker  
4 1970) with different forms and flavours exist. Cheesemaking is conjectured to date back to  
5 some 8000 years ago originating from the Middle East (Fox et al. 2000) where the first  
6 fermented milk-based foods were made. Four basic ingredients are required to produce most  
7 cheeses: milk, rennet, salt and microorganisms. These four ingredients are processed through  
8 different steps such as acidification, coagulation, syneresis and ripening (see Fox et al. 2000).  
9 Each unique combination of ingredients and processing parameters leads to a specific type of  
10 cheese with unique properties. Fox et al. (2000) mentioned how fascinating it is that “such a  
11 diverse range of products can be produced” from “basically similar raw material”. The  
12 composition and activity of the microflora is the least controllable of all the parameters. The  
13 microflora is made up of (i) starter lactic acid bacteria that are involved in acid production  
14 during cheese manufacture and that also participate in the ripening processes to various  
15 extents; and (ii) non-starter lactic acid bacteria, other bacteria, yeasts and filamentous fungi  
16 that form the secondary microflora, which plays a significant role during ripening (Fox et al.  
17 2000). Both starter and secondary flora modify the physical and chemical properties of  
18 cheese, contributing to and reacting to environmental changes that occur during the  
19 manufacture and ripening of cheese. Coppola et al. (2007) distinguished seven technological  
20 production phases of cheese that may constitute selective pressures for the microbial species  
21 that play an important role in community succession occurring throughout cheese  
22 manufacture and ripening. The fundamental features that influence the dynamics of the  
23 cheese ecosystem have been described by Beresford et al. (2001). They include (i) physical  
24 features such as moisture, salt concentration, pH or redox potential which change during  
25 cheese manufacture and are themselves influenced by the microflora, and (ii) biological

1 features such as those resulting from the interactions between microorganisms. The  
2 characteristics of a given cheese therefore depend on microflora dynamics. Although food  
3 microflora is undoubtedly not as diverse as environmental microflora, such as those found in  
4 the soil (Garbeva et al. 2004), cheese microbial populations still remain difficult to control  
5 due to their complex dynamics and to their interactions (Beresford et al. 2001). Knowledge of  
6 the structure and dynamics of the whole microbial community of cheese would promote  
7 better understanding of how cheese characteristics vary with respect to microbial growth and  
8 metabolism. For instance, greater control over microflora composition would make it  
9 possible to better select for specific organoleptic properties or to prevent quality defects or  
10 spoilage. For these reasons, cheese bacterial and fungal communities have already been  
11 partially identified using traditional methods and, to a lesser extent, molecular techniques.

12

## 13 **2. Towards culture-independent methods**

14

15 As in other fields in microbiology, species identification in cheese can be assessed through  
16 the use of either culture-dependent or culture-independent methods. Culture-dependent  
17 methods consist of isolating and culturing microorganisms prior to their identification  
18 according to either morphological, biochemical or genetic characteristics. Different cultures  
19 can even be bulked and analysed using global analysis methods, such as those as described  
20 below for culture-independent methods (Ercolini et al. 2001; 2004). These methods have  
21 already shed light on the structure of microbial populations during cheese manufacture  
22 (Andrighetto et al. 1998; Fitzsimmons et al. 1999; Mannu et al. 2000; Berthier et al. 2001;  
23 Dasen et al. 2003). However, culture-dependent methods are time-consuming, due to long  
24 culture periods and elaborate culture techniques. They are therefore not amenable to  
25 monitoring community dynamics during cheese manufacture and ripening. Moreover, species

1 occurring in low numbers are often out-competed *in vitro* by numerically more abundant  
2 microbial species (Hugenholtz et al. 1998) and some species may be unable to grow *in vitro*  
3 (Ward et al. 1990, 1992; Head et al. 1998). Hence, if culture conditions are poor and the  
4 number of isolates too low, the culture collection will not be representative of the community  
5 and the actual microbial diversity will be misinterpreted.

6         Community-level studies are relying more and more on culture-independent methods  
7 based on the direct analysis of DNA (or RNA) without any culturing step. These methods are  
8 based on protocols where total DNA (or RNA) is directly extracted from the substrate.  
9 Coupled with a global analysis, these methods make it possible to study the total diversity  
10 from the bulk extract in a single step. As they are fast and potentially more exhaustive, these  
11 methods are well suited for analysing microbial communities over time and may provide the  
12 possibility of exploring cheese microflora dynamics in detail. Most of these methods use  
13 polymerase chain reaction (PCR) amplification of total DNA. The PCR amplicons from  
14 different species are discriminated by using gel or capillary separation or by hybridization to  
15 specific probes (**Figure 1**). However, these methods have potential biases, which will be  
16 discussed below.

17

### 18 **3. 'Pick'em all'**

19

20 As previously mentioned, the isolation step of culture-dependent methods introduces biases  
21 because some species are unable to grow under the selected experimental conditions. Culture-  
22 independent methods typically aim at collecting DNA from the whole community to  
23 overcome this bias. Nevertheless, technical issues may arise: DNA may not be recovered  
24 from all genotypes or PCR amplification may be inaccurate. Some genotypes may remain  
25 undetected due to low species abundance in the substrate, low species availability due to

1 insufficient homogenization of the matrix, inadequate cell lysis that prevents release of  
2 nucleic acids, or inhibition of PCR amplification.

3 Complete homogenization of the cheese matrix can be challenging and partial  
4 homogenization may hamper cell lysis, lowering DNA availability. Therefore, existing  
5 protocols for DNA extraction from cheese include mechanical homogenization in salt  
6 solutions such as trisodium-citrate (Coppola et al. 2006; Parayre et al. 2007). However, these  
7 protocols may have to be adapted to the cheese under investigation. The extraction step is  
8 usually performed in specific buffers, such as phosphate buffers containing various  
9 detergents. Extraction can also be done with guanidine thiocyanate and *N*-laurylsarcosine  
10 (Callon et al. 2006; Duthoit et al. 2005a,b; Delbès et al. 2007; Le Bourhis et al. 2007) and/or  
11 lytic enzymes such as pronase (Florez and Mayo 2006; El Baradei et al. 2007) or proteinase  
12 K (Ercolini et al. 2003; Cocolin et al. 2007). Lysis can be mechanically improved by  
13 vigorously shaking samples in tubes containing beads (Randazzo et al. 2002; Duthoit et al.  
14 2003, Duthoit et al. 2005a,b; Feurer et al. 2004a,b; Delbès and Montel 2005; Florez and  
15 Mayo 2006; Delbès et al. 2007; Le Bourhis et al. 2007). New commercially available  
16 automated instruments exploiting different techniques, such as pressure cycling technology,  
17 which generates alternating hydrostatic pressure (Tao et al. 2007), or optimized mechanical  
18 grinding, should be tested for their capacity to improve DNA extraction from microorganisms  
19 in cheese.

20 Like for other food materials or environmental samples, DNA extraction yield and  
21 PCR sensitivity are significantly reduced by a wide range of inhibiting substances (Wilson  
22 1997). DNA extraction efficiency may be decreased by the high quantities of  
23 macromolecules in cheeses, such as casein or lipids, that can adsorb detergents, chaotropic or  
24 chelating agents necessary for nucleic acid extraction (Bonaïti et al. 2006). Moreover, even if  
25 DNA yield is high, inhibitors that have not been eliminated may lower PCR sensitivity

1 (Wilson 1997). For example, Drake et al. (1996) reported that protein breakdown products  
2 decreased the sensitivity of PCR amplification of rRNA genes of Lactobacillus spp.. DNA  
3 solutions are therefore often purified using phenol-chloroform, which eliminates proteins and  
4 various remnants (Randazzo et al. 2002, 2006; Cocolin et al. 2004; Feurer et al. 2004a,b;  
5 Callon et al. 2006; Delbès and Montel 2005; Duthoit et al. 2003, 2005a,b; Delbès et al. 2007;  
6 Le Bourhis et al. 2007). Several authors have used commercial DNA extraction kits based on  
7 ion-exchange chromatography, size-exclusion or sorption techniques (Ercolini et al. 2001;  
8 Rademaker et al. 2005) or just for DNA purification (Parayre et al. 2007).

9         It is noteworthy that most studies optimized DNA extraction to target bacterial DNA,  
10 with the exception of Callon et al. (2006), who evaluated the dynamics of yeasts, and Florez  
11 and Mayo (2006), who investigated the dynamics of the dominant microorganisms including  
12 bacteria, yeasts and filamentous fungi. In order to characterise whole microbial communities,  
13 protocols have to be adapted to extract DNA from all different types of microorganisms, as  
14 attempted by Bonaïti et al. (2006).

15         Most culture-independent methods rely on PCR amplification of a targeted sequence.  
16 This step can be affected by preferential or differential PCR amplification (Reysenbach et al.  
17 1992; Walsh et al. 1992; Ercolini 2003; Kanawaga 2003) that may hinder the detection of  
18 some genotypes when analysing bulk DNA extracted from cheese. Preferential PCR  
19 amplification can be caused by (i) primer mismatches at the annealing sites of the DNA  
20 templates of some genotypes or (ii) a lower rate of primer hybridization to certain templates  
21 due to differential denaturation of these templates (Walsh et al. 1992; Suzuki and Giovannoni  
22 1996). Systematically testing different sets of primers and enhancing DNA denaturation  
23 during PCR by using different reagents (denaturants and cosolvents) may solve these  
24 problems (Reysenbach et al. 1992; Weissensteiner and Lanchbury 1996; Hansen et al. 1998).  
25 A second type of bias that may affect PCR carried out on complex bulk DNA extracts is the



1 occurrence of (i) heteroduplexes that arise in later PCR cycles when primer concentration  
2 decreases and the concentration of PCR products is high enough to compete with the primers  
3 for annealing (Kanawaga 2003) and (ii) chimeric amplicons that also form in later PCR  
4 cycles when the concentration of incompletely extended primers is high enough to compete  
5 with the original primer for annealing, or when template concentration is high enough to  
6 allow the re-annealing of templates before or during primer extension (Kanawaga 2003). All  
7 these artefacts can generate additional signals that do not correspond to genotypes in the  
8 sample. These artefacts can be minimized by using a low number of PCR cycles.

9

#### 10 **4. Analysing bulk DNA**

11

12 Through the construction of clone libraries from PCR products amplified from bulk cheese  
13 DNA and subsequent sequencing of the different clones, it is theoretically possible to  
14 examine the actual diversity of a given community. Although the cost of cloning and  
15 sequencing is decreasing, this strategy remains expensive if used on a routine basis since it  
16 implies large-scale cloning and sequencing to ensure that the community is exhaustively  
17 screened. This strategy has been used for the study of bacterial cheese communities (Feurer et  
18 al. 2004a,b; Delbès et al. 2007; El Baradei et al. 2007), but only as a complement to other  
19 techniques (**Table 1**). Amongst the other molecular methods that may allow exhaustive  
20 screening of microbial communities with no requirement for cloning PCR products, the most  
21 commonly used in cheese microbiology are based on gel electrophoresis or chromatography.  
22 The principal techniques that can be used to describe microbial communities in cheese are  
23 briefly discussed below.

24

#### 1 **4.1. PCR-denaturing gradient gel electrophoresis and PCR-temporal temperature** 2 **gradient gel electrophoresis**

3  
4 PCR-denaturing gradient gel electrophoresis (PCR-DGGE; Myers 1987) and PCR-temporal  
5 temperature gradient gel electrophoresis (PCR-TTGE; Yoshino et al. 1991) (collectively  
6 referred to here as PCR-DG/TTGE) are based on the separation of PCR amplicons of the  
7 same size but with different sequences. In a denaturing acrylamide gel, DNA partially  
8 denatures in discrete regions called melting domains. The melting temperature of these  
9 domains is sequence specific. When the melting temperature ( $T_m$ ) of the lowest melting  
10 domain is reached, the DNA is partially denatured creating branched molecules. This  
11 branching reduces DNA mobility in the gel. Therefore, amplicons of the same size but with  
12 different nucleotide compositions can be separated based on differences in the melting  
13 behaviour of their melting domains. For PCR-DGGE, the denaturing conditions rely on the  
14 use of chemical denaturants (formamide and urea) incorporated into an acrylamide gel as a  
15 linear denaturing gradient. PCR-DGGE electrophoresis is carried out at constant temperature,  
16 typically between 55°C and 65°C (Ercolini 2004). For PCR-TTGE, the denaturing gradient is  
17 obtained by varying the temperature over time without chemicals, thus generating more  
18 reproducible data. PCR-DG/TTGE provides optimal resolution when PCR products are not  
19 completely denatured. Thus, adding a so-called GC clamp (30-40bp) to one of the PCR  
20 primers ensures that the PCR products are not completely denatured during the analysis  
21 (Myers et al. 1985a,b; Sheffield et al. 1989). Prior to PCR-DG/TTGE analysis, the location of  
22 the different melting domains within a DNA sequence can be predicted *in silico*, allowing  
23 selection of the best primer pair. A PCR-DG/TTGE database containing migration profiles  
24 corresponding to reference strains is generally created (Ogier et al. 2002, 2004) to facilitate  
25 further identification of PCR-DG/TTGE profiles. A molecular ladder can be constructed by

1 using amplicons corresponding to representative species of this database and used in each gel  
2 migration. However, this kind of database cannot be exhaustive and representative of the  
3 actual community analysed and usually requires constant updates. The advantage of PCR-  
4 DG/TTGE is that amplicons can be directly extracted from the DG/TTGE acrylamide gel and  
5 sequenced. Unidentified profiles that do not match reference profiles can thus be sequenced,  
6 compared to publicly available sequence databases and appended to the PCR-DG/TTGE  
7 profile database. The first PCR-DGGE application to microbiology was done by Muyzer et  
8 al. (1995) who studied bacterial communities of deep-sea hydrothermal vents. PCR-  
9 DG/TTGE are the molecular techniques that have been the most extensively used for the  
10 study of microbial communities in dairy products (e.g. Lafarge et al. 2004; Ogier et al. 2002;  
11 2004) and more specifically to monitor the structure and even dynamics of microbial  
12 communities in cheese (see references in **Table 1**).

13

#### 14 **4.2. Single-strand conformation polymorphism-PCR**

15

16 Single-strand conformation polymorphism-PCR (SSCP-PCR; Orita et al. 1989; Ravnik-  
17 Glavac et al. 1994) is a technique using either acrylamide gel- or capillary-based automated  
18 sequencer, based on the separation of denatured (single-stranded) PCR products. Under non-  
19 denaturing conditions, single-stranded DNA folds into tertiary structures according to their  
20 nucleotide sequences and their physicochemical environment (e.g., temperature and ion  
21 strength). This causes differences in electrophoretic mobility in non-denaturing gels. SSCP-  
22 PCR is potentially easier to carry out than PCR-DG/TTGE since there is no need for gradient  
23 gels or use of GC-clamp primers and it can be performed using an automated sequencer.  
24 However, when using an automated sequencer, one of the disadvantages of this technique lies  
25 in the difficulty of appending new data to an existing database: samples presenting unknown

1 profiles cannot be directly sequenced because they are labelled. SSCP-PCR is the second  
2 most-used method to study microbial communities of cheese (see references in **Table 1**).

3

#### 4 **4.3. Terminal restriction fragment length polymorphism**

5

6 Terminal restriction fragment length polymorphism (T-RFLP; Liu et al. 1997) is based on  
7 digestion of fluorescent end-labelled PCR products with restriction endonucleases. Either one  
8 or both 5' and 3' ends of the amplicon can be labelled by incorporating a dye on either one or  
9 both PCR primers. The digested products are separated by electrophoresis using either  
10 acrylamide gel- or capillary-based automated sequencer, with laser detection of the labelled  
11 fragments. This system only detects the end-labelled terminal restriction fragments (TRFs) of  
12 the digested PCR products and their size can be calculated based on the use of DNA size  
13 standards that are run simultaneously with the samples. The data consists of the sizes of the  
14 PCR amplicons that contain the labelled primer and are observed as electrophoregram peaks  
15 or gel bands. Variation in the presence and location of the restriction sites results in different  
16 genotypes having different TRF lengths. T-RFLP was initially developed as a fingerprinting  
17 technique (Liu et al. 1997), with the number of TRFs used as an indication of biodiversity.  
18 The use of a database of TRF profiles obtained from reference samples allows the  
19 identification of the different species of a given community (Dickie et al. 2002). As for  
20 SSCP-PCR performed using an automated sequencer, samples that present unknown profiles  
21 cannot be directly sequenced because they are cut and labelled. T-RFLP has been used to  
22 study diverse microbial communities (e.g. Liu et al. 1998) and has been extensively used by  
23 mycologists since this method is reportedly more sensitive than PCR-DG/TGGE for fungi  
24 (Brodie et al. 2003). However, some limitations—due in particular to inefficient restriction  
25 enzyme cleavage—have been pointed out (Avis et al. 2006). T-RFLP was also found to be an

1 excellent tool for culture-independent assessment of bacterial community structure and  
2 dynamics during ripening of cheese (Rademaker et al. 2005).

3

#### 4 **4.4. Denaturing high-performance liquid chromatography**

5

6 Denaturing high-performance liquid chromatography (DHPLC) (Oefner and Underhill 1995  
7 in Xiao and Oefner 2001) allows separation of amplicons using an ion-pair reversed-phase  
8 high-performance liquid chromatography (IP RP HPLC) automated detection system. It was  
9 initially used to detect single nucleotide polymorphisms (SNPs) in clinical applications (e.g.,  
10 Frueh and Noyer-Weidner 2005) based on the analysis of heteroduplex formation. DHPLC is  
11 a promising approach for microbial community analysis (Barlaan et al. 2005). PCR  
12 amplicons are injected into a chromatography column containing alkylated non-porous  
13 polystyrene/polydivinylbenzene particles. Separation of the different amplicons relies on the  
14 elution of partially denatured PCR products, which is achieved with the intervention of the  
15 ion-pairing agent, triethylammonium acetate (TEAA) and the cartridge matrix of the system.  
16 As described in Barlaan et al. (2005), under partial denaturation and a given flow rate of  
17 gradient buffers, amplicons of the same size but with different melting behaviours due to  
18 different nucleotide compositions will have different retention times. These differences are  
19 due to the reduced negative charges in the single-stranded portions of the partially denatured  
20 PCR products compared to the double-stranded molecules. The negative charges interact with  
21 the positive charges of ammonium ions of TEAA, and double-stranded DNA is more  
22 efficiently adsorbed to the stationary phase in the cartridge. DHPLC permits high-throughput  
23 automated analyses and, unlike SSCP-PCR or T-RFLP, it allows the collection of elution  
24 fractions corresponding to different amplicons that can be directly sequenced even more  
25 easily than with PCR-DG/TTGE methods. Protocols have already been developed for

1 analysing marine bacterial populations (Barlaan et al. 2005), monitoring intestinal microflora  
2 (Goldenberg et al. 2007) and, recently, also for studying the bacterial diversity occurring in  
3 natural whey cultures used for cheese manufacture (Ercolini et al., 2008). The latter study  
4 showed that DHPLC technique was at least as effective as the widely used PCR-DGGE  
5 technique in assessing species diversity of food-related microbial communities.

#### 7 **4.5. DNA microarrays**

8  
9 The complementary DNA (cDNA) microarray (or microchips) technology has dramatically  
10 changed the way gene expression can be assessed (Duggan et al. 1999). Since Guschin et al.  
11 (1997) introduced the DNA microarray approach to microbial community analysis, the DNA  
12 array-based methods hold great promise for more extensive analyses of microbial  
13 communities (Zhou et al. 2003; Bodrossy et al. 2007; Wagner et al. 2007) and potential  
14 applications cover various fields of microbiology, including food science (Bae et al. 2005).  
15 By using DNA microarrays, the identification of labelled PCR products or directly retrieved  
16 RNA relies on their hybridization to oligonucleotide probes attached to a substrate. A  
17 description of the various types of microarrays is given by Zhou et al. (2003). Unlike the  
18 previously described methods, DNA array technology potentially allows the simultaneous  
19 application of a nearly unlimited number of probes in a single hybridization experiment  
20 (Small et al. 2001; Peplies et al. 2003). This technique is therefore very well suited to even  
21 the most complex environmental samples. However, for this approach to work, each probe  
22 must specifically hybridize, under given stringency conditions, to a fully matched DNA  
23 target (Valinsky et al. 2002) which has been proven very difficult (Wagner et al. 2007).  
24 Moreover, the design and refinement of efficient probes depend on the comprehensiveness  
25 and quality of probe target database (Wagner et al. 2007). The low quality of some annotated

1 sequences in the available databases complicates probe design. Nonetheless, given the lower  
2 diversity of food-borne communities (compared to environmental communities) and given  
3 the unequalled high-throughput capacity of DNA microarrays, it would be worthwhile to  
4 devote research efforts for developing a robust probe target sequence database necessary for  
5 using DNA microarrays as a routine tool for monitoring cheese microbial communities  
6 throughout cheese manufacture.

7

## 8 **5. What is a good target?**

9

10 The discrimination between the different species from a cheese sample depends on the  
11 capacity of the analytical technique to reveal polymorphisms. However, whatever analytical  
12 technique is used, no discrimination is possible without polymorphism. The target DNA  
13 sequence must therefore be variable between species. In addition, since almost all of the  
14 above-described techniques rely on PCR amplification, the target DNA sequence must also  
15 include conserved sequences that can serve as anchors for PCR primers.

16 Unlike prokaryotic genes that do not usually harbour introns, fungal genes are  
17 interrupted by non-coding introns. Although introns are much more variable than exons and  
18 thus offer high levels of polymorphism between species, their length and their location vary  
19 greatly beyond the family level (James et al. 2006). It is therefore difficult to design primers  
20 that allow PCR amplification of introns from all the species in the community. Moreover, the  
21 analysis of exons is often discriminative enough to distinguish between species since they  
22 show sufficient variation at the interspecific level, especially at their third codon positions  
23 that evolve at rates similar to introns. Consequently, most of the DNA regions selected for  
24 studying fungal communities are located in exons.

25

## 1 **5.1 Bacterial and fungal ribosomal DNA**

2

3 Since Woese and Fox's paper (1977), most community surveys have focused on RNA genes  
4 and intergenic spacers. Bacterial 16S, 23S and 5S rRNA genes are organized into a co-  
5 transcribed operon. From one to as many as 15 copies of the operon may be dispersed in a  
6 bacterial genome (Klappenbach et al. 2000). In contrast, in fungi and more generally in  
7 eukaryotes, 18S, 28S 5.8S and 5S rRNA genes and intergenic spacers are tandemly repeated  
8 (up to hundreds of repeats) (Cihlar and Sypherd 1980). The 5S rRNA is, in some species,  
9 separated from the main unit (Kellog and Appels 1995). These repeated units are organized  
10 as multiple and dispersed arrays at chromosomal nucleolar organizer regions (NORs). In both  
11 prokaryotes and eukaryotes, rRNA genes usually show extraordinary sequence homogeneity  
12 within a species (Elder and Turner 1995; Liao 1999). This sequence homogenization is likely  
13 to be due to a process known as concerted evolution. The underlying mechanisms of  
14 concerted evolution involve recombination processes collectively referred to as mitotic or  
15 somatic recombination (see Liao 2000) including unequal crossing-over and gene conversion  
16 that are responsible for DNA repair during replication and transcription. The paradigm of  
17 concerted evolution makes it possible to treat repeated rRNA genes as one locus. For this  
18 reason, rRNA genes have been widely used in phylogeny reconstructions and species  
19 identification. However, deviations from this expected homogeneity of rDNA sequences have  
20 been detected in many taxa.

21 Although multiple-gene investigation has already been recommended for bacterial  
22 species identification (Stackebrandt et al. 2002), studies evaluating bacterial diversity and/or  
23 dynamics or investigating bacterial phylogenetics have mostly focused only on ribosomal  
24 DNA array analysis (Throbäck et al. 2004). In cheese, all bacterial community surveys are  
25 based on the analysis of 16S rRNA genes and 16S-23S intergenic region (**Table 1**). Bacterial



1 16S rRNA genes comprise nine hypervariable regions, V1-V9, that exhibit considerable  
2 sequence diversity among species (Van de Peer et al. 1996; Baker et al. 2003). These  
3 hypervariable regions are generally flanked by conserved sequences that can serve as anchors  
4 for universal or specific primer pairs (see Baker et al. 2003). They are therefore used for  
5 species identification and allow the evaluation of community diversity. No single region can  
6 differentiate among all bacteria and different regions can be used depending on the goal  
7 (Chakravorty et al. 2007). In cheese, the V3 hypervariable region is the target that has been  
8 the most extensively used, but the use of different sets of primers targeting different regions  
9 can improve the analysis (Delbès et al. 2007).

10 In fungi, the analysis of fungal rRNA genes limits identification to the genus or  
11 family level (Anderson and Cairney 2004). The fungal internal transcribed spacers (ITS)  
12 provide a greater taxonomic resolution than rRNA genes and are generally used for fungal  
13 community surveys in different environments (Anderson and Cairney 2004). The ITS is a  
14 region located between the 18S rRNA and 28S rRNA genes and including the 5.8S rRNA  
15 gene that splits the ITS into two parts: ITS1 and ITS2. The ITS region undergoes a faster rate  
16 of evolution than rRNA genes but its sequence remains homogenous within a species. Indeed,  
17 both ITS1 and ITS2 fulfil significant functions during rRNA maturation (Joseph et al. 1991;  
18 Liu and Schardl 1994) and are under selective pressure. In Penicillium spp. which are  
19 amongst the most prominent fungi in cheese, the analysis of ITS affords better discrimination  
20 than rRNA genes (Skouboe et al. 2000; Doaré-Lebrun et al. 2006). Nevertheless, Doaré-  
21 Lebrun et al. (2006) when analysing the fungal community in grape showed that the use of  
22 ITS was not sufficient for good interspecific discrimination, especially for the species of the  
23 subgenus Penicillium, a monophyletic group of moulds that represent 58 of the ≈250  
24 accepted species in the genus *Penicillium* (Seifert et al., 2007). It is therefore recommended

1 to use other targets in addition to ITS when analysing fungal communities in cheese where  
2 species of this subgenus are among the most prominent.

3

## 4 **5.2. Other fungal genes**

5

6 With the exception of rRNA genes and ITS, only a few number of sequences have been used  
7 for studying fungal communities. Recently, Seifert et al. (2007) tested the reliability of using  
8 the *COI* gene that codes for mitochondrial cytochrome *c* oxidase 1 as a barcoding tool for  
9 Penicillium spp. identification. This gene has already been used extensively in animal  
10 barcoding and could be used as a new marker to investigate fungal communities. Sequence  
11 analysis of *COI* yielded a coherent phylogeny of the taxonomically challenging Penicillium  
12 subgenus (Seifert et al. 2007), suggesting that *COI* is a powerful tool for fungal barcoding.  
13 However, the results by these authors show that *COI* is not more variable among the species  
14 of the subgenus Penicillium than ITS. The low *COI* interspecific divergence precludes  
15 distinguishing species of the subgenus Penicillium using gel/capillary separation of PCR  
16 products.

17 Amongst other possible targets, genes that encode mitochondrial rRNA genes  
18 (Lutzoni et al. 2004) could represent an alternative to nuclear rRNA genes. Moreover, the  
19 general effort for inferring phylogeny of the kingdom fungi (Lutzoni et al. 2004; Blackwell et  
20 al. 2006; Hibbet et al. 2007) has yielded a high number of sequences for different genes and  
21 for different fungal species that can be tested for species identification. The non-ribosomal  
22 genes used for inferring fungal phylogenies comprise: (i) the genes *RPB1* and *RPB2* that  
23 encode the two largest subunits of the RNA polymerase II (James et al. 2006); the gene *EF-1*  
24  $\alpha$  coding for elongation factor 1-alpha (James et al. 2006); the gene *BenA* encoding  $\beta$ -tubulin  
25 A (Einax and Voigt 2003; Samson et al. 2004); the *GPD* gene coding for glyceraldehyde 3-

1 phosphate dehydrogenase (Berbee et al. 1999). All these potential targets should be tested for  
2 their usefulness for distinguishing between different species of fungi that compose fungal  
3 communities in cheese. According to Samson et al (2004) and Seifert et al (2007) *BenA* is  
4 more variable than ITS amongst the species of the Penicillium subgenus and could be a good  
5 target for analysing cheese fungal communities using methods based on gel/capillary analysis  
6 of PCR products. The most promising sequences to analyse can be identified by *in silico* data  
7 mining of publicly available databases. However, sequence availability for a large range of  
8 species varies amongst genes. Additional sequencing is likely to be required to cover a  
9 reasonable array of species. This requires selecting and designing universal or even a set of  
10 different primers to sequence the targeted genes in the different species investigated.

11

## 12 **6. Limitations and pitfalls of culture-independent methods**

13

14 As previously mentioned, the first limitation arises from the difficulty of accessing every  
15 genotype from the community: poor DNA extraction yield, PCR inhibition by various  
16 extraction by-products or by substances coming from the cheese matrix itself, and differential  
17 PCR amplification have already been reviewed (Wintzingerode et al. 1997; Ercolini 2004).  
18 We have already made some suggestions to address, when possible, these concerns. The  
19 various techniques have limitations in terms of resolution: PCR-DG/TTGE, SSCP-PCR, T-  
20 RFLP, DHPLC and even microarrays can generate patterns in which different genotypes  
21 group together due to co-migration/co-elution (Ogier et al. 2002; Feurer et al. 2004a; Lafarge  
22 et al. 2004; Delbès et al. 2007; El-Baradei et al. 2007) or ‘co-hybridization’ (Wagner et al.  
23 2007). Since they allow direct (cloning) sequencing of migrants/eluants, PCR-DG/TTGE and  
24 DHPLC may overcome the co-elution/co-migration problem more easily. Another limitation  
25 of gel/capillary migration-based methods is obtaining profiles in which the less-common

1 amplified sequences cannot be distinguished from background noise (Feurer et al. 2004a;  
2 Callon et al. 2006). This problem increases with the diversity of the community. Ideally,  
3 multiple DNA targets should be used, as suggested by Doaré-Lebrun et al. (2006) and Delbès  
4 et al. (2007), to increase the efficiency in discriminating between species. Another remark  
5 made by several authors (Ercolini et al 2001; Feuerer et al. 2004a,b; Florez and Mayo 2006;  
6 Delbès et al. 2007) is that culture-independent methods regularly fail to identify species  
7 obtained using culture-dependent methods. These two types of methods reveal different  
8 images of the same community. Therefore, those authors suggest that using a polyphasic  
9 approach, combining culture-dependent and culture-independent methods, may be  
10 worthwhile to obtain a more accurate view of the structure of the microbial community.  
11 Nevertheless, culture-independent methods have proven to be the only ones with the capacity  
12 for monitoring the rapid dynamics of microbial communities during cheese manufacture and  
13 ripening processes, where microorganisms encounter multiple environmental shifts. Culture-  
14 independent methods need to be improved to reveal as accurately as possible the actual  
15 microbial communities.

16

## 17 **7. Microbial community activity in cheese**

18 In order to reveal metabolically active populations, some authors performed analysis on  
19 reverse-transcribed RNA. Randazzo et al. (2002) and Duthoit et al. (2005a) were among the  
20 first to use reverse transcriptase-PCR-DGGE (RT-DGGE) and reverse transcriptase-PCR-  
21 SSCP (RT-PCR-SSCP), respectively. By combining RT-PCR-DGGE and PCR-DGGE or  
22 RT-PCR-SSCP and PCR-SSCP, these authors were able to differentiate the active component  
23 (rRNA-derived) from the total diversity (rDNA-derived) of the community. Duthoit et al.  
24 (2005b) even attempted to link population dynamics and activities (as assessed by PCR-  
25 SSCP and RT-PCR-SSCP) to sensorial characteristics in order to identify which species play

1 a major role in the development of the organoleptic properties of Salers cheese. However,  
2 their analysis failed to explain the diversity of sensorial qualities in Salers cheese.  
3 Nonetheless, combining DNA- and RNA-based analyses with more quantitative methods,  
4 such as DNA- and cDNA-microarrays coupled with chemiometric and/or sensorial tests,  
5 could significantly increase our ability to identify the impact of the microbial population on  
6 organoleptic characteristics.

7

## 8 **8. Perspectives in the field of cheese microbiology**

9 Our review draws attention to the fact that only a few studies (Florez and Mayo 2006; Callon  
10 et al. 2006; Gente et al. 2007) have targeted fungal communities of cheeses. However, yeasts  
11 and filamentous fungi play a vital role in the development of organoleptic characteristics  
12 and/or, on the contrary, may be the source of quality defects. Protocols need to be developed  
13 to optimize fungal DNA extraction from cheese and new target sequences, in addition to  
14 ribosomal loci, should be investigated to better discriminate between the different fungal  
15 species found in cheese. Surveys could then be extended to the whole microbial community.

16 It is noteworthy that, at present, some species can only be detected by culture and that  
17 polyphasic methods, including both culture-independent and culture-dependent approaches,  
18 are necessary to at least allow evaluation of the efficiency of these two types of approaches.  
19 Nevertheless, although culture-independent methods still fail to exhaustively describe the  
20 microbial community composition in cheese, these methods provide a much faster  
21 assessment of community composition than culture-dependent methods do. New light has  
22 been shed on population dynamics and culture-independent methods offer a powerful tool for  
23 controlling cheese manufacture. Therefore, it is worthwhile to devote efforts to improve the  
24 resolution of culture-independent methods and facilitate their transfer to cheese industry.

1           Because cheese microbial communities are much less complex than environmental  
2 communities (in terms of the number of species encountered), they offer an ideal system to  
3 test identification methods that require setting up reference databases or designing individual  
4 nucleic probes, such as for DNA microarrays. A major task is to select the best targets  
5 allowing both universal screening of the microorganisms in cheese and discrimination  
6 between taxa at the species level. Achieving this goal requires increasing the accessibility of  
7 high-throughput DNA sequencing technology that can provide digital images of cheese  
8 ecosystems (cheese ecogenome) by identifying genes associated with species and even  
9 functions.

10

## 11 **Acknowledgements**

12

13 This work was financially supported by Bretagne Biotechnologies Alimentaires and Région  
14 Bretagne (FongiLait grant). The authors would like to thank the managing committee of  
15 Bretagne Biotechnologies Alimentaires for critical reading of the manuscript and G. Le  
16 Dréan, V. Vasseur and M.A. Lebras-Quéré for stimulating discussions.

17

## 1 **References**

2

3 Anderson, I.C., Cairney, J.W.G., 2004. Diversity and ecology of soil fungal communities:  
4 increased understanding through the application of molecular techniques. *Environ.*  
5 *Microbiol.* 6, 769-779.

6 Andrighetto, C., De Dea, P., Lombardi, A., Neviani, E., Rossetti, L., Giraffa, G., 1998.  
7 Molecular identification and cluster analysis of homofermentative thermophilic  
8 lactobacilli isolated from dairy products. *Res. Microbiol.* 149, 631-643.

9 Andrighetto, C., Marcazzan, G., Lombardi, A., 2004. Use of RAPD-PCR and TTGE for the  
10 evaluation of biodiversity of whey cultures for Grana Padano cheese. *Lett. Appl.*  
11 *Microbiol.* 38, 400-405.

12 Avis, P.G., Dickie, I.A., Mueller, G.M., 2006. A 'dirty' business: testing the limitations of  
13 terminal restriction fragment length polymorphism (TRFLP) analysis of soil fungi. *Mol.*  
14 *Ecol.* 15, 873-882.

15 Bae, J.W., Rhee, S.K., Park, J.R., Chung, W.H., Nam, Y.D., Lee, I., Kim, H., Park, Y.H.,  
16 2005. Development and evaluation of genome-probing microarrays for monitoring lactic  
17 acid bacteria. *Appl. Environ. Microbiol.* 71, 8825-8835.

18 Baker, G.C., Smith, J.J., Cowan, D.A., 2003. Review and re-analysis of domain-specific 16S  
19 primers. *J. Microbiol. Methods* 55, 541-555.

20 Barlaan, E.A., Sugimori, M., Furukawa, S., Takeuchi, K., 2005. Profiling and monitoring of  
21 microbial populations by denaturing high-performance liquid chromatography. *J.*  
22 *Microbiol. Methods* 61, 399-412.

23 Berbee, M.L., Pirseyedi, M., 1999. Cochliobolus phylogenetics and the origin of known,  
24 highly virulent pathogens, inferred from ITS and glyceraldehyde-3-phosphate  
25 dehydrogenase gene sequences. *Mycologia* 91, 964-977.

1 Beresford, T.P., Fitzsimmons, N.A., Brennan, N.L., Cogan, T.M., 2001. Recent advances in  
2 cheese microbiology. *Int. Dairy J.* 11, 259-274.

3 Berthier, F., Beuvier, E., Dasen, A., Grappin, R., 2001. Origin and diversity of mesophilic  
4 lactobacilli in Comté cheese, as revealed by PCR with repetitive and species-specific  
5 primers. *Int. Dairy J.* 11, 293-305.

6 Blackwell, M., Hibbett, D.S., Taylor, W., Spatafora, J.W., 2006. Research coordination  
7 networks: a phylogeny for kingdom fungi (Deep Hypha). *Mycologia* 98, 829-837.

8 Bodrossy, L., Sessitch, A., 2004. Oligonucleotide microarrays in microbial diagnostics. *Curr.*  
9 *Opin. Microbiol.* 7, 245-254.

10 Bonaïti, C., Parayre, S., Irlinger, F., 2006. Novel extraction strategy of ribosomal RNA and  
11 genomic DNA from cheese for PCR-based investigations. *Int. J. Food Microbiol.* 107,  
12 171-179.

13 Brodie, E., Edwards, S., Clipson, N., 2003. Soil fungal community structure in a temperate  
14 upland grassland soil. *FEMS Microbiol. Ecol.* 45, 105-114.

15 Callon, C., Delbès, C., Duthoit, F., Montel, M., 2006. Application of SSCP-PCR  
16 fingerprinting to profile the yeast community in raw milk Salers cheeses. *System. Appl.*  
17 *Microbiol.* 29, 172-180.

18 Chakravorty, S., Helb, D., Burday, M., Connell, N., Aland, D., 2007. A detailed analysis of  
19 16S ribosomal RNA gene segments for the diagnosis of pathogenic bacteria. *J.*  
20 *Microbiol. Methods* 69, 330-339.

21 Cihlar, R.L., Sypherd, P.S., 1980. The organization of the ribosomal RNA genes in the  
22 fungus *Mucor racemosus*. *Nucleic Acid Research.* 25, 793-804.

23 Cocolin, L., Diez, A., Urso, R., Rantsiou, K., Comi, G., Bergmaier, I., Beimfohr, C., 2007.  
24 Optimization of conditions for profiling bacterial populations in food by culture-  
25 independent methods. *Int. J. Food Microbiol.* 120, 100-109.



- 1 Cocolin, L., Innocente, N., Biasutti, M., Comi, G., 2004. The late blowing in cheese: a new  
2 molecular approach based on PCR and DGGE to study the microbial ecology of the  
3 alteration process. *Int. J. Food Microbiol.* 90, 83-91.
- 4 Coppola, S., Blaiotta, G., Ercolini, D., Moschetti, G., 2001. Molecular evaluation of microbial  
5 diversity occurring in different types of Mozzarella cheese. *J. Appl. Microbiol.* 90, 414-  
6 420.
- 7 Coppola, S., Blaiotta, G., Ercolini, D., 2007. Dairy products. In: Cocolin, L., Ercolini, D.  
8 (Eds.), *Molecular Techniques in the Microbial Ecology of Fermented Foods*. Springer,  
9 New York, USA, pp. 31-90.
- 10 Coppola, S., Fusco, V., Andolfi, R., Aponte, M., Blaiotta, G., Ercolini, D., Moschetti, G.,  
11 2006. Evaluation of microbial diversity during the manufacture of Fior di Latte di  
12 Agerola, a traditional raw milk pasta-filata cheese of the Naples area. *J. Dairy Res.* 73,  
13 264-272.
- 14 Dasen, A., Berthier, F., Grappin, R., Williams, A.G., Banks, J., 2003. Genotypic and  
15 phenotypic characterization of the dynamics of the lactic acid bacterial population of  
16 adjunct-containing Cheddar cheese manufactured from raw and microfiltered pasteurized  
17 milk. *J. Appl. Microbiol.* 94, 595-607.
- 18 Delbès, C., Montel, M.C., 2005. Design and application of Staphylococcus-specific single  
19 strand conformation polymorphism-PCR analysis to monitor Staphylococcus populations  
20 diversity and dynamics during production of raw milk cheese. *Lett. Appl. Microbiol.* 41,  
21 169-174.
- 22 Delbès, C., Ali-Mandjee, L., Montel, M.C., 2007. Monitoring bacterial communities in raw  
23 milk and cheese by culture-dependent and -independent 16S rRNA gene-based analyses.  
24 *Appl. Environ. Microbiol.* 73, 1882-1891.

1 Dickie, I.A., Xu, B., Koide, R.T., 2002. Vertical niche differentiation of ectomycorrhizal  
2 hyphae in soil as shown by T-RFLP analysis. *New Phytol.* 156, 527-535.

3 Doaré-Lebrun, E., El Arbi, A., Charlet, M., Guérin, L., Pernelle, J.J., Ogier, J.C.C., Bouix,  
4 M., 2006. Analysis of fungal diversity of grapes by application of temporal temperature  
5 gradient gel electrophoresis – potentialities and limits of the method. *J. Appl. Microbiol.*  
6 101, 1340-1350.

7 Drake, M., Small, C., Spence, K.D., Swanson, B.G., 1996. Rapid detection and identification  
8 of *Lactobacillus* spp. in dairy products by using the polymerase chain reaction. *J. Food*  
9 *Prot.* 59, 1031-1036.

10 Duggan, D.J., Bittner, M., Chen, Y., Meltzer, P., Trent, J.M., 1999. Expression profiling  
11 using cDNA microarrays. *Nature Genet.* 21, 10-14.

12 Duthoit, F., Godon, J., Montel, M., 2003. Bacterial community dynamics during production  
13 of registered designation of origin Salers cheese as evaluated by 16S rRNA single-strand  
14 conformation polymorphism analysis. *Appl. Environ. Microbiol.* 69, 3840-3848.

15 Duthoit, F., Tessier, L., Montel, M.C., 2005a. Diversity, dynamics and activity of bacterial  
16 populations in ‘Registered Designation of Origin’ Salers cheese by single-strand  
17 conformation polymorphism analysis of 16S rRNA genes. *J. Appl. Microbiol.* 98, 1198-  
18 1208.

19 Duthoit, F., Callon, C., Tessier, L., Montel, M.C., 2005b. Relationships between sensorial  
20 characteristics and microbial dynamics in ‘Registered Designation of Origin’ Salers  
21 cheese. *Int. J. Food Microbiol.* 103, 259-270.

22 Einax, E., Voigt, K., 2003. Oligonucleotide primers for the universal amplification of beta-  
23 tubulin genes facilitate phylogenetic analyses in the regnum Fungi. *Org. Divers. Evol.* 3,  
24 185-194.

- 1 El Baradei, G., Delacroix-Buchet, A., Ogier, J.C., 2007. Biodiversity of bacterial ecosystems  
2 in traditional Egyptian Domiati cheese. *Appl. Environ. Microbiol.* 73, 1248-1255.
- 3 Elder, J.F. Jr, Turner, B.J., 1995. Concerted evolution of repetitive DNA sequences in  
4 Eukaryotes. *Eukaryotes. Q. Rev. Biol.* 70, 297-320.
- 5 Ercolini, D., 2004. PCR-DGGE fingerprinting: novel strategies for detection of microbes in  
6 food. *J. Microbiol. Methods* 56, 297-314.
- 7 Ercolini, D., Frisso, G., Mauriello, G., Salvatore, F., Coppola, S., 2008. Microbial diversity in  
8 natural whey cultures used for the production of Caciocavallo Silano PDO cheese. *Int. J.*  
9 *Food Microbiol.* doi: 10.1016/j.ijfoodmicro.2008.03.007.
- 10 Ercolini, D., Hill, P.J., Dodd, C.E.R., 2003. Bacterial community structure and location in  
11 Stilton cheese. *Appl. Environ. Microbiol.* 69, 3540-3548.
- 12 Ercolini, D., Mauriello, G., Blaiotta, G., Moschetti, G., Coppola, S., 2004. PCR-DGGE  
13 fingerprints of microbial succession during a manufacture of traditional water buffalo  
14 mozzarella cheese. *J. Appl. Microbiol.* 96, 263-270.
- 15 Ercolini, D., Moschetti, G., Blaiotta, G., Coppola, S., 2001. The potential of a polyphasic  
16 PCR-DGGE approach in evaluating microbial diversity of natural whey cultures for  
17 water-buffalo Mozzarella cheese production: bias of culture-dependent and culture-  
18 independent analyses. *Syst. Appl. Microbiol.* 24, 610-617.
- 19 Ercolini, D., Blaiotta, G., Moschetti, G., Coppola, S., 2002. Evaluation of PCR-DGGE  
20 analysis for molecular typing cheeses. *Ann. Microbiol.* 52, 81-87.
- 21 Feurer, C., Vallaes, T., Corrieu, G., Irlinger, F., 2004a. Does smearing inoculum reflect the  
22 bacterial composition of the smear at the end of the ripening of a French soft, red-smear  
23 cheese? *J. Dairy Sci.* 87, 3189-3197.
- 24 Feurer, C., Irlinger, F., Spinnler, H.E., Glaser, P., Vallaes, T., 2004b. Assessment of the rind  
25 microbial diversity in a farmhouse-produced vs a pasteurized industrially produced soft

1 red-smear cheese using both cultivation and rDNA-based methods. *J. Appl. Microbiol.*  
2 97, 546-556.

3 Fitzsimmons, N.A., Cogan, T.M., Condon, S., Beresford, T., 1999. Phenotypic and genotypic  
4 characterization of non-starter lactic acid bacteria in mature Cheddar cheese. *Appl.*  
5 *Environ. Microbiol.* 65, 3418-3426.

6 Florez, A., Mayo, B., 2006. Fungal diversity and succession during the manufacture and  
7 ripening of traditional, Spanish, blue-veined Cabrales cheese, as determined by PCR-  
8 DGGE. *Int. J. Food Microbiol.* 10, 165-171.

9 Fox, P.F., Guinee, T.P., Cogan, T.M., McSweeney, P.L.H., 2000. *Fundamentals of Cheese*  
10 *Science*. Fox, P., Guinee, T., Cogan, T.M. (Eds.),. Aspen Publications, Gaithersburg,  
11 MD, USA.

12 Frueh, F.W., Noyer-Weidner, M., 2005. Use of denaturing high-performance liquid  
13 chromatography for rapid detection and identification of seven *Candida* species. *J. Clin.*  
14 *Microbiol.* 43, 5912-5915.

15 Garbeva, P., van Veen, J.A., van Elsas, J.D., 2004. Microbial diversity in soil: selection of  
16 microbial populations by plant and soil type and implications for disease  
17 suppressiveness. *Annu. Rev. Phytopathol.* 1, 243-270.

18 Guschin, D.Y., Mobarry, B.K., Proudnikov, D., Stahl, D.A., Rittmann, B.E., Mirzabekov,  
19 A.D., 1997. Oligonucleotide microchips as genosensors for determinative and  
20 environmental studies in microbiology. *Appl. Environ. Microbiol.* 63, 2397-2402.

21 Gente, S., Larpin, S., Cholet, O., Gueguen, M., Vernoux, J., Demasures, N., 2007.  
22 Development of primers for detecting dominant yeasts in smear-ripened cheeses. *J. Dairy*  
23 *Res.* 74, 137-145.

1 Goldenberg, O., Herrmann, S., Marjoram, G., Noyer-Weidner, M., Hong, G., Bereswill, S.,  
2 Göbel, U.B., 2007. Molecular monitoring of the intestinal flora by denaturing high  
3 performance liquid chromatography. *J. Microbiol. Methods* 68, 94-105.

4 Hansen, M.C., Tolker-Nielsen, T., Givskov, M., Molin, S., 1998. Biased 16S rDNA PCR  
5 amplification caused by interference from DNA flanking the template region. *FEMS*  
6 *Microbiol. Ecol.* 26, 141-149.

7 Head, I.M., Saunders, J.R., Pickup, R.W., 1998. Microbial evolution, diversity, and ecology: a  
8 decade of ribosomal RNA analysis of uncultivated microorganisms. *Microbial Ecol.* 35,  
9 1-21.

10 Hibbet, D., Binder, M., Bischoff, J., Blackwell, M., Cannon, P., Eriksson, O., Huhndorf, S.,  
11 James, T., Kirk, P., Lucking, R., Thorsten, H., Lumbsch, T., Lutzoni, F., Matheny, P.,  
12 McLaughlin, D., Powell, M., Redhead, S., Schoch, C., Spatafora, J., Stalpers, J.,  
13 Vilgalys, R., Aime, M., Aptroot, A., Bauer, R., Begerow, D., Benny, G., Castlebury, L.,  
14 Crous, P., Dai, Y., Gams, W., Geiser, D., Griffith, G., Gueidan, C., Hawksworth, D.,  
15 Hestmark, G., Hosaka, K., Humber, R., Hyde, K., Ironside, J., Koljalg, U., Kurtzman, C.,  
16 Larsson, K., Lichtwardt, R., Longcore, J., Miadlikowska, J., Miller, A., Moncalvo, J.M.,  
17 Mozley-Standridge, S., Oberwinkler, F., Parmasto, E., Reeb, V., Rogers, J., Roux, C.,  
18 Ryvarden, L., Sampaio, J., Schussler, A., Sugiyama, J., Thorn, R., Tibell, L., Untereiner,  
19 W., Wlaker, C., Wang, Z., Weir, A., Weiss, M., White, M., Winka, K., Yao, Y., Zhang,  
20 N., 2007. A higher-level phylogenetic classification of the Fungi. *Mycol. Res.* 111, 509-  
21 547.

22 Hugenholtz, P., Goebel, B.M., Pace, N.R., 1998. Impact of culture-independent studies on the  
23 emerging phylogenetic view of bacterial diversity. *J. Bacteriol.* 180, 4765-4774.

24 James, T.Y., Kauff, F., Schoch, C.L., Matheny, P.B., Hofstetter, V., Cox, C.J., Celio, G.,  
25 Gueidan, C., Fraker, E., Miadlikowska, J., Lumbsch, H.T., Rauhut, A., Reeb, V., Arnold,

1 A.E., Amtoft, A., Stajich, J.E., Hosaka, K., Sung, G., Johnson, D., O'Rourke, B.,  
2 Crockett, M., Binder, M., Curtis, J.M., Slot, J.C., Wang, Z., Wilson, A.W., Schussler, A.,  
3 Longcore, J.E., O'Donnel, K., Mozley-Standridge, S., Porter, D., Letcher, P.M., Powell,  
4 M.J., Taylor, J.W., White, M.M., Griffith, G.W., Davies, D.R., Humber, R.A., Morton,  
5 J.B., Sugiyama, J., Rossman, A.Y., Rogers, J.D., Pfister, D.H., Hewitt, D., Hansen, K.,  
6 Hambleton, S., Shoemaker, R.A., Kohlmeyer, J., Volkmann-Kohlmeyer, B., Spotts, R.A.,  
7 Serdani, M., Crous, P.W., Hugues, K.W., Matsuura, K., Langer, E., Langer, G.,  
8 Untereiner, W.A., Lucking, R., Budel, B., Geiser, D.M., Aptroot, A., Diederich, P.,  
9 Schmitt, I., Schultz, M., Yahr, R., Hibbett, D.S., Lutzoni, F., McLaughlin, D.J.,  
10 Spatafora, J., Vilgalys, R., 2006. Reconstructing the early evolution of Fungi using a six-  
11 gene phylogeny. *Nature* 443, 818-822.

12 Joseph, N., Krauskopf, E., Vera, M., Michot, B., 1999. Ribosomal internal transcribed spacer  
13 2 (ITS2) exhibits a common core of secondary structure in vertebrate and yeast. *Nucleic  
14 Acid Research* 27, 4533-4540.

15 Kanagawa T., 2003. Bias and artifacts in multitemplate polymerase chain reaction (PCR). *J.  
16 Biosci. Bioengin.* 96, 317-323.

17 Kellogg, E.A., Appels, R., 1995. Intraspecific and interspecific variation in 5S RNA genes are  
18 decoupled in diploid wheat relatives. *Genetics* 140, 325-343.

19 Klappenbach, J.A., Dunbar, J.M., Schmidt, T.M., 2000. rRNA operon copy number reflects  
20 ecological strategies of bacteria. *Appl. Environ. Microbiol.* 66, 1328-1333.

21 Lafarge, V., Ogier, J.C., Girard, V., Maladen, V., Leveau, J., Delacroix-Buchet, A., 2004. Le  
22 potentiel de la TTGE pour l'étude bactérienne de quelques laits crus. *Le Lait* 84, 169-  
23 178.

1 Le Bourhis, A., Doré, J., Carlier, J., Chamba, J., Popoff, M., Tholozan, J., 2007. Contribution  
2 of C. beijerinckii and C. sporogenes in association to C. tyrobutyricum to the butyric  
3 fermentation in Emmental type cheese. *Int. J. Food Microbiol.* 113, 154-163.

4 Le Bourhis, A., Saunier, K., Doré, J., Carlier, J., Chamba, J., Popoff, M., Tholozan, J., 2005.  
5 Development and validation of PCR primers to assess the diversity of Clostridium spp. in  
6 cheese by temporal temperature gradient gel electrophoresis. *Appl. Environ. Microbiol.*  
7 71, 29-38.

8 Liao, D., 1999. Concerted evolution: molecular mechanism and biological implications. *Am.*  
9 *J. Hum. Genet.* 64, 24-30.

10 Liao, D., 2000. Gene conversion drives within genic sequences: concerted evolution of  
11 ribosomal RNA genes in bacteria and archaea. *J. Mol. Evol.* 51, 305-317.

12 Liu, J., Schardl, C., 1994. A conserved sequence in internal transcribed spacer 1 of plant  
13 nuclear genes. *Plant Mol. Biol.* 26, 775-778.

14 Liu, W.T., Marsh, T.L., Forney L.J., 1998. Determination of the microbial diversity of  
15 anaerobic-aerobic activated sludge by a novel molecular biological technique. *Water Sci.*  
16 *Technol.* 37, 417-422.

17 Liu, W.T., Marsh, T.L., Cheng, H., Forney, L.J., 1997. Characterization of microbial  
18 diversity by determining terminal restriction fragment length polymorphisms of genes  
19 encoding 16S rRNA. *Appl. Environ. Microbiol.* 63, 4516-4522.

20 Lutzoni, F., Kauff, F., Cox, C.J., McLaughlin, D., Celio, G., Dentinger, B., Padamsee, M.,  
21 Hibbet, M., James, T.Y., Baloch, E., Grube, M., Reeb, V., Hofstetter, V., Schoch, C.,  
22 Arnold, A.E., Miadlikowska, J., Spatafora, J., Johnson, D., Hambleton, S., Crockett, M.,  
23 Shoemaker, R., Sung, J., Lucking, R., Lumbsch, T., O'Donnell, K., Binder, M.,  
24 Diederich, P., Ertz, D., Gueidan, C., Hansen, K., Harris, R.C., Hosaka, K., Lim, Y.,  
25 Matheny, B., Nishida, H., Pfister, D., Rogers, J., Rossman, A., Schmitt, I., Sipman, H.,

1 Stone, J., Sugiyama, J., Yahr, R., Vilgalys, R., 2004. Assembling the fungal tree of life:  
2 progress, classification and evolution of subcellular traits. *Am. J. Bot.* 91, 1446-1480.

3 Mannu, L., Comunian, R., Maria, M.F., 2000. Mesophilic lactobacilli in Fiore Sardo cheese:  
4 PCR identification and evolution during ripening. *Int. Dairy J.* 10, 383-389.

5 Mauriello, G., Moio, L., Genovese, A., Ercolini, D., 2003. Relationships between flavoring  
6 capabilities, bacterial composition, and geographical origin of natural whey cultures used  
7 for traditional water-buffalo Mozzarella cheese manufacture. *J. Dairy Sci.* 86, 486-497.

8 Muyzer, G., Teske, A., Wirsén, C.O., Jannasch, H.W., 1995. Phylogenetic relationships of  
9 *Thiomicrospora* species and their identification in deep-sea hydrothermal vent samples  
10 by denaturing gradient gel electrophoresis of 16S rDNA fragments. *Arch. Microbiol.*  
11 164, 165-172.

12 Myers, R.M., Fischer, S.G., Maniatis, T., Lerman, L.S., 1985a. Modification of the melting  
13 properties of duplex DNA by attachment of a GC-rich DNA sequence as determined by  
14 denaturing gradient gel electrophoresis. *Nucleic Acid Research* 13, 3111-3129.

15 Myers, R.M., Fischer, S.G., Lerman, L.S., Maniatis, T., 1985b. Nearly all single base  
16 substitutions in DNA fragments joined to a GC-clamp can be detected by denaturing  
17 gradient gel electrophoresis. *Nucleic Acid Research* 13, 3131-3145.

18 Myers, R.M., Maniatis, T., Lerman, L.S., 1987. Detection and localization of single base  
19 changes by denaturing gradient gel electrophoresis. *Method. Enzymol.* 155, 501-527.

20 Oefner, P.J., Underhill, P.A., 1995. Comparative DNA sequencing by denaturing high-  
21 performance liquid chromatography (DHPLC). *Am. J. Hum. Genet.* 57, A266.

22 Ogier, J.C., Lafarge, V., Girard, V., Rault, A., Maladen, V., Gruss, A., Leveau, J., Delacroix-  
23 Buchet, A., 2004. Molecular fingerprinting of dairy microbial ecosystems by use of  
24 temporal temperature and denaturing gradient gel electrophoresis. *Appl. Environ.*  
25 *Microbiol.* 70, 5628-5643.



1 Ogier, J.C., Son, O., Gruss, A., Tailliez, P., Delacroix-Buchet, A., 2002. Identification of the  
2 bacterial microflora in dairy products by temporal temperature gradient gel-  
3 electrophoresis. *Appl. Environ. Microbiol.* 68, 3691-3701.

4 Orita, M., Iwahana, H., Kanazawa, H., Hayashi, K., Sekiya, T., 1989. Detection of  
5 polymorphisms of human DNA by gel electrophoresis as single-strand conformation  
6 polymorphisms. *Proc. Natl. Acad. Sci. USA* 86, 2766-2770.

7 Parayre, S., Falentin, H., Madec, M., Sivieri, K., Le Dizes, A., Sohier, D., Lortal, S., 2007.  
8 Easy DNA extraction and optimisation of PCR-Temporal temperature gel electrophoresis  
9 to identify the predominant high and low GC-content bacteria from dairy products. *J.*  
10 *Microbiol. Methods* 69, 431-441.

11 Peplies, J., Glockner, F.O., Amann, R., 2003. Optimization strategies for DNA microarrays-  
12 based detection of bacteria with 16S rRNA-targeting oligonucleotide probes. *Appl.*  
13 *Environ. Microbiol.* 69, 1397-1407.

14 Rademaker, J.L.W., Peinhopf, M., Rijnen, L., Bockelmann, W., Noordman, W.H., 2005. The  
15 surface microflora dynamics of bacterial smear-ripened Tilsit cheese determined by T-  
16 RFLP DNA population fingerprint analysis. *Int. Dairy J.* 15, 785-794.

17 Randazzo, C.L., Torriani, S., Akkermans, A.D.L., de Vos, W.M., Vaughan, E.E., 2002.  
18 Diversity, dynamics, and activity of bacterial communities during production of an  
19 artisanal Sicilian cheese as evaluated by 16S rRNA analysis. *Appl. Environ. Microbiol.*  
20 68, 1882-1892.

21 Randazzo, C.L., Vaughan, E.E., Caggia, C., 2006. Artisanal and experimental Pecorino  
22 Siciliano cheese: microbial dynamics during manufacture assessed by culturing and  
23 PCR-DGGE analyses. *Int. J. Food Microbiol.* 109, 1-8.

24 Ravnik-Glavac, M., Glavac, D., Dean, M., 1994. Sensitivity of single-strand conformation  
25 polymorphism and heteroduplex method for mutation detection on the cystic fibrosis

1 gene. Hum. Mol. Genet. 3, 801-807.

2 Reysenbach, A.L., Giver, L.J., Wickham, G.S., Pace, N.R., 1992. Differential amplification of  
3 rRNA genes by polymerase chain reaction. Appl. Environ. Microbiol. 58, 3417-3418.

4 Samson, R.A., Seifert, K.A., Kuijpers, A., Houbraken, J., Frisvad, C., 2004. Phylogenetic  
5 analysis of Penicillium subgenus Penicillium using partial  $\beta$ -tubulin sequences. Stud.  
6 Mycol. 49, 175-200.

7 Sandine, W.E., Elliker, P.R., 1970. Microbially induced flavors and fermented foods: flavor  
8 in fermented dairy products. J. Agric. Food Chem. 18, 557-566.

9 Saubusse, M., Millet, L., Delbès, C., Callon, C., Montel, M.C., 2007. Application of single  
10 strand conformation polymorphism-PCR method for distinguishing cheese bacterial  
11 communities that inhibit Listeria monocytogenes. Int. J. Food Microbiol. 116, 126-135.

12 Seifert, K.A., Samson, R.A., Dewaard, J.R., Houbraken, J., Levesque, C.A., Moncalvo, J.M.,  
13 Louis-Seize, G., Hebert, P.D.N., 2007. Prospects for fungus identification using COI  
14 DNA barcodes, with Penicillium as a test case. Proc. Natl. Acad. Sci. USA 104, 3901-  
15 3906.

16 Sheffield, V.C., Beck, J.S., Stone, E.M., Myers, R.M., 1989. Attachment of a 40 bp G+C rich  
17 sequence (GC-clamp) to genomic DNA fragments by polymerase chain reaction results  
18 in improved detection of single-base changes. Proc. Natl. Acad. Sci. USA 86, 232-236.

19 Skouboe, P., Taylor, J.W., Frisvad, J.C., Lauritsen, D., Larsen, L., Alboek, C., Boysen, M.,  
20 Rossen, L., 2000. Molecular methods for differentiation of closely related Penicillium  
21 species. In: Samson, R.A., Pitt, J.I. (Eds.), Integration of Modern Taxonomic Methods  
22 for Penicillium and Aspergillus Classification. Amsterdam: Harwood Academic  
23 Publishers, pp. 179-188.

1 Small, J., Call, D.R., Brockman, F.J., Staub, T.M., Chandler, D.P., 2001. Direct detection of  
2 16S rRNA in soil extracts by using oligonucleotide microarrays. *Appl. Environ.*  
3 *Microbiol.* 67, 4708-4716.

4 Stackebrandt, E., Frederiksen, W., Garrity, G., Grimont, P.A.D., Kämpfer, P., Maiden,  
5 M.C.J., Nesme, X., Rosselo-Mora, R., Swings, J., Trüper, H.G., Vauterin, L., Ward,  
6 A.C., Whitman, W.B., 2002. Report of the ad hoc committee for the re-evaluation of the  
7 species definition in bacteriology. *Int. J. Syst. Evol. Biol.* 52, 1043-1047.

8 Suzuki, M.T., Giovannoni, S.J., 1996. Bias caused by template annealing in the amplification  
9 of mixtures of 16S rRNA genes by PCR. *Appl. Environ. Microbiol.* 62, 625-630.

10 Tao, F., Li, C., Smejkal, G., Lazarev, A., Lawrence, N., Schumacher, R.T., 2007. Pressure  
11 cycling technology (PCT) applications in extraction of biomolecules from challenging  
12 biological samples. *High Pressure Bioscience and biotechnology*. In: *Proceedings of the*  
13 *4<sup>th</sup> international conference on high pressure and biotechnology*, Vol 1, pp. 166-173

14 Throbäck, I.N., Enwall, K., Jarvis, A., Hallin, S., 2004. Reassessing PCR primers targeting  
15 *nirS*, *nirK* and *nosZ* genes for community surveys of denitrifying bacteria with DGGE.  
16 *FEMS Microbiol. Ecol.* 49, 401-417.

17 Valinsky, L., Della Vedova, G., Scupham, A.J., Alvey, S., Figueroa, A., Yin, B., Hartin, R.J.,  
18 Chrobak, M., Crowley, D.E., Jiang, T., Borneman, J., 2002. Analysis of bacterial  
19 community composition by oligonucleotide fingerprinting of rRNA genes. *Appl.*  
20 *Environ. Microbiol.* 68, 3243-3250.

21 Van de Peer, Y., Chapelle, S., De Wachter, R., 1996. A quantitative map of nucleotide  
22 substitution rates in bacterial rRNA. *Nucleic Acid Research* 24, 3381-3391.

23 Wagner, M., Smidt, H., Loy, A., Zhou, J., 2007. Unravelling microbial communities with  
24 DNA-microarrays: challenges and future directions. *Microbial Ecol.* 53, 498-506.

- 1 Weissensteiner, T., Lanchbury, J.S., 1996. Strategy for controlling preferential amplification  
2 and avoiding false negatives in PCR typing. *Biotechniques* 21, 1102-1108.
- 3 Wintzingerode, v F., Göbel, U.B., Stackebrandt, E., 1997. Determination of microbial  
4 diversity in environmental samples: pitfalls of PCR-based analysis. *FEMS Microbiol.*  
5 *Rev.* 21, 213-229.
- 6 Walsh, P.S., Erlich, H.A., Higuchi, R., 1992. Preferential PCR amplification of alleles:  
7 mechanisms and solutions. *PCR Methods Appl.* 1, 241-250.
- 8 Ward, D.M., Bateson, M.M., Weller, R., Ruff-Roberts, A.L., 1992. Ribosomal RNA analysis  
9 of microorganisms as they occur in nature. *Adv. Microbial Ecol.* 12, 219–286.
- 10 Ward, D.M., Weller, R., Bateson, M.M., 1990. 16S rRNA sequences reveal uncultured  
11 inhabitants of a well-studied thermal community. *FEMS Microbiol. Rev.* 75, 105–116.
- 12 Wilson, I.G., 1997. Inhibition and facilitation of nucleic acid amplification. *Appl. Environ.*  
13 *Microbiol.* 63, 3741-3751.
- 14 Woese, C.R., Fox, G.E., 1977. Phylogenetic structure of the prokaryotic domain: the primary  
15 kingdoms. *Proc Natl. Acad. Sci. USA* 74, 5088-5090.
- 16 Xiao, W., Oefner, P.J., 2001. Denaturing high-performance liquid chromatography: A review.  
17 *Hum. Mutat.* 17, 439-474.
- 18 Yoshino, K., Nishigaki, K., Husimi, Y., 1991. Temperature sweep gel electrophoresis: a  
19 simple method to detect point mutations. *Nucleic Acid Research* 19, 3153.
- 20 Zhou, J., 2003. Microarrays for bacterial detection and microbial community analysis. *Curr.*  
21 *Opin. Microbiol.* 6, 288-294.

22