



HAL
open science

Isolation and characterization of nine microsatellite loci of *Terapon jarbua* (Forsskål, 1775) from Socotra Island (Gulf of Aden) using multiplex PCR

Edouard Lavergne, Isabelle Calves, Uwe Zajonz, Jean Laroche

► To cite this version:

Edouard Lavergne, Isabelle Calves, Uwe Zajonz, Jean Laroche. Isolation and characterization of nine microsatellite loci of *Terapon jarbua* (Forsskål, 1775) from Socotra Island (Gulf of Aden) using multiplex PCR. 2011, pp.418-421. hal-00573769

HAL Id: hal-00573769

<https://hal.univ-brest.fr/hal-00573769v1>

Submitted on 21 Jan 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title:**

2 Isolation and characterization of nine microsatellite loci of *Terapon jarbua* (Forsskål, 1775)
3 from Socotra Island (Gulf of Aden) using multiplex PCR.

4

5 **Author's names with initials:**6 E. LAVERGNE ^{1,2,3} (EL)7 I. CALVES ³ (IC)8 U. ZAJONZ ^{2,1} (UZ)9 J. LAROCHE ³ (JL)

10

11 **Full postal addresses:**

12 ¹ Senckenberg Forschungsinstitut und Naturmuseum, Sektion Ichthyologie –
13 Senckenberganlage 25, 60325 Frankfurt am Main, Deutschland

14 ² Biodiversität und Klima Forschungszentrum (BiK-F) – Senckenberganlage 25, 60325
15 Frankfurt am Main, Deutschland

16 ³ UMR 6539 CNRS/UBO/IRD - LEMAR (Laboratoire des Sciences de l'Environnement
17 Marin), IUEM (Institut Universitaire Européen de la Mer), Université de Bretagne
18 Occidentale – Technopôle Brest Iroise, Rue Dumont d'Urville 29280 Plouzané, France

19

20 **Keywords:** *Terapon jarbua*, microsatellites, population genetics, Socotra Island

21

22

23

24

25 **Corresponding author:**

26 Edouard Lavergne

27 Senckenberg Forschungsinstitut und Naturmuseum, Sektion Ichthyologie –

28 Senckenberganlage 25, 60325 Frankfurt am Main, Deutschland

29 Fax: +49 69 75 42 12 53

30 E-mail: edouard.lavergne@senckenberg.de

31

32 **Abstract**

33 Ten polymorphic microsatellite loci were identified and characterized from 22 individuals of
34 *Terapon jarbua* (Forsskål, 1775) from Socotra Island (Gulf of Aden, Yemen). Microsatellite
35 polymorphism was tested, revealing 4 to 19 alleles per locus. The observed heterozygosity
36 values ranged from 0.318 to 0.909. Nine loci out of ten conformed to Hardy-Weinberg
37 proportions. They did not show evidence for null alleles and gametic disequilibrium. These
38 loci will be used in an ongoing study of the population structure of this species; associated
39 with a study assessing habitat connectivity based on otolith microchemistry of *T. jarbua*.
40 Results are expected to inform estuarine conservation efforts on Socotra Island and in the
41 Gulf of Aden region.

42

43 *Terapon jarbua* (Terapontidae, Perciformes) inhabits marine and brackish waters of the Indo-
44 Pacific, from the Red Sea and East coast of Africa to Samoa. Its juveniles have been observed
45 to thrive even in coastal freshwater courses. The type locality of the species is Jeddah, Saudi
46 Arabia, and the type series is deposited in the Zoological Museum of Copenhagen
47 (Klausewitz & Nielsen 1965, Nielsen 1974). Although advances have been made in the
48 taxonomy and biology of the Terapontidae (Vari 1978, Whitfield & Blaber 1978, Miu 1990),

49 no in-depth studies of the reproductive ecology, life history strategy and population structure
50 of *T. jarbua* have been conducted as of yet. The microsatellite markers presented herein are
51 therefore the first developed for this species, and will be especially supportive to further
52 genetic studies of the species.

53 DNA libraries for *Terapon jarbua* enriched for microsatellite sequences containing AAC,
54 ATG, CATC and TAGA repeat motifs were constructed by Genetic Identification Services
55 following the method described by Jones et al. (2002). Resulting recombinant clones were
56 selected at random and sequenced on an Applied Biosystems™ 377 DNA Sequencer, using
57 Amersham's DYEnamic™ ET Terminator Cycle Sequencing Kit (Amersham Biosciences P/N
58 US81050). Initial polymerase chain reaction (PCR) primers were designed for flanking
59 regions of microsatellite containing sequences using DESIGNER PCR v1.03 (Research Genetics
60 Inc.).

61 Genomic DNA was extracted from muscle tissue, preserved in 95% ethanol of 22 individuals
62 collected in March 2007 by seine net at Matief Estuary (12° 26' 48.5'' N and
63 54° 18' 17.6'' E) on Socotra Island (Fig. 1). Extractions were performed using AcroPrep™ 96
64 well Filter Plates (1mL) with 1µm Glass Fiber media (PALL® 5051), following the extraction
65 protocol for DNA barcoding by Ivanova et al. (2006). PCRs conducted on a GeneAmp PCR
66 system 9700 (Applied Biosystems™) were optimized for each primer individually on four
67 randomly selected samples. After optimisation, PCR reactions were multiplexed in a total
68 reaction volume of 10µL, using 5µL of Master-mix, 1µL of Solution Q both from the Qiagen
69 Multiplex PCR kit, 1µL of multiplexed primer-mix (Table 1) containing specific primers and
70 the labelled universal primers 6Fam-TAGTCGACGACCGTTA, Yakima Yellow YY-
71 TCGGATAGCTAGTCGT, and Dargonfly Orange DO-CTGGCCGTCGTTTTAC (Chang et
72 al. 2004) in order to avoid the expenses of using specific fluorescent primers, 1µL of template

73 DNA (30-50 ng. μL^{-1}) and 2 μL of water. Touchdown PCR conditions consisted of an initial
74 denaturing step at 95°C (15') followed by 11 cycles at 94°C (30''), 63-53°C (1'30'') and
75 72°C (1'), followed by 25 cycles at 94°C (30''), 53°C (1'30'') and 72°C (1'). A final
76 elongation step at 60°C (30 min) ended the PCR. 3 μL of PCR products were added to 12 μL
77 of formamide and 0.2 μL of a 50-500bp size standard (GeneScanTM-500 LIZTM) to visualise
78 microsatellite alleles using an ABI 3130 Genetic Analyzer (Applied BiosystemsTM). Alleles
79 were then scored using GeneMapper[®] Software v4.0 (Applied BiosystemsTM). The number of
80 alleles and the observed and expected heterozygosity values were calculated using
81 GENETIX v4 (Belkhir 2004); deviation from the Hardy–Weinberg proportions (Fisher's exact
82 test) and gametic disequilibrium (Fisher's exact test) among loci were tested using GENEPOP
83 v4 (Raymond & Rousset 1995, Rousset 2008). Both tests were corrected for multiple
84 simultaneous tests by calculating the q-value of each test which measures the minimum *false*
85 *discovery rate* (FDR) that is incurred when calling that test significant. The bootstrap method
86 was chosen as recommended by the authors for a limited number of p-values (Storey 2002).
87 The q-values were calculated using the R package QVALUE (www.r-project.org, Ihaka &
88 Gentleman 1996, Storey 2002, Storey & Tibshirani 2003, Storey 2003, Storey et al. 2004).
89 This correction was preferred over the commonly used sequential Bonferroni correction (Rice
90 1989) following Moran (2003). Null allele frequencies were calculated based on Brookfield
91 (1996) using the program MICRO-CHECKER (Van Oosterhout et al. 2004).
92 Ten out of 12 loci were reliably amplified and found to be polymorphic for *T. jarbua*
93 (Table 2). The number of alleles per locus ranged from 4 to 19, with observed and expected
94 heterozygosity values varying respectively from 0.318 to 0.909, and from 0.328 to 0.941.
95 Exact tests after correction indicated that one locus (B107) deviated significantly from Hardy-
96 Weinberg proportions (q-value < 0.01). Exact test for gametic disequilibrium yielded five

97 weakly significant p-values ($p\text{-value} < 0.05$) out of 45 pair wise comparisons; none of which
98 were ultimately found to be significant following the FDR correction ($q\text{-value} > 0.05$). No
99 locus showed evidence for a null allele. Therefore, nine of the ten markers presented in this
100 study can be applied in studying the genetic structure of *T. jarbua* populations. Such studies
101 are expected to be instrumental in future estuary conservation and management efforts (1) in
102 Yemen including the Socotra Archipelago, representing a UNESCO World Heritage (2008),
103 which is a treasure of marine biodiversity of regional and global importance (Zajonz & Krupp
104 2006), and (2) in the wider Indian Ocean.

105

106 **Acknowledgments**

107 The present study was mainly funded by the Total Foundation, Paris. It was partially
108 conducted at the Socotra Field Research Station and the Molecular Laboratory of the
109 Biodiversity and Climate Research Centre (BiK-F) based on financial support of the research
110 funding programme "LOEWE – Landes-Offensive zur Entwicklung Wissenschaftlich-
111 ökonomischer Exzellenz" of Hesse's Ministry of Higher Education, Research, and the Arts.
112 Additional mobility funds were provided by scholarships from the GRADE - Goethe Graduate
113 Academy of the Goethe University Frankfurt a.M. and the International Doctoral College of
114 the European University of Brittany (UEB). A European Community fund (CPER-FEDER
115 project) allowed the LEMAR laboratory to acquire the high performance DNA sequencer used
116 in this study. We would like to express our gratitude to the Ministry of Water and
117 Environment and the Environment Protection Authority of Yemen for granting research and
118 sample export permits. The Socotra Conservation and Development Program (SCDP-UNDP)
119 is thanked for providing indispensable support during the field work. We wish to express our
120 gratitude to Fouad Naseeb and Motea Sheikh Aideed for their efficient assistance in the field,

121 to Dr. Le Maréchal for providing us the universal primer sequences and to Dr. Leila
122 Meistertzheim for her useful comments on the manuscript. Dr. Friedhelm Krupp is cordially
123 acknowledged for his support to the PhD project of the first author.

124

125 **References**

126 Belkhir K, Borsa P, Chikhi L, Raufaste N & Bonhomme F (1996-2004) GENETIX v4.05.
127 logiciel sous Windows[™] pour la génétique des populations. Laboratoire Génome, Populations.
128 Interactions, CNRS UMR 5000, Université de Montpellier II, Montpellier (France).

129 Brookfield JFY (1996) A simple new method for estimating null allele frequency from
130 heterozygote deficiency. *Molecular Ecology*, **5**, 453-455.

131 Chang EH, Menezes M, Meyer NC, Cucci RA, Vervoort VS, Schwartz CE & Smith RJ
132 (2004) Branchio-oto-renal syndrome: The mutation spectrum in EYA1 and its phenotypic
133 consequences. *Human Mutation*, **23**, 582-589.

134 Ihaka R & Gentleman R (1996) R: A language for data analysis and graphics. *Journal of*
135 *Computational and Graphical Statistics*, **5**, 299-314.

136 Ivanova NV, de Waard J, Hebert PDN (2006) An inexpensive, automation-friendly protocol
137 for recovering high-quality DNA. *Molecular Ecology Notes*, **6**, 998-1002.

138 Jones K, Levine K & Banks J (2002) Characterization of 11 polymorphic tetranucleotide
139 microsatellites for forensic applications in California elk (*Cervus elaphus canadensis*).
140 *Molecular Ecology Notes*, **2**, 425-427.

141 Klausewitz W & Nielsen J (1965) On Forsskål's collection in the Zoological Museum of
142 Copenhagen. Spolia zoological museum Hauniensis, Copenhagen, pp.1-29.

143 Miu TS, Lee SC & Tzeng WN (1990) Reproductive Biology of *Terapon jarbua* from the
144 Estuary of Tamshui River. *Journal of the Fishery Society of Taiwan*, **17(1)**, 9-20.

- 145 Moran MD (2003) Arguments for rejecting the sequential Bonferroni in ecological studies.
146 *Oikos*, **100**(2), 403-405.
- 147 Nielsen JG (1974) Fish types in the Zoological Museum of Copenhagen. Zoological Museum,
148 University of Copenhagen, Denmark, 115 p.
- 149 Raymond M & Rousset F (1995) GENEPOP Version 1.2: population genetics software for exact
150 tests and ecumenicism. *Journal of Heredity*, **86**, 248-249.
- 151 Rice WR (1989) Analyzing tables of statistical tests. *Evolution*, **43**, 223-225.
- 152 Rousset F (2008) GENEPOP'007: a complete reimplementation of the GENEPOP software for
153 Windows and Linux. *Molecular Ecology Resources*, **8**, 103-106.
- 154 Storey JD (2002) A direct approach to false discovery rates. *Journal of the Royal Statistical*
155 *Society, Series B*, **64**, 479-498.
- 156 Storey JD & Tibshirani R (2003) Statistical significance for genome-wide experiments.
157 *Proceeding of the National Academy of Sciences*, **100**, 9440-9445.
- 158 Storey JD (2003) The positive false discovery rate: A Bayesian interpretation and the qvalue.
159 *Annals of Statistics*, **31**, 2013-2035.
- 160 Storey JD, Taylor JE & Siegmund D (2004) Strong control, conservative point estimation and
161 simultaneous conservative consistency of false discovery rates: A unified approach. *Journal*
162 *of the Royal Statistical Society, Series B*, **66**, 187-205.
- 163 Van Oosterhout C, Hutchinson WF, Wills DPM & Shipley P (2004) MICRO-CHECKER:
164 software for identifying and correcting genotyping errors in microsatellite data. *Molecular*
165 *Ecology Notes*, **4**, 535-538.
- 166 Vari RP (1978) The terapon perches (Percoidei. Teraponidae). A cladistic analysis and
167 taxonomic revision. *Bulletin of the American Museum of Natural History*, 159 (5), 175-340.

- 168 Whitfield AK & Blaber SJM (1978) Scale-eating habits of the marine teleost *Terapon jarbua*
169 (Forsskål, 1775). *Journal of Fish Biology*, **12**, 61-70.
- 170 Zajonz U & Krupp F (2006) Fish biogeography of Socotra. In: Socotra – A natural history of
171 the islands and their people. Cheung C & De Vantier L (Eds.), Van Damme K (Science Ed.),
172 Odyssey Books and Guides, Airphoto International Ltd, 408 pp.

173 **Figure 1: Map of Socotra Archipelago and sampling site location (Matief)**

Table1: Multiplex Primer-mix

Multiplex PCR 1	Multiplex PCR 2
5µL R C102 primer, 100nM	5µL R A4 primer, 100nM
5µL R C108 primer, 100nM	5µL R B107 primer, 100nM
5µL R C3 primer, 100nM	5µL R D108 primer, 100nM
5µL F C102 primer + univ. ext., 10nM	5µL F A4 primer + univ. ext., 10nM
5µL F C108 primer + univ. ext., 10nM	5µL F B107 primer + univ. ext., 10nM
5µL F C3 primer + univ. ext., 10nM	5µL F D108 primer + univ. ext., 10nM
5µL 6FAM-univ. primer, 100nM	5µL 6FAM-univ. primer, 100nM
5µL YY-univ. primer, 100nM	5µL YY-univ. primer, 100nM
5µL DO-univ. primer, 100nM	5µL DO-univ. primer, 100nM
117µL H ₂ O	117µL H ₂ O
150 µL	150µL

Multiplex PCR 3	Multiplex PCR 4
5µL R B103 primer, 100nM	5µL R B106 primer, 100nM
5µL R C103 primer, 100nM	5µL R C105 primer, 100nM
5µL R D102 primer, 100nM	5µL R D3 primer, 100nM
5µL F B103 primer + univ. ext., 10nM	5µL F B106 primer + univ. ext., 10nM
5µL F C103 primer + univ. ext., 10nM	5µL F C105 primer + univ. ext., 10nM
5µL F D102 primer + univ. ext., 10nM	5µL F D3 primer + univ. ext., 10nM
5µL 6FAM-univ. primer, 100nM	5µL 6FAM-univ. primer, 100nM
5µL YY-univ. primer, 100nM	-
5µL DO-univ. primer, 100nM	7,5µL DO-univ. primer, 100nM
117µL H ₂ O	119,5µL H ₂ O
150µL (Total volume)	150µL (Total volume)

Primer and universal extension association is shown in Table 2

Table 2: Primer sequences for 10 microsatellite loci and allele statistics in one population (N = 22) of *Terapon jarbua*

Loci	EMBL #	Repeat motif	5' Universal Extension	Primers 5' → 3'	<i>i</i>	A	R	H_E	H_O	P_{HW}	Q_{HW}
C102	FR719958	(CTAT) ₁₃ (CCAT) ₂₈ (CTAT) ₂	F: CTGGCCGTCGTTTTACGTCTCCCTCCCTCATGTCTG R: TTGCCACAGTGGACCTGTAG	1	19	171-257	0.938	0.909	0.0662	0.0513	
C108	FR719959	(ATCC) ₉ ATTT(ATCC) ₂	F: TCGGATAGCTAGTCGTCCATCCATTCATCCATCTAC R: GCTTTGGAGTATTTTGCAGTT	1	7	274-336	0.763	0.727	0.2318	0.0773	
C3	FR719960	(CATC) ₇	F: TAGTCGACGACCGTTACATAATGAGCGAGGTCAGAT R: ATCACGGAGGTTCTAAGAGTC	1	6	278-305	0.791	0.727	0.0770	0.0513	
A4	FR719961	(AAC) ₁₄	F: TCGGATAGCTAGTCGTACCTGCCTACTACAGCCTCAG R: CACTCCACTTGCCCATTTT	2	5	262-271	0.711	0.636	0.2146	0.0773	
B107	FR719962	(CAT)CAA(CAT) ₇	F: CTGGCCGTCGTTTTACCCAAGTTCCTGATGCTAAAAG R: AGACGATGATGGGATTATTTG	2	10	205-231	0.868	0.773	0.003*	0.0008*	
B103	FR719963	(CAT) ₂ CTT(CAT) ₉ CAC(CAT)	F: TCGGATAGCTAGTCGTGGGCTGTAACAGTATGCAATG R: ATGCAGCACCTTCAGAGTTTA	3	4	215-227	0.669	0.727	0.5839	0.1557	
C103	FR719964	(TCCA) ₈	F: CTGGCCGTCGTTTTACCTTTCAATAGCCAGGACTACC R: TCTTCCACACTGAGACTGCT	3	4	183-198	0.665	0.545	0.1209	0.0645	
B106	FR719965	(CAT) ₂ CA(CAT) ₄ CAC(CAT)	F: CTGGCCGTCGTTTTACAGAGGAGGACCACATAAACAC R: TTCCACCAGATGAGAGGAG	4	11	112-159	0.859	0.727	0.0540	0.0513	
C105	FR719966	(ATCC) ₅ CTCC(ATCC) ₁₂	F: CTGGCCGTCGTTTTACAGCTTTGTGAGGCTAATACCAG R: AAGTCTTCTTCAACCCTGTGAG	4	6	242-314	0.328	0.318	0.5475	0.1557	
D3	FR719967	(TCTA) ₁₀	F: TAGTCGACGACCGTTACAGTCCAGTAATGTCGTTTGT R: AGTGTTAGACAGGAGCACATG	4	19	283-349	0.941	0.909	0.2256	0.0773	

i, multiplex PCR index (Table 1); A, allele nb; R, size range; H_E , expected heterozygosity; H_O , observed heterozygosity; P_{HW} , p-value of HW exact test; Q_{HW} , q-value of HW exact test; *, Significant deviation from HWE.

