



HAL
open science

Marine culturable yeasts in deep-sea hydrothermal vents: species richness and association with fauna

Gaëtan Burgaud, Danielle Arzur, Lucile Durand, Marie-Anne Cambon-Bonavita, Georges Barbier

► **To cite this version:**

Gaëtan Burgaud, Danielle Arzur, Lucile Durand, Marie-Anne Cambon-Bonavita, Georges Barbier. Marine culturable yeasts in deep-sea hydrothermal vents: species richness and association with fauna. FEMS Microbiology Ecology, 2010, 73 (1), pp.121-133. 10.1111/j.1574-6941.2010.00881.x. hal-00609796

HAL Id: hal-00609796

<https://hal.univ-brest.fr/hal-00609796>

Submitted on 1 Sep 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Marine culturable yeasts in deep-sea hydrothermal vents:

Species richness and association with fauna

Gaëtan Burgaud^{1*}, Danielle Arzur¹, Lucile Durand², Marie-Anne Cambon-Bonavita² & Georges Barbier^{1*}

¹Laboratoire Universitaire de Biodiversité et Ecologie Microbienne (EA3882), IFR 148, Université Européenne de Bretagne, Université de Brest, ESMISAB, Technopole Brest-Iroise, 29280 Plouzané, France.
²Ifremer, Centre de Brest, DEEP/LM2E, UMR 6197, BP 70, 29280 Plouzané, France

*Corresponding authors: georges.barbier@univ-brest.fr; gaetan.burgaud@univ-brest.fr

Keywords: Yeasts - Hydrothermal vents - Fauna - 26S rRNA gene – FISH

Running title: Culturable yeasts from hydrothermal vents

26 **Abstract**

27 The diversity of culturable yeasts at deep-sea hydrothermal sites has been investigated and
28 raises emerging hints regarding interactions with endemic fauna. Samples were collected
29 during diverse oceanographic cruises at Mid-Atlantic Ridge, South Pacific Basins and East
30 Pacific Rise. A culture collection of thirty-two isolates mostly associated with animals was
31 performed. The phylogenetic analyses of 26S rRNA gene sequences revealed that the yeasts
32 belonged to *Ascomycota* and *Basidiomycota* phyla with the identification of several genera:
33 *Rhodotorula*, *Rhodospiridium*, *Candida*, *Debaryomyces*, and *Cryptococcus*. Those genera are
34 usually isolated from deep-sea environments. To our knowledge, this is the first report of
35 yeasts associated with deep-sea hydrothermal animals.

36 **Introduction**

37 Yeasts are ubiquitous microorganisms that represent a part of the microbiota in all natural
38 ecosystems such as soils, freshwaters and marine waters from ocean surface to deep-sea.
39 Marine yeasts are divided into *obligate* and *facultative* groups. “Obligate marine yeasts” are
40 yeasts that have never been isolated from anywhere else than marine environment, whereas
41 “facultative marine yeasts” are also known from terrestrial habitats (Kohlmeyer and
42 Kohlmeyer, 1979). Based on these definitions, Kohlmeyer and Kohlmeyer (1979) have
43 examined yeasts occurring in marine environments and have gathered a list of 176 species
44 isolated from diverse marine habitats. Out of those, only 25 were obligate marine yeasts
45 widely represented by the genera *Metschnikowia*, *Rhodospiridium*, *Candida* and *Torulopsis*.

46 The existence of 1.5 million fungal species as hypothesized by Hawksworth (2002) is a
47 commonly used and accepted figure. If this estimate is correct, less than 5% of the fungi have
48 been described up to now and almost exclusively from terrestrial environments. In this
49 ecosystem, fungi are known to utilize a wide spectrum of simple and more complex organic
50 compounds. The decomposition activities of fungi are clearly important in relation to the
51 redistribution of elements among organisms and environmental compartments (Gadd, 2007).
52 Bearing in mind those parameters, our hypothesis is that deep-sea and especially
53 hydrothermal vents, which remain underexplored habitats for fungi, could be ecological
54 niches hosting specific fungal communities.

55 Deep-sea hydrothermal vents are localized at sea floor spreading centers called *riffts* where
56 seawater seeps into cracked regions caused by the presence of hot basalt and magma.

57 Seawater carrying dissolved minerals is then emitted from springs. Two major types of
58 emissions have been found. Warm fluids diffuse at temperatures ranging from 6 to 23°C into
59 seawater at 2-4°C when hot vents called black smokers emit hydrothermal fluid at 270-
60 380°C (Munn, 2003). Thermal gradients in hydrothermal vents are so important that just a
61 few centimeters away, the temperature can fall to 2-4°C allowing mesophilic or psychrophilic
62 organisms as well as thermophilic and hyperthermophilic prokaryotes to grow and interact
63 with all biotic or abiotic components of these ecosystems. Dense animal communities cluster
64 around those hot springs. These communities are supported by the chemolithoautotrophic
65 activities of prokaryotes (Jorgensen and Boetius, 2007).

66 The occurrence of fungi (filamentous fungi and yeasts) at deep-sea hydrothermal vents
67 remains an underexplored topic. Over the last years, the interest for the diversity of microbial
68 eukaryotes in these ecosystems emerged using PCR amplification of SSU ribosomal RNA
69 genes and sequence analysis (Edgcomb *et al.* 2002; Lopez-Garcia *et al.*, 2003; 2007). These
70 papers revealed a scarce fungal diversity but some sequences were novel. Only two papers
71 have specifically dealt with fungal diversity at deep-sea hydrothermal vents based on culture-
72 dependent methods (Gadanho & Sampaio, 2005; Burgaud *et al.*, 2009). Culturable yeasts
73 affiliated to *Ascomycota* and *Basidiomycota* phyla were reported from hydrothermal waters.
74 Some papers assessing fungal diversity at deep-sea vents were also published. Bass *et al.*
75 (2007) reported the presence of sequences affiliated to *Debaryomyces hansenii* and novel
76 sequences closed to *Malassezia furfur* in hydrothermal sediments. Le Calvez *et al.* (2009)
77 reported that fungal diversity from deep-sea vent animals was widely constituted of sequences
78 affiliated to *Chytridiomycota* and *Basidiomycota* phyla. The latter phylum was mostly
79 represented by yeasts with, for example, the *Cryptococcus* and *Filobasidium* genera that form
80 dense clusters.

81

82 The occurrence of yeasts in other deep-sea environments has been much more studied.
83 Nagahama *et al.* (2001b) reported that culturable fungal diversity was dominated by
84 ascomycetous yeasts in surface sediments in water depths exceeding 2000 meters (*Candida*,
85 *Debaryomyces*, *Kluyveromyces*, *Saccharomyces* and *Williopsis*). Inversely, diversity was
86 dominated by basidiomycetous yeasts on the subsurface of sediments in water depths
87 exceeding 2000 meters and from deep-sea clams, tubeworms and mussels (*Rhodotorula*,
88 *Sporobolomyces*, *Cryptococcus* and *Pseudozyma*). Recent studies have clearly demonstrated
89 that *Cryptococcus* was the dominant genus sequenced from sediments collected at deep
90 methane cold seeps (Takishita *et al.*, 2006; 2007). Those observations are in agreement with

91 Bass *et al.* (2007) who suggest that yeast forms dominate fungal diversity in deep oceans.
92 Several yeasts mostly isolated from deep-sea sediments represented new species in the
93 *Ascomycota* or *Basidiomycota* phyla (Nagahama *et al.*, 1999; 2001a; 2003a; 2003b; 2006a;
94 2008).

95

96 In this study, we decided to assess the presence of yeasts at deep-sea hydrothermal vents
97 based on a culture-based approach with an emphasis on yeasts in interactions with the
98 endemic animal fauna thriving in these extreme ecosystems. A recent paper (Gadanho and
99 Sampaio, 2005) has dealt with the diversity of yeasts in deep-sea vent waters but, to our best
100 knowledge, this is the first report of the culturable yeasts isolated from deep-sea animals.
101 Those interactions with the fauna are discussed based on the cultures obtained from the
102 samples collected during different oceanographic the cruises at Mid-Atlantic Ridge, South-
103 West Pacific Lau Basin and East Pacific Rise.

104 **Materials and methods**

105 *Environmental sampling*

106 210 hydrothermal samples were collected during 6 oceanographic cruises at several dates and
107 locations (For hydrothermal vents locations, see Tivey, 2007): (i) BIOLAU in the Lau Basin,
108 South-west Pacific (12/05/1989–27/05/1989; 20°3.0'S, 176°7.8'W; -2620 m); (ii)
109 DIVANAUT2 (19/06/ 1994–01/07/1994) on the MAR at Menez Gwen (37°51'N, 31°31'W; -
110 900 m) and Lucky Strike (37°17'N, 32°16'W; -1650 m) hydrothermal sites; (iii) HERO on
111 the EPR at Elsa site (30/09/1991–04/11/1991; 12°48'N, 103°57'W; -2630 m); (iv) MARVEL
112 (29/08/1997–13/09/1997) on the MAR at Menez Gwen and Lucky Strike sites; (v) EXOMAR
113 (25/07/2005–28/08/2005) on the MAR at Rainbow (36°08'N, 34°00'W, -2300 m), TAG
114 (26°02'N, 44°54'W, -3630 m) and Lost City (30°04'N, 42°12'W, -900 m) sites; (vi)
115 MoMARDREAM-Naut (08/07/2007–19/07/2007) on the MAR at Rainbow site. Depending
116 on cruises, deep-sea sampling was performed using the Deep Submergence Vehicle “Nautile”
117 or the Remote Operated Vehicle (ROV) “Victor 6000” and the N/O “Atalante” and
118 “Pourquoi Pas?” research vessel.

119 The deep-sea samples were processed as described by Burgaud *et al* (2009) taking care to
120 avoid contamination in applying strict sterile sampling conditions.

121

122 *Enrichment conditions and isolation*

123 The samples were processed directly after the Nautile or ROV recovery. The ollected samples
124 mainly composed of deep-sea hydrothermal vent animals (*Rimicaris exoculata* and

125 *Chorocaris chacei* shrimps and *Bathymodiolus azoricus* mussels) were used to inoculate the
126 GYPS culture medium that led to the best isolation rate during a previous study (Burgaud *et*
127 *al.*, 2009). This medium contained per liter: glucose (Sigma) 1 g, yeast extract (AES) 1 g,
128 peptone (AES) 1 g, starch (Fisher) 1 g, sea salts (Sigma) 30 g. This medium was
129 supplemented per litre with agar 15 g and chloramphenicol (Sigma) 500 mg, pH was also
130 adjusted to 7.5. Cultures were done aerobically at 4°C, 15°C, 25°C (ambient temperature), 35
131 and 45°C (only during EXOMAR) at atmospheric pressure until fungal strains were
132 visualized. During the MoMARDREAM-Naut cruise, some dissections were realized on
133 board on animal samples in order to investigate the yeast location.
134 Each purified strain from our collection (Table 1) has been integrated to the ‘Souchothèque de
135 Bretagne’ culture collection
136 (<http://www.ifremer.fr/souchotheque/internet/htdocs/generique.php?pagebody=catalogue.php>)
137 and are available with an accession number associated to their GenBank number.
138

139 ***Physiological characterization and statistical analysis***

140 All experiments were done in triplicate. The yeasts were grown in liquid GYPS broth media.
141 The effect of temperature on growth was determined at 5°C, 15°C, 25°C and 35°C at 30 g.L⁻¹
142 sea salts. The effect of salinity was analyzed modifying sea salts concentrations in media from
143 0 to 60 g.L⁻¹ with steps of 15 g.L⁻¹ at optimal temperature for each strain. Optical densities
144 (OD) were measured at 600 nm with Nanocolor 100D (Macherey-Nagel, Hoerd, France) at
145 17, 22, 25 and 28 hours of growth under each condition of salinity and temperature.
146

147 ***DNA extraction and 26S rDNA sequencing***

148 DNA of each strain was extracted using FastDNA Spin Kit (MP Biomedicals, Illkirch,
149 France) specific for fungi and yeasts. Amplifications of the D1/D2 region of 26S rDNA were
150 carried out with rDNA primers ITS5 (5'-GGA AGT AAA AGT CGT AAC AAG-3'), LR6
151 (5'-CGC CAG TTC TGC TTA CC-3'), NL1 (5'-GCA TAT CAA TAA GCG GAG GAA
152 AAG-3') and NL4 (5'-GGT CCG TGT TTC AAG ACG G-3') as described by Gadanho &
153 Sampaio (2005). All PCR reactions were performed in 20 µL reaction volumes containing 19
154 µL of 1X PCR Buffer (Promega), 2 mM of MgCl₂, 0.2 mM of each dNTPs (Promega), 0.6
155 µM of primers (forward and reverse), 1.25 U of Taq DNA Polymerase (Promega) and 1µL of
156 DNA. The polymerase chain reactions were performed on PTC-200 (Biorad, France). The
157 amplification consisted in an initial denaturation step at 94°C for 2 min, followed by 30

158 iterations of 15 sec at 94°C, 30 sec at 54°C, 1 min at 72°C and a final extension step of 2 min
159 at 72°C. A negative control with sterile distilled water replacing DNA was added. Two kinds
160 of amplification were generated using ITS5-NL4 and NL1-LR6 primers. The amplified DNA
161 fragments were separated by electrophoresis in 0.8% agarose (w/v) gel (Promega) in 0.5X
162 Tris-Borate-EDTA (TBE) Buffer at 90 V for 1h and stained with ethidium bromide. A
163 molecular size marker was used for reference (Lambda DNA/EcoR1+Hind III Markers,
164 Promega). The DNA banding patterns were visualized under UV transillumination and
165 picture files were generated using Gel-Doc 2000 (Biorad, France).

166 The sequencing of the D1/D2 region of the 26S rDNA was then realized using NL1 on the
167 ITS5-NL4 fragments and NL4 on the NL1-LR6 fragments. The sequences were obtained by
168 “Big Dye Terminator” technology (Applied Biosystems). This work was done at
169 “Biogenouest” sequencing facility in the “Station Biologique de Roscoff” (www.sb-roscoff.fr).
170

171 *Phylogenetic analyses*

172 Sequences were edited and cleaned using Sequencher v 4.8 (Gene Codes). Sequences were
173 then imported to MEGA 4.0 software (Tamura *et al.*, 2007). Each sequence was analyzed in
174 order to find GenBank sequences with close BLAST-N hits (Altschul *et al.*, 1990).
175 Similarities between sequences were assessed using pairwise distance calculation with
176 MEGA 4.0. The sequences were trimmed to ensure that all sequences had the same start and
177 end-point. All the D1/D2 regions of the 26S rDNA sequences were aligned using ClustalW
178 v.1.83 (Thompson *et al.*, 1994). After visual checking and manual curation, an alignment
179 composed of 62 taxa and 579 characters was analysed for the Bayesian estimation of
180 phylogeny using MrBayes v.3.1.2 software (Ronquist and Huelsenbeck, 2003). A two-million
181 generation option has been set to run the Metropolis-coupled Monte Carlo Markov Chain
182 method (*mcmc*). After generation 2 000 000, the standard deviation of split frequencies was P
183 = 0.005997 indicating that a convergence had occurred (P-value of < 0.05). The alignment
184 was analysed using MODELTEST v.3.7 (Posada and Crandall, 1998), in order to obtain the
185 more realistic evolutionary model used for phylogenetic analyses (GTR + G model; gamma-
186 distribution shape parameter = 0.3978). Phylogeny was then evaluated using two different
187 methods: (i) Bayesian inference with MrBayes v.3.1.2 analysis using 2 000 000 generations
188 and the *mcmc* method. The tree search included two *mcmc* searches with four chains (setting
189 default temperature for heating the chains) and a sampling frequency of 100 generations. A
190 ‘burnin’ of 5000 (25% of the 2 000 000 generations/100 sample frequency) was set in order to
191 exclude the first 5000 trees generated. (ii) Maximum likelihood with 100 bootstrap iterations

192 using PHYML (Guindon *et al.*, 2005) and the parameters obtained with MODELTEST v.3.7.
193 The final phylogenetic tree topology was realized using MrBayes v.3.1.2 analysis results.
194 Nodes in the tree show Bayesian posterior probabilities and ML bootstraps respectively.

195 ***Fluorescent probe design and evaluation***

196
197 For the detection of yeasts isolated from deep-sea vent animals by FISH, we designed
198 oligonucleotide probes using the Primrose software ([http://www.bioinformatics-
199 toolkit.org/Primrose/index.html](http://www.bioinformatics-toolkit.org/Primrose/index.html)) as described by Ashelford *et al.* (2002) using a set of high-
200 quality, full-length rRNA sequences of probe target organisms. The PrimRose design tool
201 permitted to produce oligonucleotide probes for the three principal clusters of our collection
202 (Table 3). These probes exhibited no mismatches with the target organisms but exhibited
203 mismatches with the next most similar sequences in the GenBank database proving that the
204 designed probes were *in silico* highly specific. The target sites of newly designed probes were
205 checked for accessibility using the prediction maps based on the 26S rRNA of *Saccharomyces*
206 *cerevisiae* (Inacio *et al.*, 2003). Each probe was in a relative accessible area of the 26S rRNA
207 secondary structure (Fig S1). As it was not possible to test the probes with culture isolates that
208 exhibited zero or one mismatch with the probes, we used an alternative method and tested the
209 probes against all strains from our collection displaying two or more mismatches with the
210 oligonucleotides. All newly designed probes were labelled at the 5' terminus with the
211 fluorescent marker Cy3. All probes were synthesized by (Proligo, France) and stored in sterile
212 distilled water at -20°C. The newly designed probes were checked under *in situ* conditions
213 with target and non-target species. The universal probe Euk516-Fluorescein (5'-
214 ACCAGACTTGCCCTCC-3'; Amann *et al.*, 1995) and the non-Euk516-Cy3 (5'-
215 CCTCCCGTTCAGACCA-3') probes were used as positive and negative control respectively.
216 The average cell brightness was measured using different formamide concentrations from 0 to
217 80% with steps of 10%. Systematic evaluation of the signal intensities was done by recording
218 images of independent visual fields (encompassing at least 100 cells), followed by digital
219 image analysis using the daime software (Daims *et al.*, 2006). During this step, the intensities
220 of the image pixels analyzed enable determination of single cell fluorescence in relative units
221 (RU).

222 ***Fluorescence In Situ Hybridization***

223 *On environmental samples.* Interior branchiostegites of *Rimicaris exoculata* shrimps and
224 byssus of *Bathymodiolus azoricus* mussels were processed for FISH analyses. Following
225 harvest and dissections, animal subsamples were fixed with 4% paraformaldehyde solution in
226 phosphate-buffered saline (PBS) for 3 hours at 4°C in a dark room. After fixation, tissues
227 were washed three times with PBS and stored at -20°C in a storage buffer containing PBS and
228 96% ethanol (1:1).

229 *On membrane filters.* The seawater surrounding shrimps (MoMAR08, Rainbow) was sampled
230 in 5 L sterile sampling bags using a peristaltic pump. Immediately after dives, seawater
231 samples for *in situ* hybridizations were mixed with 3 % formaldehyde (final concentration) 2
232 hours at 4°C. Fixed seawater was then filtered on polycarbonate membranes 0.22 µm
233 (Nuclepore®, 47 mm diameter; Whatman, Maidstone, Kent, UK) and rinsed with a PBS 2X -
234 sterile seawater (v:v) buffer. Then filters were dehydrated using ethanol series (50 %, 80 %
235 and absolute, 3 min each). Dried filters were stored at -20°C until hybridization treatments.
236 Three membranes were treated in this study. The filtered volume was 0.8 L for membrane A,
237 1 L for membrane B and 1.5 L for membrane C. The filtered seawater on membranes A and B
238 was from the same sample.

239 The samples (environmental samples and membrane filters) were cut in squares and paste
240 with one drop of 0.2% low-gelling point agarose (35-40°C) on slides (Menzel-Glaser,
241 Germany). All slides were then dipped in 0.2% agarose and air dried. Samples were then
242 subjected to dehydration with increasing concentrations of ethanol (50, 80, and 96%, for 3
243 min each). Working solutions of probes had a concentration of 30 ng of DNA per liter. The
244 hybridization buffer, containing 0.9 M NaCl, 20 mM Tris-HCl (pH 7.2), 0.03% SDS, and 0,
245 10, 20, 30, 40, 50, 60, 70 and 80% formamide, and the fluorescent probe were gently mixed
246 in a ratio of 10:1 (vol/vol) to get a final oligonucleotide concentration of 3 ng per liter. For
247 hybridization, slides were placed in sampling tubes and incubated at 46°C in the dark for
248 exactly 3 hours. Following hybridization, the slides were washed with prewarmed washing
249 buffer (20 mM Tris/HCl, 5 mM EDTA (pH 8.0) and 900, 450, 215, 102, 46, 18, 5, 0.6 and 0
250 mM NaCl corresponding respectively to 0, 10, 20, 30, 40, 50, 60, 70 and 80% formamide
251 stringencies) for 20 min at 48°C. Slides were rinsed with double-distilled water, air dried,
252 DAPI stained (final concentration 1 µg/ml) and mounted with the antifading reagent Citifluor
253 AF 2 (Citifluor, France) before observations under fluorescent microscope.

254 **Results**

255 **Yeast isolation**

256 Yeasts were not found in all the studied sites as shown in Table 1. No yeast was isolated from
257 samples collected during HERO (on the East Pacific Rise at Elsa site), DIVANAUT2 and
258 MARVEL (Menez-Gwenn and Lucky Strike) cruises or at TAG site during the EXOMAR
259 oceanographic cruise. The hydrothermal site that yielded the highest number of isolates was
260 clearly Rainbow (29 isolates out of 32 strains). Rainbow is also the site where the highest
261 number of samples was processed (97/210). The yeast collection obtained from deep-sea
262 samples raised thirty-two isolates that could be divided in pigmented yeasts (18) and non-
263 pigmented yeasts (14). Pigmented yeasts consisted widely of red-pigmented yeasts (16),
264 black-pigmented yeast (1) and brown-pigmented yeast (1).

265 Regarding yeast isolation versus type of substrate, strains were obtained mostly from
266 hydrothermal shrimps *Rimicaris exoculata* (11), *Chorocaris chacei* (3), *Mirocaris fortunata*
267 (1) and from hydrothermal mussels *Bathymodiolus azoricus* (7). Carbonate colonization
268 modules deployed for 1 year near Rainbow vent yielded a few yeasts (4); sponges led to the
269 isolation of three yeasts. Finally, seawater, gastropods and coral samples permitted to obtain
270 one strain each (Table 1). Those results indicate that yeasts were much more associated with
271 animals rather than mineral substrates. Statistical distribution tests have been performed in
272 order to find out the distribution type of yeasts in hydrothermal sites. The variance to mean
273 ratio (s^2/m) was calculated for each site (Cancela da Fonseca, 1966). Values of s^2/m
274 significantly different of 1 corresponds with $(s^2/m) - 1 > 2(2n/(n - 1)^2)^{1/2}$ and were obtained
275 only for Rainbow site. For this hydrothermal site, an aggregate distribution was observed
276 ($s^2/m=1.32$) indicating that the culturable yeasts isolated were located in specific niches in
277 this hydrothermal site (mainly shrimps and mussels).

278

279 During the MoMARDREAM-Naut cruise, dissections of body components were processed
280 for all shrimps (branchiostegites, scaphognathites, exopodites, gills, stomach and digestive
281 tract) and mussels (interior and external faces of shells) to investigate the localization of
282 yeasts in deep-sea animals. For shrimps, a large majority of strains were grown from the inner
283 side of the branchiostegites that can be divided in 3 different compartments: (a) an antero-
284 ventral area, which was relatively clear; (b) a posterior area, which always remained light
285 beige; (c) an antero-dorsal area with an intensely rusty coloration (for schematic views, see
286 Zbinden *et al.*, 2004; Corbari *et al.*, 2008). Yeast isolates resulted from this study were all
287 cultivated from the antero-dorsal area characterized by high amounts of minerals and a dense
288 bacterial mat.

289

290 The yeasts were also isolated from *Bathymodiolus azoricus* (7) during the MoMADREAM-

291 Naut oceanographic cruise (Table 1). Most of them (6) were cultivated from external face of
292 the mussel shells and more precisely from the byssus that is a network of filaments allowing
293 attachment to rocks. This tangle gathers a lot of particles and organic matter in decomposition
294 (personal observation). Only one yeast was isolated from the interior of a mussel (Mo32).
295

296 **Physiological analysis**

297

298 Three categories of strains were identified (Table 2) based on the definition of halotolerant
299 and halophilic microorganisms (Margesin & Schinner, 2001; Kushner, 1978). Non halophiles
300 are strains with maximal growth without sea salts and a decreasing growth rate with increased
301 sea salts concentration in media. Halotolerant yeasts are strains able to grow in the absence as
302 well as in the presence of salt. Halophiles required salt for an optimal growth. Regarding
303 halophily, optimal salinities, optimal temperatures and OD measurement, 9 physiological
304 groups were defined. Most of the isolated strains were non halophiles (23 strains) and
305 halotolerant (2 strains, maximal OD at 30 g/l sea salts) growing efficiently at an optimal
306 temperature of 25°C. Four strains had poor maximal growth at 25°C including 1 non
307 halophile, 2 halotolerant (maximal OD at 30 and 60 g/l sea salts) and 1 halophile (maximal
308 OD at 30 g/l sea salts). Three strains had maximal and efficient growth at 35°C, including 1
309 non halophile, 1 halotolerant (maximal OD at 45 g/l sea salts) and 1 halophile (maximal OD
310 at 30 g/l sea salts).

311

312 **Identification**

313

314 For species identification, a sequence analysis of the D1/D2 domain of the 26S rRNA gene
315 was done (Fig 1). A total of 12 phylotypes was found among the collection of yeasts isolated
316 from deep-sea hydrothermal vents. Eleven phylotypes could be assigned to a known yeast
317 species and one represents a new yeast species.

318

319 Within *Basidiomycota*, the *Sporidiobolales* order was the dominant cluster composed of 16
320 strains. A majority of strains (Ex2, Ex3, Ex4, Ex5, Ex6, Ex7, Ex9, Ex11, Ex12, Mo32, Mo35
321 and Mo37) was identified as *Rhodotorula mucilaginosa* (100% similarity). A large majority
322 of *R. mucilaginosa* was isolated from deep-sea shrimps (14) and the others from deep-sea
323 mussels (2). As member of the *Sporidiobolales* order, isolates affiliated to *Rhodospodium*
324 *diobovatum* were also isolated (Mo24, Mo33 and Mo38) with 100% similarity. These 3

325 strains were isolated respectively from *Rimicaris exoculata* exuviae in decomposition on
326 smocker rocks, *Bathymodiolus azoricus* and a sponge. One strain isolated from *R. exoculata*
327 was identified as *Sporobolomyces roseus* based on 26S rRNA genes (Mo22) with 100%
328 similarity with the reference strain. Four strains (Mo26, Mo27, Mo28 and Mo29) were
329 affiliated to the *Filobasidiales* order and identified as *Cryptococcus uzbekistanensis* (100%
330 similarity). These four strains were all isolated from a carbonate colonization module. Finally,
331 one isolate (Mo36) from *B. azoricus* mussel was identified as *Leucosporidium scottii* in the
332 *Leucosporidiales* order.

333 The *Ascomycota* phylum gathered 9 strains belonging to the *Saccharomycetales* order. Within
334 this order, 4 strains (Mo20, Mo21, Mo40 and Bio2) isolated respectively from *R. exoculata*,
335 *Mirocaris fortunata*, a deep-sea coral and the gills of the gastropod *Ifremeria nautilei* were
336 identified as *Debaryomyces hansenii* (100% similarity). *Candida atlantica* isolates were
337 found in *R. exoculata* exuviae in decomposition (Mo25) and *B. azoricus* (Mo31). One strain
338 isolated from a deep-sea sponge (Ex15) was identified as *Pichia guilliermondii* (100%
339 similarity). Finally, among the *Saccharomycetales* order, one strain was identified as *Candida*
340 *viswanathii* (Bio1) with 100% similarity. One halophilic strain (Mo39) isolated from a deep-
341 sea coral represents a new species in the *Candida* genus and thus was identified as *Candida*
342 sp. This strain has 95% similarity with the reference sequence of *Candida atmosphaerica* (23
343 mismatches on 505 bp). Mo30 isolated from *Bathymodiolus azoricus* was identified as
344 *Phaeotheca triangularis* (mitosporic *Ascomycota*) with 100% similarity. In the *Dothideales*
345 order, one strain (Mo34) isolated from *Bathymodiolus azoricus* was identified as *Hortaea*
346 *werneckii* with 99.98% similarity (one mismatch on 560bp).

347 Sequencing of the 26S rRNA genes indicated the presence of *Ascomycota* and *Basidiomycota*
348 in our culture collection. In term of quantity, the phylum *Basidiomycota* (21) was two times
349 higher than the *Ascomycota* (11). In term of species richness, ascomycetous yeasts belonged
350 to 7 different clusters while basidiomycetous yeasts belonged to 5 clusters.

351

352 **Fluorescence *in situ* hybridizations**

353 We processed numerous assays to detect fungi on deep-sea hydrothermal vent animal samples
354 using different existing fluorescent probes from different databases. The Euk516-Cy3 probe
355 gave positive results on pure cultures but strong background fluorescence on hydrothermal
356 samples led to the renouncement of its use. The probe MY1574 targeting *Eumycota*
357 organisms (Baschien *et al.*, 2008) showed very weak fluorescence on pure cultures. Thus, we

358 decided to design our own probes (Table 3) based on our culture collection that was divided
359 in 3 main clusters: **MitoFilo** (*Cryptococcus* / Mitosporic *Filobasidiales* order), **MitoSporidio**
360 (*Rhodotorula*, *Rhodosporeidium* / Mitosporic *Sporidiobolales* order) and **Sacch**
361 (*Debaryomyces*, *Pichia* / *Saccharomycetales* order). The probes designed revealed a strong
362 specificity for the target organisms. The optimal conditions for the *in situ* hybridization
363 protocol use stringent conditions of 20% formamide (Fig S1).

364 Our aim was to check the applicability of the FISH method to the *in situ* detection of yeasts in
365 deep-sea hydrothermal fauna samples. Hydrothermal body components of endemic shrimps
366 (*Rimicaris exoculata*) and mussels (*Bathymodiolus azoricus*) were fixed for FISH
367 experiments directly after dissection. The pieces of shrimps and mussels that gave the higher
368 number of fungi isolation (interior branchiostegites of shrimps and byssus of mussels) were
369 analyzed for yeast cell fluorescence. Although shrimp and mussel samples from Rainbow site
370 led to the highest rate of isolation, no FISH signal was ever observed. The FISH detection
371 limit of 10^3 - 10^4 target cells per ml is relatively high (Daims *et al.*, 2005) and thus, the absence
372 of FISH signals does not necessarily mean that the target organisms were not present in the
373 samples.

374 To test this hypothesis, several volumes of water were concentrated on polycarbonate
375 membrane filters to yield sufficient cells for FISH experiments with these new probes.
376 Membrane filters were embedded in low gelling-point agarose to minimize cell loss. Yeast
377 cells could be visualized in a low quantity on these membrane filters (Fig 2). Such results are
378 another evidence of the yeast cells presence in hydrothermal vents but at low concentration.
379 Using FISH on membrane filters, yeast cells detected were affiliated to 3 genera:
380 *Rhodosporeidium*, *Rhodotorula* and *Cryptococcus*.

381

382 **Discussion**

383

384 *Occurrence of yeasts in deep-sea hydrothermal vents*

385 In this study, the main aim was to isolate yeast strains from deep-sea hydrothermal endemic
386 fauna knowing that yeasts can be isolated from seawater surrounding hydrothermal fauna
387 (Gadanhó and Sampaio, 2005). Yeast isolation was successful even if the retrieved species
388 richness was relatively low. Thirty-two strains were isolated mostly from *Rimicaris exoculata*

389 shrimps. The association with shrimps is probably favorable for yeasts that could benefit from
390 nutrients due to the water circulation in the gill chamber. Most of our strains were isolated
391 from the Rainbow hydrothermal site which confirms previous results (Gadanhó and Sampaio,
392 2005). The Rainbow hydrothermal field hosted in ultramafic rocks is a unique vent enriched
393 in CH₄, H₂, CO, Fe and depleted in H₂S (Charlou *et al.*, 2002). The high yeast isolation ratio
394 may indicate that yeasts thrive in hydrothermal sites depleted in H₂S. The isolation rate of
395 non-pigmented yeasts on sulfur-free media significantly higher than those on sulfur-based
396 media in a previous study (Gadanhó & Sampaio, 2005) support such hypothesis.

397

398 Several yeasts were also isolated from mussels and more precisely from the byssus
399 constituted of filaments with a high concentration of minerals and organic matter. These
400 yeasts may have a role in the decomposition of organic material entrapped in mussel byssi in
401 deep-sea vents. These results seem promising as they confirm the data obtained in previous
402 studies and suggest that yeasts may interact with deep-sea hydrothermal vent fauna.

403

404 ***Pattern of the culturable yeast communities***

405 *New species.*

406 The yeast that was firstly isolated from stomach of a marine fish was described as *D. hansenii*
407 and deposited in the Centraalbureau voor Schimmelcultures (CBS 5307) database. In a recent
408 paper, based on the intergenic spacer (IGS) region of the rRNA gene, this strain was re-
409 evaluated as *Candida sp.* (Nguyen *et al.*, 2009). This strain is identical to another one isolated
410 from deep-sea hydrothermal vent waters and annotated MARY089 (Gadanhó and Sampaio,
411 2005). These two strains isolated from different marine environments were finally reported as
412 a single new undescribed species within the *Candida* genus. In our collection, strain Mo39,
413 isolated from deep-sea coral near Rainbow hydrothermal vents (Table 1), has the same 26S
414 rRNA gene sequence as CBS 5307 and MARY089. Mo39 is halophilic and thus supposed to
415 be an autochthonous marine yeast species. This new ecotype can be characterized as an
416 obligate marine yeast and its complete description is currently under progress.

417

418 *Known species*

419 Two strains (Mo25 and Mo31) isolated from *Rimicaris exoculata* and *Bathymodiolus azoricus*
420 samples were identified as *Candida atlantica*. This result seems in keeping with previous
421 published reports that have isolated this species from coastal seawater in the South of
422 Portugal (Gadanhó *et al.*, 2003) and in deep-sea hydrothermal vent waters (Gadanhó and
423 Sampaio, 2005). The very first *C. atlantica* strain was isolated from shrimp eggs in the North
424 Atlantic Ocean (Siepmann and Höhnk, 1962). *C. atlantica* seemed to be a marine obligate

425 yeast and some interactions with shrimps seemed to occur. Our physiological analysis has
426 revealed that Mo25 and Mo31 were non-halophiles, which does not mean that they are unable
427 to grow in marine environments. They may have a role in deep-sea environments in
428 interaction with endemic crustaceans even if they are not in optimal growth conditions. One
429 isolate (Bio1) isolated from seawater surrounding mussels at Lau Basin in the South-West
430 Pacific was clearly identified as *Candida viswanathii*. Kohlmeyer & Kohlmeyer (1979)
431 characterized this yeast as marine facultative. More recently, *C. viswanathii* was isolated from
432 a shrimp (*Peneaus braziliensis*) in the Gulf of Mexico. Its synonym, *Candida lodderae* was
433 recently reported in deep-sea hydrothermal vent waters at Rainbow site (Gadanhó and
434 Sampaio, 2005) and characterized as the most abundant yeast.

435

436 *Leucosporidium scottii* isolates (Mo36) were retrieved only in the oceanic regions close to
437 Antarctica and are known to be psychrophilic and probably autochthonous marine species
438 (Lachance and Starmer, 1998). Such strains known for their presence in cold polar marine
439 environments could be another evidence that confirms the hypothesis of global exchanges
440 from polar environments to deep-sea vents based on results from bacteria (Maruyama *et al.*,
441 2000) and filamentous fungi (Burgaud *et al.*, 2009). *Hortaea werneckii* (Mo34) was
442 characterized as halophilic in our physiological study. This is not surprising as this black
443 yeast-like fungus was characterized as halophilic or extremely halotolerant in different studies
444 (Gunde-Cimerman *et al.*, 2000; Kogej *et al.*, 2005) where it was frequently isolated from
445 hypersaline waters of solar salterns. In a molecular survey, *H. werneckii* was identified (based
446 on internal transcribed spacers and 5.8 S rRNA gene) in deep-sea methane seep sediments at a
447 depth of 2965 meters (Lai *et al.*, 2007). *Phaeotheca triangularis* (Mo30) was also frequently
448 isolated from salted environments (Gunde-Cimerman *et al.*, 2000) and characterized as
449 halophile. This confirmed previous results on *P. triangularis* showing a better growth with
450 5% additional salts (Zalar *et al.*, 1999). In our study, Mo30 was characterized as halotolerant
451 with 4.5% sea salts optimal concentration and thus hypothesized as marine adapted yeast.
452 This is the first report about the presence of *Phaeotheca triangularis* at deep-sea vents.

453 Mo22 is described as *Sporobolomyces roseus*. The genus *Sporobolomyces* is composed of
454 strains mainly isolated from the pylophane (Bai *et al.*, 2002). However, a previous study has
455 proved that strains of the genus *Sporobolomyces* are frequently isolated from marine
456 ecosystems and the frequency of isolation increases when distance from the coastline and
457 depth increase (Hernandez-Saavedra *et al.*, 1992). Moreover, yeasts from this genus were
458 found in benthic invertebrates collected from deep-sea floor in the Pacific Ocean (Nagahama

459 *et al.*, 2001b). Our strain was isolated from a deep-sea hydrothermal shrimp in the Atlantic
460 Ocean and characterized as halotolerant with an optimal salinity of 6% sea salts. This may
461 indicate that yeasts of this genus are also able to live in deep-sea vents and interact with
462 endemic crustaceans.

463
464 A previous study of yeasts in oceanic environments (Fell, 1976) reported that yeast
465 communities appeared to be constituted of ubiquitous and endemic species. Typical
466 ubiquitous strains were the ascomycetous yeast *Debaryomyces hansenii* and the
467 basidiomycetous ones *Cryptococcus* and *Rhodotorula*. Kohlmeyer and Kohlmeyer (1979)
468 confirmed this statement and characterized these genera mainly as facultative marine yeasts.
469 Some of these results, especially for *Rhodotorula* yeasts showing a strong ubiquity, were
470 confirmed based on their presence in several habitats such as deep-sea vents (Gadanhó and
471 Sampaio, 2005), deep-sea sediments (Nagahama *et al.*, 2001b), coastal waters (Gadanhó *et*
472 *al.*, 2003; 2004) and oligotrophic lakes (Libkind *et al.*, 2003). Our results confirm their
473 ubiquity and indicate that these strains seem to be allochthonous. Strain Ex15 identified as
474 *Pichia guilliermondii* has also been characterized as non halophile and may be another
475 allochthonous yeast strain as reported by Kohlmeyer and Kohlmeyer (1979).

476
477 The members of the genus *Rhodospiridium* have been characterized as non halophiles (Mo24
478 and Mo33) and halotolerant (Mo38). Based on previous reports, this genus seemed to be
479 restricted to marine environments (Gadanhó and Sampaio, 2005). *R. diobovatum* in deep-sea
480 vents seemed to be able to colonize different substrates (shrimps, mussels and sponges). The
481 isolation of a strain from shrimp exuviae in decomposition may indicate a role as a recycler of
482 organic material and so a probable implication in carbon cycle in deep-sea environments.

483 484 ***Adaptation to marine conditions***

485 The isolation of culturable yeasts led to an old question about marine yeasts “Are there any
486 indigenous marine yeasts ?” (Kohlmeyer & Kohlmeyer, 1979) and to the resulting question
487 “Which are the indigenous species ?”. Based on our results, one can suggest that halophilic
488 strains are marine indigenous yeasts and that others, halotolerant and non-halophiles, are
489 ubiquitous terrestrial strains present in deep-sea waters due to sedimentation or other natural
490 or anthropogenic phenomena. But almost all yeast species can grow well in media with NaCl
491 concentrations exceeding those normally present in the sea (Kohlmeyer & Kohlmeyer, 1979).

492 Few yeast species with a physiological dependence on NaCl or other seawater components
493 have been reported (Nagahama, 2006b). Thus, our results appeared in good agreement with
494 such statements. Only 2 strains described as halophiles (Mo34 and Mo39) in our study can be
495 described as obligate marine yeasts.

496 ***FISH observations***

497

498 FISH using labeled oligonucleotide probes targeting rRNA has been used as a powerful
499 technique for assessing both microbial identity and spatial distributions *in situ* in complex
500 environmental contexts (Yang *et al.*, 2008). Our results indicate a very low-level of yeasts at
501 deep-sea vents. As a first conclusion, regarding diversity and quantification (added to
502 previous results of Gadanho and Sampaio, 2005), it seems that yeasts at deep-sea vents
503 represent a minor community that might not be major actors in biogeochemical cycles.
504 However, fluorescent signals are correlated to the cellular content of ribosomes and
505 consequently to the microbial growth rates. Recently, the detection limits of conventional
506 FISH with Cy3-labeled probe EUB338 were found to be approximately 370 16S rRNA
507 molecules per cell for *Escherichia coli* hybridized on glass microscope slides and 1,400 16S
508 rRNA copies per *E. coli* cell in environmental samples (Hoshino *et al.*, 2008). So, in addition
509 to a low concentration of yeast cells, low detection of yeasts may be caused by low ribosome
510 content of most yeasts in the deep-sea environment due to low-level metabolic activities of
511 yeasts living under extreme environmental abiotic factors (high hydrostatic pressure, low
512 temperatures,...). Our attempts to cultivate the yeast strains resulted from this study under
513 elevated hydrostatic pressure have been successful, but ribosomal activities were lower under
514 high hydrostatic pressure than at atmospheric pressure. Such results may account for the low
515 fungal detection using FISH (unpublished data). Consequently, care must be taken when
516 dealing with diversity and biomass estimations when using FISH alone.

517 The quantification of yeasts using FISH has been impossible due to a non homogeneous
518 repartition of microorganisms on filters. Moreover, bacterial and yeast cells were only visible
519 in some regions of the filters without minerals due to strong autofluorescence. However we
520 can say that yeast concentrations are really low, as shown by the only few cells visualized
521 after filtration of seawater surrounding shrimps. This result is in keeping with the relatively
522 low diversity revealed by Gadanho and Sampaio (2005) ranging from 0 to 10 cfu/L for pink
523 yeasts and from 0 to 6000 cfu/L for non-pigmented yeasts. To better analyze the fungal
524 presence in deep-sea animals, one could work with phylum-specific probes on histological

525 sections of animals and use the CARD-FISH (Amann & Fuchs, 2008) or the DOPE-FISH
526 (Stoecker *et al.*, 2010) methods to amplify probe signals.

527 These data raise emerging questions regarding the ecological role of such microorganisms in
528 deep-sea vents and about the old question of the ubiquity or endemism of those strains. Yeasts
529 at deep-sea vents may be facultative parasites or opportunistic pathogens of endemic deep-sea
530 animals as it has already been hypothesized in previous works (Van Dover *et al.*, 2007;
531 Burgaud *et al.*, 2009). However, a role in the decomposition of abundant organic material may
532 occur.

533 Considering all the results obtained, we can say that yeasts may seem to interact with deep-
534 sea hydrothermal endemic fauna even if the density is low. These yeasts are mainly composed
535 of ubiquitous species but obligate marine yeasts have also been harvested. However, the
536 results obtained using *in situ* hybridization have allowed us to visualize these ubiquitous species
537 showing that they are able to live and grow in deep-sea hydrothermal vents. Yeasts associated
538 with endemic animals in deep-sea vents may be exposed to favorable conditions and could
539 benefit from a stable source of nutrients (Nagahama *et al.*, 2001b). Yeasts were reported from
540 dead and healthy individuals which may also indicate their facultative saprophytism and so
541 emphasize the wide role of fungi in the decomposition of organic matter from terrestrial
542 environments to deep-sea hydrothermal vents. Even if yeasts were isolated from animal body
543 components, they were not visualized using FISH. To better understand the interaction with
544 animals and fungi in deep-sea vents, we need to work on tissues as in Van Dover *et al.* (2007)
545 and also with probes specific to fungal phyla (*Ascomycota*, *Basidiomycota* and
546 *Chytridiomycota*). In conclusion, several questions regarding the role of yeasts in deep-sea
547 hydrothermal vents and the endemism or ubiquity of the isolated yeasts remain a difficult task
548 without clear answers. Their culture under high hydrostatic pressures would be an interesting
549 study to better characterize their lifestyle and role at deep-sea vents.

550

551 **Acknowledgements**

552

553 We thank the chief scientists of the BIOLAU, DIVANAUT2, HERO, MARVEL, EXOMAR
554 and MoMARDREAM-Naut cruises, the pilots and support crews of the oceanographic vessels
555 and Deep Submergence Vehicles of Ifremer. We greatly acknowledge Dr Jerome Mounier for
556 his valuable help and comments on FISH and Jerome Lepioufle for valuable advice on the
557 manuscript. We thank all the members of the GDR Ecchis for their advice and suggestions.
558 We would like to thank the editor and the anonymous reviewers who have provided helpful

559 comments on the refinement of this manuscript. We also thank ANR Deep-Oases, Ifremer,
560 Region Bretagne and French Research Ministry for their financial support. We finally thank
561 Françoise Gail and the CHEMECO/European Science Foundation EURODEEP for
562 discussions and financial support.

563

564

565 **References**

- 566 Altschul S, Gish W, Miller W, Myers E & Lipman D (1990) Basic local alignment search tool. *J Mol Bio* **215**:
567 403–410.
568
- 569 Amann RI, Ludwig W & Schleifer K (1995) Phylogenetic identification and *in situ* detection of individual
570 microbial cells without cultivation. *Microbiol Rev*, **59**: 143-169.
571
- 572 Amann RI & Fuchs BM (2008) Single-cell identification in microbial communities by improved fluorescence in
573 situ hybridization techniques. *Nat Rev Microbiol*, **6**: 339-348.
574
- 575 Ashelford KE, Weightman AJ, & Fry JC (2002) PRIMROSE: a computer program for generating and estimating
576 the phylogenetic range of 16S rRNA oligonucleotide probes and primers in conjunction with the RDP-II
577 database. *Nucleic Acids Res* **30**: 3481-3489.
578
- 579 Bai F, Zhao JH, Takashima M, Jia JH, Boekhout T & Nakase T (2002) Reclassification of the *Sporobolomyces*
580 *roseus* and *Sporidiobolus pararoseus* complexes, with the description of *Sporobolomyces phaffii* sp. nov. *Int J*
581 *Syst Evol Microbiol* **52**: 2309-2314.
582
- 583 Baschien C, Manz W, Neu T, Marvanova L & Szewzyk U (2008) *In Situ* Detection of Freshwater Fungi in an
584 Alpine Stream by New Taxon-Specific Fluorescence In Situ Hybridization Probes. *Appl Environ Microb* **74**:
585 6427-6436.
586
- 587 Bass D, Howe A, Brown N, Barton H, Demidova M, Michelle H, Li L, Sanders H, Watkinson SC, Willcock S &
588 Richards TA (2007) Yeast forms dominate fungal diversity in the deep oceans. *Proc R Soc B* **274**: 3069-3077.
589
- 590 Burgaud G, Le Calvez T, Arzur D, Vandenkoornhuysse, P & Barbier G (2009) Diversity of culturable marine
591 filamentous fungi from deep-sea hydrothermal vents. *Environ Microbiol* **11**: 1588-1600.
592
- 593 Cancela da Fonseca JP (1966) L'outil statistique en biologie du sol. III. Indices d'intérêt écologique. *Revue*
594 *d'Ecologie et de Biologie du Sol* **3**: 381–407.
595
- 596 Charlou JL, Donval JP, Fouquet Y, Jean-Baptiste P & Holm N (2002) Geochemistry of high H₂ and CH₄ vent
597 fluids issuing from ultramafic rocks at the Rainbow hydrothermal field (36j14VN, MAR). *Chem Geol*, **191**: 345-
598 359.
599
- 600 Corbari L, Zbinden M, Cambon-Bonavita MA, Gaill F & Compère P (2008) Bacterial symbionts and mineral
601 deposits in the branchial chamber of the hydrothermal vent shrimp *Rimicaris exoculata*: Relationship to moult
602 cycle. *Aquat Biol*, **1**: 225-238.
603
- 604 Daims H, Stoecker K & Wagner M (2005) Fluorescence *In situ* Hybridisation for the Detection of Prokaryotes.
605 In: *Advanced Methods in Molecular Microbial Ecology*. BIOS Scientific Publishers, Abingdon, UK. pp. 213-
606 239.
- 607 Daims H, Lückner S, & Wagner M (2006) *daime*, a novel image analysis program for microbial ecology and
608 biofilm research. *Environ Microbiol* **8**: 200-213.
609
- 610 Edgcomb VP, Kysela DT, Teske A, & de Vera Gomez A (2002) Benthic eukaryotic diversity in the Guaymas
611 Basin hydrothermal vent environment. *Proc Natl Acad Sci USA* **99**: 7658-7662.
612
- 613 Fell JW (1976) Yeasts in oceanic regions. In: Jones EBG (ed) *Recent advances in aquatic mycology*. Elec,
614 London, pp 93-124.
- 615 Gadanho M, Almeida JM, & Sampaio JP (2003) Assessment of yeast diversity in a marine environment in the
616 south of Portugal by microsatellite-primed PCR. *Antonie van Leeuwenhoek* **84**: 217-227.
617
- 618 Gadanho M & Sampaio JP (2004) Application of temperature gradient gel electrophoresis to the study of yeast
619 diversity in the estuary of the Tagus river, Portugal. *FEMS Yeast Res* **5**: 253-261.
620
- 621 Gadanho M & Sampaio J (2005) Occurrence and Diversity of Yeasts in the Mid-Atlantic Ridge Hydrothermal
622 Fields Near the Azores Archipelago. *Microb Ecol* **50**: 408-417.
623
- 624 Gadd GM (2007) Geomycology: biogeochemical transformations of rocks, minerals and radionuclides by fungi,
625 bioweathering and bioremediation. *Mycol Res* **111**: 3-49.
626
- 627 Guindon S, Lethiec F, Duroux P & Gascuel O (2005) PHYML online – a web server for fast maximum
628 likelihoodbased phylogenetic inference. *Nucleic Acids Res* **33**: 557–559.
629

- 630 Gunde-Cimerman N, Zalar P, de Hoog S & Plemenitas A (2000) Hypersaline waters in salterns - natural
631 ecological niches for halophilic black yeasts. *FEMS Microbiol Ecol* **32**: 235-240.
632
- 633 Hawksworth DL (2002) The magnitude of fungal diversity: the 1.5 million species estimate revisited. *Mycol Res*
634 **105**: 1422-1432.
635
- 636 Hernandez-Saavedra NY, Hernandez- Saavedra D & Ochoa JL (1992) Distribution of *Sporobolomyces* (Kluyver
637 et van Niel) genus in the western coast of Baja California Sur, Mexico. *Syst Appl Microbiol* **15**: 319-322.
638
- 639 Hoshino T, Yilmaz LS, Noguera DR, Daims H & Wagner M (2008) Quantification of Target Molecules Needed
640 To Detect Microorganisms by Fluorescence In Situ Hybridization (FISH) and Catalyzed Reporter Deposition-
641 FISH. *Appl Environ Microb* **74**: 5068-5077.
642
- 643 Joergensen BB, & Boetius A (2007) Feast and famine-microbial life in the deep-sea bed. *Nat Rev Microbiol* **5**:
644 770-781.
645
- 646 Kogej T, Ramos J, Plemenitas A, & Gunde-Cimerman N (2005) The halophilic fungus *Hortaea werneckii* and
647 the halotolerant fungus *Aureobasidium pullulans* maintain low intracellular cation concentrations in hypersaline
648 environments. *Appl Environ Microb* **71**: 6600-6605.
649
- 650 Kohlmeyer J & Kohlmeyer E (1979) Marine Mycology: The Higher Fungi. New-York, USA; Academic Press.
651
- 652 Inacio J, Behrens S, Fuchs BM, Fonseca A, Spencer-Martins I & Amann R (2003) In situ accessibility of
653 *Saccharomyces cerevisiae* 26S rRNA to Cy3-labeled oligonucleotide probes comprising the D1 and D2
654 domains. *Appl Environ Microb* **69**: 2899-2905.
- 655 Kushner DJ (1978) Life in high salt and solute concentrations. In: Kushner DJ (ed) *Microbial life in extreme*
656 *environments*. Academic Press, London, pp 317-368.
- 657 Lachance MA & Starmer WT (1998) Ecology and yeasts. In: Kurtzman CP, Fell JW (eds) *The yeasts, a*
658 *taxonomic study*, 4th edn. Elsevier, Amsterdam, The Netherlands, pp 21-30.
- 659 Lai X, Cao L, Tan H, Fang S, Huang Y & Zhou S (2007) Fungal communities from methane hydrate-bearing
660 deep-sea marine sediments in South China Sea. *ISME J* **1**: 756-762.
661
- 662 Le Calvez T, Burgaud G, Mahe S, Barbier G & Vandenkoornhuysen P (2009) Fungal Diversity in Deep Sea
663 Hydrothermal Ecosystems. *Appl Environ Microb* **75**: 6415-6421.
664
- 665 Libkind D, Brizzio S, Ruffini A, Gadanho M, van Broock M & Sampaio JP (2003) Molecular characterization of
666 carotenogenic yeasts from aquatic environments in Patagonia, Argentina. *Antonie van Leeuwenhoek* **84**: 313-
667 322.
668
- 669 Lopez-Garcia P, Philippe H, Gail F & Moreira D (2003) Autochthonous eukaryotic diversity in hydrothermal
670 sediment and experimental microcolonizers at the Mid-Atlantic Ridge. *Proc Natl Acad Sci USA* **100**: 697-702.
671
- 672 Lopez-Garcia P, Vereshchaka A & Moreira D (2007) Eukaryotic diversity associated with carbonates and fluid-
673 seawater interface in Lost City hydrothermal field. *Environ Microbiol* **9**: 546-554.
674
- 675 Margesin R & Schinner F (2001) Potential of halotolerant and halophilic microorganisms for biotechnology.
676 *Extremophiles* **5**: 73-83.
677
- 678 Maruyama A, Honda D, Yamamoto H, Kitamura K & Higashihara T (2000) Phylogenetic analysis of
679 psychrophilic bacteria from the Japan Trench, including a description of the deep-sea species *Psychrobacter*
680 *pacificensis* sp. nov. *Int J Syst Evol Microbiol* **50**: 835-846.
681
- 682 Munn, CB (2003) *Marine Microbiology – Ecology and Applications*. Oxford, UK: Bios-Garland Scientific.
683
- 684 Nagahama T, Hamamoto M, Nakase T & Horikoshi K (1999) *Kluyveromyces nonfermentans* sp. nov., a new
685 yeast species isolated from the deep sea. *Int J Syst Evol Microbiol* **49**: 1899-1905.
686
- 687 Nagahama T, Hamamoto M, Nakase T & Horikoshi K (2001a) *Rhodotorula lamellibrachii* sp. nov., a new yeast
688 species from a tubeworm collected at the deep-sea floor in Sagami Bay and its phylogenetic analysis. *Antonie*
689 *van Leeuwenhoek* **80**: 317-323.
690
- 691 Nagahama T, Hamamoto M, Nakase T, Takami H & Horikoshi K (2001b) Distribution and identification of red
692 yeasts in deep-sea environments around the northwest Pacific Ocean. *Antonie van Leeuwenhoek* **80**: 101-110.
693

694 Nagahama T, Hamamoto M, Nakase T, Takaki Y & Horikoshi K (2003a) *Cryptococcus surugaensis* sp. nov., a
695 novel yeast species from sediment collected on the floor of Suruga Bay. *Int J Syst Evol Microbiol* **53**: 2095-
696 2098.

697
698 Nagahama T, Hamamoto M, Nakase T & Horikoshi K (2003b) *Rhodotorula benthica* sp. nov. and *Rhodotorula*
699 *calyptogenae* sp. nov., novel yeast species from animals collected from the deep-sea floor, and *Rhodotorula*
700 *lysiniphila* sp. nov., which is related phylogenetically. *Int J Syst Evol Microbiol* **53**: 897-903.

701
702 Nagahama T, Hamamoto M & Horikoshi K (2006a) *Rhodotorula pacifica* sp. nov., a novel yeast species from
703 sediment collected on the deep-sea floor of the north-west Pacific Ocean. *Int J Syst Evol Microbiol* **56**: 295-299.

704
705 Nagahama T (2006b) Yeast biodiversity in freshwater, marine and deep-sea environments. *Biodiversity and*
706 *Ecophysiology of Yeasts*.

707
708 Nagahama T, Abdel-Wahab MA, Nogi Y, Miyazaki M, Uematsu K, Hamamoto M & Horikoshi K (2008)
709 *Dipodascus tetrasporeus* sp. nov., an ascosporegenous yeast isolated from deep-sea sediments in the Japan
710 Trench. *Int J Syst Evol Microbiol* **58**: 1040-1046.

711
712 Nguyen H, Gaillardin C & Neuvéglise C (2009) Differentiation of *Debaryomyces hansenii* and *Candida famata*
713 by rRNA gene intergenic spacer fingerprinting and reassessment of phylogenetic relationships among *D.*
714 *hansenii*, *C. famata*, *D. fabryi*, *C. flareri* (= *D. subglobosus*) and *D. prosopidis*: description of *D. vietnamensis* sp.
715 nov. closely related to *D. nepalensis*. *FEMS Yeast Res* **9**: 641-662.

716
717 Posada D & Crandall K (1998) Applications note. MODELTEST: testing the model of DNA substitution.
718 *Bioinformatics* **14**: 817-818.

719
720 Ronquist F & Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models.
721 *Bioinformatics* **19**: 1572-1574

722
723 Siepmann R & Höhnk W (1962) Über Hefen und einige Pilze (Fungi imp., Hyphales) aus dem Nordatlantik.
724 *Veroeff Inst für Meeresforschung in Bremerhaven* **8**: 79-97.

725
726 Stoecker K, Dorninger C, Daims H & Wagner M (2010) Double Labeling of Oligonucleotide Probes for
727 Fluorescence *In Situ* Hybridization (DOPE-FISH) Improves Signal Intensity and Increases rRNA Accessibility.
728 *Appl Environ Microb* **76**: 922-926.

729
730 Takishita K, Tsuchiya M, Reimer JD & Maruyama T (2006) Molecular evidence demonstrating the
731 basidiomycetous fungus *Cryptococcus curvatus* is the dominant microbial eukaryote in sediment at the
732 Kuroshima Knoll methane seep. *Extremophiles* **10**: 165-169.

733
734 Takishita K, Yubuki N, Kakizoe N, Inagaki Y & Maruyama T (2007) Diversity of microbial eukaryotes in
735 sediment at a deep-sea methane cold seep: surveys of ribosomal DNA libraries from raw sediment samples and
736 two enrichment cultures. *Extremophiles* **11**: 563-576.

737
738 Tamura K, Dudley J, Nei M & Kumar, S (2007) MEGA4: molecular evolutionary genetics analysis (MEGA)
739 software, version 4.0. *Mol Biol Evol* **24**: 1596-1599.

740
741 Tivey, MK (2007) Generation of seafloor hydrothermal vent fluids and associated mineral deposits.
742 *Oceanography* **20**: 50-65.

743
744 Thompson JD, Higgins DG & Gibson, T.J. (1994) CLUSTAL W: improving the sensitivity of progressive
745 multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix
746 choice. *Nucleic Acids Res* **22**: 4673-4680.

747
748 Van Dover CL, Ward ME, Scott JL, Underdown J, Anderson B, Gustafson C, Whalen M & Carnegie RA (2007)
749 A fungal epizootic in mussels at a deep-sea hydrothermal vent. *Mar Ecol* **28**: 54-62.

750
751 Zalar P, de Hoog GS & Gunde-Cimerman N (1999) Ecology of halotolerant dothideaceous black yeasts. *Stud*
752 *Mycol* **43**: 38-48.

753
754 Zbinden M, Le Bris N, Gaill F & Compere P (2004) Distribution of bacteria and associated minerals in the gill
755 chamber of the vent shrimp *Rimicaris exoculata* and related biogeochemical processes. *Mar Ecol Progr Ser* **284**:
756 237-251.

757

758 **Tables and Figures**

759

760 Table 1. Culture collection of yeasts from deep-sea hydrothermal vents.

761

762

763

764

765

766

767	Location (Depth)	Sample processed (type)	Strain
768	South Pacific West; (Lau Basin; -2620m)	B2E07: Seawater surrounding mussels	Bio1
769		B9E07: Gastropod (<i>Ifremeria nautiliei</i>) gills	Bio2
770	Mid-Atlantic Ridge (Rainbow; -2300m)	EX6E01 to EX6E04: <i>Rimicaris exoculata</i>	Ex2 to Ex7
771		EX6E05: <i>Chorocaris chacei</i>	Ex9, Ex11 and Ex12
772		MoPR1: <i>Rimicaris exoculata</i>	Mo20
773		MoPR1: <i>Mirocaris fortunata</i>	Mo21
774		MoPR2: <i>Rimicaris exoculata</i>	Mo22
775		MoPR3: Sloughs of shrimp on smocker rocks	Mo24 and Mo25
776		MoPR5: Colonization module TRAC (Carbonates)	Mo26 to Mo29
777		MoPR6: <i>Bathymodiolus azoricus</i>	Mo30 to Mo36
778		MoPR8: <i>Rimicaris exoculata</i>	Mo37
779		MoPR9: Sponge	Mo38 and Mo39
780	Mid-Atlantic Ridge (Lost-City; -700m)	MoPR9: Coral	Mo40
781		EX18E02: Siliceous sponge	Ex15
782			
783			
784			
785			
786			
787			
788			
789			
790			
791			
792			
793			
794			
795			

796 Table 2. Physiological analysis of the yeast collection. This table shows distribution of
797 halotolerant and halophilic strains of the collection depending on their optimal salinities (g/l
798 sea salts), optimal temperatures (°C) and maximal optical densities of cultures on GYPS broth
799 medium (120 rpm on a rotary shaker) measured at 600nm at 4 different incubation times (17h,
800 22h, 25h and 28h).

801

802

803

		Low OD (<1.1)	High OD (>2.0)	
	Optimum	25°C	25°C	35°C
Non halophile	0-15 g/l	Mo25	Bio1, Bio2, Ex2, Ex3, Ex4, Ex5, Ex6, Ex7, Ex9, Ex11, Ex12, Mo20, Mo21, Mo24, Mo26, Mo27, Mo28, Mo29, Mo31, Mo32, Mo33, Mo35, Mo40	Ex15
Halotolerant	30 g/l	Mo36	Mo37, Mo38	
	45 g/l			Mo30
	60 g/l	Mo22		
Halophile	30 g/l	Mo34		Mo39

804

805

806 Table 3. Yeast oligonucleotide probes and their sequences, target organisms and binding
807 positions on the 26S rRNA.

808

809

810

811

Probe	Hybridization stringency (% formamide)	rRNA subunit, binding position ^(a) and relative probe accessibility ^(b)	Probe sequence (5'-3')	Target organisms (Genus/Species)
Sacch	20	26S; 162-177 ; 44 to 66%	GGCATCTCATCGCACG	<i>Debaryomyces</i> <i>Pichia</i>
MitoFilo	10	26S; 397-412 ; 60%	ACACCGCAGAACGGCT	Members of the genus <i>Cryptococcus</i> ^(c)
MitoSporidio	20	26S; 164-179 ; 44 to 66%	TGGGCGTCCGCACCAT	Members of the genera <i>Rhodotorula</i> and <i>Rhodospiridium</i> ^(d)

(a) Nucleotide position according to *Saccharomyces cerevisiae* 26Sr RNA between NL1 and NL4 primers.

(b) According to Inacio *et al.*, 2003.

(c) *Cryptococcus saitoi*, *C. randhawii*, *C. uzbekistanensis*, *C. adeliensis*, *C. vishniacii*, *C. socialis*, *C. friedmannii* and *C. uniguttulatus*.

(d) *Rhodotorula mucilaginosa*, *R. glutinis*, *R. graminis*, *R. dairenensis*, *Rhodospiridium babjevae* and *R. diobovatum*.

Figure 1: Phylogenetic tree of deep-sea yeast isolates (coloured terminals) and close relatives obtained by analysis of the D1/D2 domain of the 26S rRNA gene. Topology was built using MrBayes v.3.1.2 from a ClustalW 1.83 alignment. Node support values are given in the following order: MrBayes posterior probabilities/PHYML 100 bootstraps. Black squares represent nodes supported by an excess of 0.95 posterior probabilities and 95% bootstraps. *Mucor flavus* (EU071390) belonging to the *Zygomycota* phylum was used as outgroup. All sequences are listed with their GenBank accession numbers.




Figure 2: Fluorescence *in situ* hybridization with specific oligonucleotide probes on membrane filters. (a, b and c) Membrane filter labelled with DAPI and hybridized using MitoSporidio probe indicating the presence of bacteria and yeast cells (blue). Yeasts belonging to *Rhodotorula* and *Rhodospiridium* genera are visualized in pink (composite of blue and red). (d) Membrane filter labelled with DAPI and hybridized with MitoFilo indicating the presence of yeasts belonging to *Cryptococcus* genera. White arrows indicate the yeast cells.






Figure S1 : Target sites of the fluorescent oligonucleotide probes designed on a model of the *Saccharomyces cerevisiae* 26S rRNA secondary structure in which the D1 and D2 domains (delimited by NL1 and NL4) are enlarged (Inacio *et al.*, 2003). Each probe was evaluated without formamide in order to check whether the probe binds to the ribosomes of the target cells. The optimal hybridization conditions were determined in a series of FISH experiments with increasing formamide concentrations for a probe target and a non-target organism : (i) Sacch probe, *Debaryomyces hansenii* (Target) and *Candida atlantica* (Non-Target) with two mismatches ; (ii) MitoSporidio probe, *Rhodospiridium diobovatum* (Target) and *Cryptococcus uzbekistanensis* (Non-Target) with five mismatches and (iii) MitoFilo probe, *Cryptococcus uzbekistanensis* (Target) and *Rhodospiridium diobovatum* (Non-Target) with seven mismatches. Relative probe accessibility was determined for each probe : MitoFilo, about 60% ; MitoSporidio and Sacch, 44 to 66%.

