



HAL
open science

Transcriptomic Response of *Enterococcus faecalis* V583 to Low Hydrogen Peroxide Levels.

Xue Yan, Aurélie Budin-Verneuil, Nicolas Verneuil, Michael S Gilmore,
Sébastien Artigaud, Yanick Auffray, Vianney Pichereau

► **To cite this version:**

Xue Yan, Aurélie Budin-Verneuil, Nicolas Verneuil, Michael S Gilmore, Sébastien Artigaud, et al..
Transcriptomic Response of *Enterococcus faecalis* V583 to Low Hydrogen Peroxide Levels.. *Current
Microbiology*, 2014, 70 (2), pp.156-168. 10.1007/s00284-014-0691-8 . hal-01089248

HAL Id: hal-01089248

<https://hal.univ-brest.fr/hal-01089248>

Submitted on 5 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Transcriptomic response of *Enterococcus faecalis* V583 to low hydrogen peroxide levels

Yan Xue ¹, Budin-Verneuil Aurelie ¹, Verneuil Nicolas ¹, Gilmore Michael S. ², Artigaud Sebastien ³, Auffray Yannick ¹, Pichereau Vianney ^{1,3,*}

¹ Univ Caen Basse Normandie, Unite Rech Risques Microbiens U2RM, Equipe Stress Virulence, F-14032 Caen, France.

² Harvard Univ, Sch Med, Dept Ophthalmol, Boston, MA USA.

³ Univ Bretagne Occidentale UEB, Lab Sci Environm Marin LEMAR, Inst Univ Europeen Mer, IFREMER,IRD,UBO,CNRS,UMR 6539, F-29280 Plouzane, France.

* Corresponding author : Vianney Pichereau, email address : vianney.pichereau@univ-brest.fr

Abstract :

Enterococcus faecalis is a Gram-positive commensal bacterium inhabiting the gastrointestinal tracts of human and other mammals, but is also increasingly recognized as an opportunistic human pathogen. Oxidative stress is one of the major challenges encountered by enterococci, both in their natural environment and during infection. In this paper, we evaluated the transcriptomic response of *E. faecalis* to oxidative stress, and showed that transcript abundance was reduced for 93 genes and increased for 39 genes during growth in medium containing 1.75 mM H₂O₂. The presence of hydrogen peroxide affected several metabolic pathways, including a large decrease in ethanolamine utilization and methylglyoxal metabolism, and an increase in transcript abundance for several transport systems. In particular, four operons encoding iron transporters appeared highly induced. By contrast, in our experimental conditions, the expression of most of the genes known to be involved in the enterococcal response to oxidative stress, did not appear significantly altered.

45 **1. Introduction**

46 *Enterococcus faecalis* is a Lactic Acid Bacterium naturally inhabiting the gastrointestinal
47 tracts of humans and other mammals, and used as a constituent of some probiotic food
48 supplements [17,18,30]. However, *E. faecalis* also can cause life-threatening infections in
49 humans, especially in the hospital environment [35,46]. Oxidative stress is one of the major
50 challenges encountered by *E. faecalis* during its life cycle [23,34,50]. Nearly two decades of
51 research have revealed that this bacterium possesses genes encoding antioxidant enzymes
52 such as *sodA* (Superoxide dismutase, [41]), *kata* (Catalase, [14]), *gor* (Glutathione reductase),
53 *nox* (NADH oxidase), *trx* (Thioredoxine reductase), *dps* (DNA-binding protein), *msrA*
54 (Peptide methionine-S-sulfoxide reductase) [39], *npr* (NADH peroxidase, [44]), *ohr* (Organic
55 hydroperoxide resistance protein, [43]), *ahpC* (Alkyl hydroperoxide reductase, [57]), *ahpF*
56 (Peroxiredoxine reductase), and several transcriptional regulators involved in the oxidative
57 stress response, as *hypR* [55], *perR*, *fur*, *zur* [56], *soxS* [54].

58 Global experimental approaches that allow monitoring of gene and protein expression (i.e.,
59 transcriptomics and proteomics, respectively) in response to a particular environmental
60 condition, constitute powerful tools to understand the metabolic and regulatory networks
61 involved in stress adaptation on a genome-wide scale. To date, only one proteomic study
62 examined changes in expression of 23 proteins in *E. faecalis* cells exposed to hydrogen
63 peroxide (H₂O₂) [12]. In this paper, we report global changes in transcript abundance in *E.*
64 *faecalis* V583 during growth in the presence of H₂O₂.

65

66 **2. Materials and methods**

67 *2.1. Bacterial strains and growth conditions*

68 In this work, we used *E. faecalis* V583 ery^S, an erythromycin sensitive derivative of the
69 clinical isolate *E. faecalis* V583 [45]. Cells were cultivated with shaking in M17 medium
70 supplemented with 0.5% (w/vol) glucose (GM17), supplemented with ± 1.75 mM H₂O₂.

71 *Microarray experiments*

72 V583 ery^S growth was monitored spectrophotometrically at 600 nm. When the culture
73 reached mid-exponential phase (Fig. 1), cells were harvested and total RNA was extracted,
74 purified, and residual DNA enzymatically removed as described previously [59]. Two
75 independent experiments were performed for each condition. cDNA preparation,
76 fragmentation, labeling and hybridization were performed as described in the Affymetrix
77 manual ("GeneChip Expression Analysis Technical Manual: Procaryotic Target preparation"
78 P/N 702232 Rev 2). Washing and scanning were performed using a GeneChip® Fluidics
79 Station 450. The arrays were read at 570 nm with a resolution of 1.56 μ m using an Affymetrix
80 GeneChip Scanner 3000 7G (Affymetrix). Results were analyzed and compared using the
81 GeneChip Operating Software (GCOS) version 1.4. Data were also analysed using R (version
82 3.0.0; R core team, 2013) and the Bioconductor software (version 2.12; [15]), to perform a
83 t-test with a Benjamini and Hochberg multiple testing correction cut-off of $P < 0.05$ (data not
84 shown). In all, we considered differences in transcript abundance significant if by GCOS they
85 exhibited both a p-value below 0.05, and an absolute expression ratio significantly greater
86 than 2 fold.

87 *2.3. Real-Time quantitative PCR*

88 Real-Time qPCR was performed exactly as described by Giard *et al.* [16]. Specific primers
89 were designed using the Primer3 software available at the web site
90 http://frodo.wi.mit.edu/cgi-bin/primer3/primer3_www.cgi (Table 1) with the following

91 parameters: amplicon length (99 to 101 bp), primer length (19 to 21 nucleotides), and primer
92 melting temperature (T_m from 59 to 61 °C). Total RNAs, treated with DNase I (GE
93 Healthcare) and purified using the DNA-free kit (Ambion), were reverse transcribed using the
94 Omniscript enzyme (Qiagen) and random hexamer primers according to the manufacturer's
95 recommendations. RT qPCR was performed using the QuantiTect SYBR Green RT PCR kit
96 (Qiagen). Quantification of 23S rRNA transcript levels was used as an internal control. All
97 experiments were performed twice, and in duplicate with two different RNA samples, using
98 the BioRad iCycler iQ detection system (Bio-Rad Laboratories, Richmond, CA). The value
99 used for the comparison of genes expressions in stressed and unstressed cells was the number
100 of PCR cycles required to reach the threshold cycle (C_T). To relate the C_T value to the
101 abundance of an mRNA species, C_T was converted to “ n -fold difference” by comparing
102 mRNA abundance in the control cells to that obtained with the H_2O_2 stressed cells. The n -fold
103 difference was calculated by the formula ($n = 2^{-x}$) when the $C_{T \text{ mutant}} < C_{T \text{ V583}}$ and ($n = -2^x$)
104 when $C_{T \text{ mutant}} > C_{T \text{ V583}}$, with $x = (C_{T \text{ mutant}} - C_{T \text{ V583}})$.

105

106 **3. Results and discussion**

107 *3.1. Growth of E. faecalis V583 ery^S under oxidative stress.*

108 *E. faecalis* cells were cultured in the presence of increasing concentrations of H_2O_2 , from 1.5
109 mM to 2 mM to identify a condition where H_2O_2 had a detectable effect on growth, but where
110 cells were also able to successfully respond to the challenge. Results presented in the Fig. 1
111 showed that the inhibitory effect of H_2O_2 was mainly observable on the duration of the Lag
112 phase, which increased from approx. 30 min to 3 h in the presence of 1.5 mM to 2 mM H_2O_2 .
113 We aimed at determining the effects of low doses of H_2O_2 on the global genes expression
114 profiles of exponentially growing *E. faecalis* cells, which were previously shown to display
115 differences in stress tolerance [12]. Therefore, total RNAs were extracted from

116 mid-exponential growth phase cells (Fig. 1), and then processed for transcriptome analyses.

117 *3.2. Transcriptome changes in response to H₂O₂ exposure.*

118 Whole genome expression profiling of *E. faecalis* V583 was performed using affymetrix
119 chips, using total RNAs extracted from both unstressed- and stressed- (1.75 mM H₂O₂)
120 cultures at an OD_{600nm} of 0.6 (reached two and four hours after inoculation, respectively, Fig.
121 1). Comparative analysis showed that 132 genes were significantly differentially expressed in
122 response to 1.75 mM H₂O₂, considering a ratio threshold of two fold (Tables 2 and 3). We
123 selected 25 of the mostly deregulated genes, and subjected them to RT-qPCR (Table 4). It is
124 noteworthy that we showed a very strong correlation ($r^2 = 0.86$) between the changes in genes
125 expression observed with the whole genome transcriptomic- and the qPCR- experimental
126 approaches.

127 Among the 132 genes for which transcript abundance changes significantly, 93 showed
128 reduction and 39 were found to be increased. We classified these 132 genes into 9 groups
129 according to the categories described in the comprehensive microbial resource of The Institute
130 of Genome Research (<http://www.tigr.org/tigr-scripts/CMR2/CMRHomePage.spl/>). Figure 2
131 shows the number of differentially regulated genes in each functional class. The main
132 observation is the large dominance of genes encoding proteins for which the function is
133 annotated as unknown (60), and the abundance of genes encoding (i) transport and binding
134 proteins (22 genes increased and 3 reduced), (ii) proteins involved in biosynthesis and
135 metabolism (4 increased and 24 reduced), and (iii) proteins involved in regulation and cell
136 signaling (8). Transcripts for only two genes annotated as involved in stress adaptation and
137 virulence were found to be changed.

138 *3.3. Classification of genes by their regulation patterns.*

139 Table 2 and Table 3 identify genes associated with increased and decreased levels of mRNA,
140 respectively, as a result of exposure to 1.75 mM H₂O₂. The most highly affected transcripts

141 are changed by about 10 fold. For example, the *ef3082-ef3085* operon encoding the 4
142 components of an iron compound ABC transport system, displayed increased transcript
143 abundance from 9 to 11 fold, while the greatest decreases all related to ethanolamine
144 metabolism, and displayed a reduction in transcript abundance reaching almost 12 fold.

145 To understand these changes in the global context of cellular metabolism, we searched for
146 functional association networks using the STRING program ([13], available at
147 <http://string-db.org/>). Results presented in Fig. 3 show that of the 132 genes for which
148 significant transcript changes were observed, only 14 were disconnected from any node, and
149 only 22 belonged to networks that included less than 4 partners (thus showing that 73% of the
150 genes affected by H₂O₂ stress belonged to larger networks).

151 Gene ontology enrichment analyses showed that the main biological processes and molecular
152 functions associated with global transcriptional change, again were those related to metal ion
153 transport and ethanolamine metabolism P-values < 10⁻⁵ for each).

154 3.3.1. Metabolism

155 Ethanolamine metabolism

156 Both microarray and real time q-PCR (Table 3 and Table 4) showed that H₂O₂ decreased the
157 the abundance of transcripts for 18 genes belonging to the *eut* operon involved in
158 ethanolamine metabolism. Ethanolamine (EA) is an abundant compound in the human
159 intestinal tract [24,26] and in processed food [7], and can be used as a source of carbon,
160 nitrogen, and energy. However, growth of the V583 strain, and of RR17 (*ef1633*) a null
161 mutant of *ef1633*, showed that EA provided with CoB12 allowed *E. faecalis* to grow, albeit
162 poorly, only under anaerobic conditions, and the RR17 response regulator was required for
163 this effect. A promoter upstream of RR17 is induced by EA-CoB12 in anaerobiosis [38]. The
164 observation that most of the genes from EF1617 (*eutQ*) to EF1638 (*eutP*) were highly
165 repressed by H₂O₂ in this study, could stem from a negative impact of H₂O₂ on either EA

166 and/or CoB12 in the medium, and reflects an important modulation of metabolism in
167 enterococcal cells in response to oxidative stress. It is noteworthy that the *Eut* system was also
168 shown to be negatively deregulated in (i) a microarray study of the *Fsr* system [5], and (ii) in
169 *relA* mutants unable to produce (p)ppGpp [59,60] and unpublished data), showing that the
170 regulation of the *Eut* operon is governed by many inputs. This new data adds to the
171 understanding of the role of this system in the physiology of bacteria subjected to changing
172 environments.

173 Methylglyoxal metabolism

174 The production of MG (methylglyoxal) in bacterial cells is maintained in balance with the
175 capacity for detoxification and protection against this toxic electrophile, which can react with
176 the nucleophilic centres of macromolecules such as DNA, RNA and proteins [4,11]. Table 3
177 and Table 4 show that the abundance of transcripts for 3 genes encoding glyoxalase family
178 proteins (among the 9 genes identified in the genome with related function) is significantly
179 reduced during the growth in the presence of H₂O₂. It should be noted that all other
180 glyoxalase genes, and the *ef0939* gene encoding the methyl glyoxal synthase, were also
181 negatively impacted, but by differences < 2 fold (Table 5). This indicates an important role
182 for methylglyoxal metabolism under this stress. In enteric bacteria, three methylglyoxal
183 detoxification routes have been identified, *i.e.* the glutathione-dependent glyoxalase I-II
184 system [19], (ii) glutathione-independent glyoxalase III [31], and (iii) methylglyoxal
185 reductases and dehydrogenases [19,32]. The 5 glyoxalase genes for which mRNA abundance
186 decreased in the presence of H₂O₂ share sequence identity with genes of the glyoxalase I-II
187 system. Although methylglyoxal is considered as an extremely toxic electrophile, potentially
188 leading to the death of bacterial cell, several studies suggest that this compound is important
189 for bacterial homeostasis. Indeed, methylglyoxal is thought to be important in bacteria for
190 growth regulation, D-lactate production [8], the uncoupling of anabolism and catabolism [49],

191 the prevention of accumulation of phosphorylated intermediates [6,21], virulence [3],
192 programmed cell death [22,37], and oxidative stress [58]. Therefore, the reduction in the
193 abundance of transcripts encoding cellular glyoxalases suggests that tuning of methylglyoxal
194 intracellular content in H₂O₂ stressed cells is important in *E. faecalis* oxidative stress
195 adaptation.

196 3.3.2. Transport

197 Iron uptake

198 Table 2 and Table 4 show a notable increase in mRNA encoded by 10 genes belonging to 4
199 putative operons, with functions related to iron transport. Iron is an essential element for most
200 microbes, since many enzymes use iron as a cofactor [1]. During pathogenic processes, iron
201 acquisition systems are often found to be highly expressed to compensate for low iron
202 availability within the host [47]. Interestingly, the greatest increase in mRNA encoding an
203 iron transporter in our study (ef3082-3085), was also shown to be affected by growth in blood
204 [53]. Conversely, the expression of the *mntH* gene (ef1057), that was recently shown to be
205 strongly repressed in response to iron excess in *E. faecalis* OG1-RF [29], was not modified in
206 our experimental conditions. There is also an intimate relationship between iron metabolism
207 and oxidative stress. First, through the Fenton/Haber-Weiss reaction, iron can promote the
208 formation of hydroxyl radicals, which indiscriminately damage all cellular components [9,51].
209 Second, iron acts as a co-factor for several enzymes, including some involved in oxidative
210 stress defence, such as KatA or Dps.

211 The induction by H₂O₂ of genes with functions in iron uptake was also observed in *Bacillus*
212 *subtilis*, and could reflect an adaptive response to iron limitation [33]. Taken together, our
213 results suggest that H₂O₂-stressed cells encounter an intracellular iron limitation that could
214 result from iron recruitment by certain oxidative defence proteins (*e.g.* catalase), necessitating
215 increased transport capacity to meet the competing intracellular iron needs.

216 Stress related transporters

217 We observed in Table 2 and Table 3 that hydrogen peroxide resulted in increased abundance
218 of transcripts of *kdpA*, *kdpB*, *kdpC*, *kdpD*, encoding the high affinity ABC potassium transport
219 system Kdp, and *ef0575-ef0576* (cationic ABC transporter). At the same time, H₂O₂ exposure
220 resulted in reduced levels of *cadA* (cadmium-translocating P-type ATPase), *ef0986* (cation
221 transporter) and *nhaC* (Na⁺/H⁺ antiporter) mRNA. Select changes were verified by RT-qPCR
222 (Table 4). All bacterial *kdp* operons investigated so far are repressed during growth in media
223 of high external K⁺ concentration ([K⁺]_e) [2]. Increased abundance of mRNA encoding the
224 *kdp* operon under our experimental condition, suggests that H₂O₂ led to an intracellular K⁺
225 limitation. Thus, the Kdp-ATPase is an efficient K⁺ scavenging system that is expressed when
226 other K⁺ transporters cannot support cellular requirement for K⁺, which not only plays a vital
227 role in bacterial osmotic adaptation [10,40] but is also important for pH regulation, gene
228 expression and activation of cellular enzymes [42]. The most notable deficiency observed in
229 the *nhaC* deletion strain of *Bacillus firmus* was its poor growth at pH 7.5 and Na⁺
230 concentrations up to 25 mM [20], suggesting its contribution in osmotic stress. The cadmium
231 cation is toxic to most microorganisms, probably by binding to essential respiratory proteins
232 [52] and through oxidative damage by production of reactive oxygen species [48]. One of the
233 best-characterized bacterial cadmium resistance mechanisms is determined by the
234 cadmium-transporting ATPase encoded by *cadA* found in *Staphylococcus aureus* [36] and
235 *Listeria monocytogenes* [27,28]. Various stresses (NaCl, heat, ethanol, acidity and alkalinity)
236 induced weak or strong H₂O₂ cross-protection in *E. faecalis* ATCC19433 [12]. Combined
237 results suggest that the pre-treatment of H₂O₂ could increase the bacterial osmotolerance and
238 resistance to antibiotic, and decrease the bacterial resistance to cadmium.

239 3.3.3. Genes involved in the adaptation to oxidative stress

240 Expression of other genes clearly involved in the adaptation to oxidative stress did not

241 significantly change in GM17 medium supplemented with 1.75 mM H₂O₂ (Table 6). These
242 genes may already be expressed at a high level in laboratory growth. Interestingly, although
243 not reaching the criteria used for significance in this study, the abundance of transcripts
244 encoding *msrB* (Peptide methionine-S- sulfoxide reductase [25]) was reduced 1.65 fold by
245 H₂O₂. The repression of *msrB* in 2.5 mM H₂O₂ adaptation for 30 minutes and the resistance of
246 Δ *msrB* double crossing over mutant to the lethal challenge of 7 mM H₂O₂ for 6 hours
247 observed in *E. faecalis*, suggests that *msrB* contributes negatively to adaptation to oxidative
248 stress [61]. It therefore was not surprising that when *E. faecalis* is facing the oxidative stress
249 of H₂O₂, a reduction in transcripts for *msrB* was observed.

250 3.4. Concluding remarks

251 Global changes in the abundance of mRNA for all genes in exponentially growing *E. faecalis*
252 cells, cultured in the presence of a low dose (1.75 mM) of H₂O₂, was measured. At this time
253 (4 h), *E. faecalis* had almost completely adapted as evinced by restoration of its growth rate
254 (Fig. 1). Analysis of changes in transcript abundance in response to this stress revealed that (i)
255 the expression of most of the well-known oxidative stress genes of *E. faecalis* was not
256 significantly altered, possibly because they already were highly expressed in rapid growth in
257 laboratory conditions, or because they may be regulated post-transcriptionally to minimize the
258 time delay associated with response to a potentially lethal condition. Among the responses we
259 did observe, we found that under H₂O₂ stress, *E. faecalis* restricted the metabolism of
260 ethanolamine, and limited the metabolism of methylglyoxal, but increased its use of transport
261 systems to satisfy the specific demand for essential metal ions, such as iron. Concomitantly,
262 we observed decreased expression of some genes known to contribute to adaptation to other
263 stresses (*e.g.* *nhaC*, *cadA*). Future work will aim at determining whether and how (i)
264 metabolic changes in ethanolamine and methylglyoxal metabolism, and iron transport,
265 impact the adaptation of *E. faecalis* to these specific conditions; and (ii) how changes in

266 expression of genes clearly involved in the adaptation to other stresses affect cross adaptation.
267 It will also be of importance to determine the role of the 60 genes encoding hypothetical
268 proteins that were shown to be deregulated in these conditions.

269

270 **Acknowledgements**

271 X.Y. was the recipient of a doctoral fellowship from INRA (MICA Department) and the
272 Conseil Régional de Basse-Normandie, France. Microarray design, development and
273 production was supported by NIH grant AI083214 “Harvard-wide program on antibiotic
274 resistance.” The authors deeply thank Annick Blandin and Marie-Jeanne Pigny for expert
275 technical assistance, and Christelle Thibault from IGBMC, Strasbourg (France), for DNA
276 chip hybridization.

277

278 **References**

- 279 [1] Andrews SC, Robinson AK, Rodríguez-Quñones F (2003) Bacterial iron homeostasis.
280 FEMS Microbiol Rev 27:215–237.
- 281 [2] Ballal A, Basu B, Apte SK (2007) The Kdp-ATPase system and its regulation. J Biosci
282 32:559–568.
- 283 [3] Baskaran S, Prasanna Rajan D, Balasubramanian KA (1989) Formation of
284 methylglyoxal by bacteria isolated from human faeces. J Med Microbiol 28:211–215.
- 285 [4] Booth IR, Ferguson G, Miller S, Li C, Gunasekera B, Kinghorn S (2003) Bacterial
286 production of methylglyoxal: a survival strategy or death by misadventure? Biochem
287 Soc Trans 31:1406–1408.
- 288 [5] Bourgogne A, Hilsenbeck SG, Dunny GM, Murray BE (2006) Comparison of OG1RF
289 and an isogenic *fsrB* deletion mutant by transcriptional analysis: the Fsr system of
290 *Enterococcus faecalis* is more than the activator of gelatinase and serine protease. J
291 Bacteriol 188:2875–2884.
- 292 [6] Burke RM, Tempest DW (1990) Growth of *Bacillus stearothermophilus* on glycerol in
293 chemostat culture: expression of an unusual phenotype. J Gen Microbiol
294 136:1381–1385.
- 295 [7] Collier PD, Cromie DDO, Davies AP (1991) Mechanism of formation of
296 chloropropanols present in protein hydrolysates. J Am Oil Chem Soc 68:785–790.
- 297 [8] Cooper RA (1984) Metabolism of methylglyoxal in microorganisms. Annu Rev
298 Microbiol 38:49–68.
- 299 [9] Cornelis P, Wei Q, Andrews SC, Vinckx T (2011) Iron homeostasis and management of
300 oxidative stress response in bacteria. Metallomics 3:540–549.
- 301 [10] Epstein W (1986) Osmoregulation by potassium transport in *Escherichia coli*. FEMS
302 Microbiol Lett 39:73–78.
- 303 [11] Ferguson GP, Töttemeyer S, MacLean MJ, Booth IR (1998) Methylglyoxal production in
304 bacteria: suicide or survival? Arch Microbiol 170:209–218.
- 305 [12] Flahaut S, Laplace JM, Frère J, Auffray Y (1998) The oxidative stress response in
306 *Enterococcus faecalis*: relationship between H₂O₂ tolerance and H₂O₂ stress proteins.
307 Lett Appl Microbiol 26:259–264.
- 308 [13] Franceschini A, Szklarczyk D, Frankild S, Kuhn M, Simonovic M, Roth A, et al. (2013)
309 STRING v9.1: protein-protein interaction networks, with increased coverage and
310 integration. Nucleic Acids Res 41:D808–815.
- 311 [14] Frankenberg L, Brugna M, Hederstedt L (2002) *Enterococcus faecalis* Heme-Dependent
312 Catalase. J Bacteriol 184:6351–6356.

- 313 [15] Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, et al. (2004)
314 Bioconductor: open software development for computational biology and
315 bioinformatics. *Genome Biol* 5:R80.
- 316 [16] Giard JC, Riboulet E, Verneuil N, Sanguinetti M, Auffray Y, Hartke A (2006)
317 Characterization of Ers, a PrfA-like regulator of *Enterococcus faecalis*. *FEMS Immunol*
318 *Med Microbiol* 46:410–418.
- 319 [17] Giraffa G (2003) Functionality of *enterococci* in dairy products. *Int J Food Microbiol*
320 88:215–222.
- 321 [18] Hugas M, Garriga M, Aymerich MT (2003) Functionality of *enterococci* in meat
322 products. *Int J Food Microbiol* 88:223–233.
- 323 [19] Inoue Y, Kimura A (1995) Methylglyoxal and regulation of its metabolism in
324 microorganisms. *Adv Microb Physiol* 37:177–227.
- 325 [20] Ito M, Guffanti AA, Zemsy J, Ivey DM, Krulwich TA (1997) Role of the
326 *nhaC*-encoded Na⁺/H⁺ antiporter of alkaliphilic *Bacillus firmus* OF4. *J Bacteriol*
327 179:3851–3857.
- 328 [21] Kadner RJ, Murphy GP, Stephens CM (1992) Two mechanisms for growth inhibition by
329 elevated transport of sugar phosphates in *Escherichia coli*. *J Gen Microbiol*
330 138:2007–2014.
- 331 [22] Kang Y, Edwards LG, Thornalley PJ (1996) Effect of methylglyoxal on human
332 leukaemia 60 cell growth: modification of DNA G1 growth arrest and induction of
333 apoptosis. *Leuk Res* 20:397–405.
- 334 [23] Klebanoff SJ (1980) Oxygen metabolism and the toxic properties of phagocytes. *Ann*
335 *Intern Med* 93:480–489.
- 336 [24] Kofoid E, Rappleye C, Stojiljkovic I, Roth J (1999) The 17-gene ethanolamine (*eut*)
337 operon of *Salmonella typhimurium* encodes five homologues of carboxysome shell
338 proteins. *J Bacteriol* 181:5317–5329.
- 339 [25] Laplace JM, Hartke A, Giard JC, Auffray Y (2000) Cloning, characterization and
340 expression of an *Enterococcus faecalis* gene responsive to heavy metals. *Appl Microbiol*
341 *Biotechnol* 53:685–689.
- 342 [26] Lawhon SD, Frye JG, Suyemoto M, Porwollik S, McClelland M, Altier C (2003) Global
343 regulation by CsrA in *Salmonella typhimurium*. *Mol Microbiol* 48:1633–1645.
- 344 [27] Lebrun M, Audurier A, Cossart P (1994) Plasmid-borne cadmium resistance genes in
345 *Listeria monocytogenes* are present on Tn5422, a novel transposon closely related to
346 Tn917. *J Bacteriol* 176:3049–3061.
- 347 [28] Lebrun M, Loulergue J, Chaslus-Dancla E, Audurier A (1992) Plasmids in *Listeria*
348 *monocytogenes* in relation to cadmium resistance. *Appl Environ Microbiol*
349 58:3183–3186.

- 350 [29] López G, Latorre M, Reyes-Jara A, Cambiazo V, González M (2012) Transcriptomic
351 response of *Enterococcus faecalis* to iron excess. *Biometals* 25:737–747.
- 352 [30] Mercenier A, Pavan S, Pot B (2003) Probiotics as biotherapeutic agents: present
353 knowledge and future prospects. *Curr Pharm Des* 9:175–191.
- 354 [31] Misra K, Banerjee AB, Ray S, Ray M (1995) Glyoxalase III from *Escherichia coli*: a
355 single novel enzyme for the conversion of methylglyoxal into D-lactate without reduced
356 glutathione. *Biochem J* 305:999–1003.
- 357 [32] Misra K, Banerjee AB, Ray S, Ray M (1996) Reduction of methylglyoxal in *Escherichia*
358 *coli* K12 by an aldehyde reductase and alcohol dehydrogenase. *Mol Cell Biochem*
359 156:117–124.
- 360 [33] Mostertz J, Scharf C, Hecker M, Homuth G (2004) Transcriptome and proteome analysis
361 of *Bacillus subtilis* gene expression in response to superoxide and peroxide stress.
362 *Microbiology* 150:497–512.
- 363 [34] Nathan CF (1983) Mechanisms of macrophage antimicrobial activity. *Trans R Soc Trop*
364 *Med Hyg* 77:620–630.
- 365 [35] Noskin GA, Till M, Patterson BK, Clarke JT, Warren JR (1991) High-level gentamicin
366 resistance in *Enterococcus faecalis* bacteremia. *J Infect Dis* 164:1212–1215.
- 367 [36] Nucifora G, Chu L, Misra TK, Silver S (1989) Cadmium resistance from *Staphylococcus*
368 *aureus* plasmid pI258 *cadA* gene results from a cadmium-efflux ATPase. *Proc Natl Acad*
369 *Sci USA* 86:3544–3548.
- 370 [37] Okado A, Kawasaki Y, Hasuike Y, Takahashi M, Teshima T, Fujii J, et al. (1996)
371 Induction of apoptotic cell death by methylglyoxal and 3-deoxyglucosone in
372 macrophage-derived cell lines. *Biochem Biophys Res Commun* 225:219–224.
- 373 [38] del Papa MF, Perego M (2008) Ethanolamine activates a sensor histidine kinase
374 regulating its utilization in *Enterococcus faecalis*. *J Bacteriol* 190:7147–7156.
- 375 [39] Paulsen IT, Banerjee L, Myers GSA, Nelson KE, Seshadri R, Read TD, et al. (2003) Role
376 of mobile DNA in the evolution of vancomycin-resistant *Enterococcus faecalis*. *Science*
377 299:2071–2074.
- 378 [40] Pichereau V, Hartke A, Auffray Y (2000) Starvation and osmotic stress induced
379 multiresistances. Influence of extracellular compounds. *Int J Food Microbiol* 55:19–25.
- 380 [41] Poyart C, Quesnes G, Trieu-Cuot P (2000) Sequencing the gene encoding
381 manganese-dependent superoxide dismutase for rapid species identification of
382 *enterococci*. *J Clin Microbiol* 38:415–418.
- 383 [42] Prince WS, Villarejo MR (1990) Osmotic control of *proU* transcription is mediated
384 through direct action of potassium glutamate on the transcription complex. *J Biol Chem*
385 265:17673–17679.

- 386 [43] Rincé A, Giard JC, Pichereau V, Flahaut S, Auffray Y (2001) Identification and
387 characterization of gsp65, an organic hydroperoxide resistance (ohr) gene encoding a
388 general stress protein in *Enterococcus faecalis*. J Bacteriol 183:1482–1488.
- 389 [44] Ross RP, Claiborne A (1991) Cloning, sequence and overexpression of NADH
390 peroxidase from *Streptococcus faecalis* 10C1. Structural relationship with the
391 flavoprotein disulfide reductases. J Mol Biol 221:857–871.
- 392 [45] Sahm DF, Kissinger J, Gilmore MS, Murray PR, Mulder R, Solliday J, et al. (1989) *In*
393 *vitro* susceptibility studies of vancomycin-resistant *Enterococcus faecalis*. Antimicrob
394 Agents Chemother 33:1588–1591.
- 395 [46] Schaberg DR, Culver DH, Gaynes RP (1991) Major trends in the microbial etiology of
396 nosocomial infection. Am J Med 91:72S–75S.
- 397 [47] Skaar EP (2010) The battle for iron between bacterial pathogens and their vertebrate
398 hosts. PLoS Pathog 6:e1000949.
- 399 [48] Stohs SJ, Bagchi D (1995) Oxidative mechanisms in the toxicity of metal ions. Free
400 Radic Biol Med 18:321–336.
- 401 [49] Tempest DW, Neijssel OM (1984) The status of YATP and maintenance energy as
402 biologically interpretable phenomena. Annu Rev Microbiol 38:459–486.
- 403 [50] Thomas EL, Lehrer RI, Rest RF (1988) Human neutrophil antimicrobial activity. Rev
404 Infect Dis 10 Suppl:S450–456.
- 405 [51] Touati D (2000) Iron and oxidative stress in bacteria. Arch Biochem Biophys 373:1–6.
- 406 [52] Vallee BL, Ulmer DD (1972) Biochemical effects of mercury, cadmium, and lead. Annu
407 Rev Biochem 41:91–128.
- 408 [53] Vebø HC, Snipen L, Nes IF, Brede DA (2009) The transcriptome of the nosocomial
409 pathogen *Enterococcus faecalis* V583 reveals adaptive responses to growth in blood.
410 PLoS One 4:e7660.
- 411 [54] Verneuil N, Le Breton Y, Hartke A, Auffray Y, Giard JC (2004) Identification of a new
412 oxidative stress transcriptional regulator in *Enterococcus faecalis*. Lait 84:69–76.
- 413 [55] Verneuil N, Rincé A, Sanguinetti M, Auffray Y, Hartke A, Giard JC (2005) Implication
414 of *hypR* in the virulence and oxidative stress response of *Enterococcus faecalis*. FEMS
415 Microbiol Lett 252:137–141.
- 416 [56] Verneuil N, Rincé A, Sanguinetti M, Posteraro B, Fadda G, Auffray Y, et al. (2005)
417 Contribution of a PerR-like regulator to the oxidative-stress response and virulence of
418 *Enterococcus faecalis*. Microbiology 151:3997–4004.
- 419 [57] Verneuil N, Sanguinetti M, Le Breton Y, Posteraro B, Fadda G, Auffray Y, et al. (2004)
420 Effects of the *Enterococcus faecalis hypR* gene encoding a new transcriptional regulator

- 421 on oxidative stress response and intracellular survival within macrophages. *Infect*
422 *Immun* 72:4424–4431.
- 423 [58] Wu L, Juurlink BHJ (2002) Increased methylglyoxal and oxidative stress in hypertensive
424 rat vascular smooth muscle cells. *Hypertension* 39:809–814.
- 425 [59] Yan X, Zhao C, Budin-Verneuil A, Hartke A, Rincé A, Gilmore MS, Auffray Y,
426 Pichereau V (2009) The (p)ppGpp synthetase RelA contributes to stress adaptation and
427 virulence in *Enterococcus faecalis* V583. *Microbiology* 155:3226–3237.
- 428 [60] Yan X, Budin-Verneuil A, Auffray Y, Pichereau V (2014) Proteome phenotyping of
429 Δ relA mutants in *Enterococcus faecalis* V583. *Can J Microbiol* 60:525–531.
- 430 [61] Zhao C, Hartke A, La Sorda M, Posteraro B, Laplace JM, Auffray Y, et al. (2010) Role
431 of methionine sulfoxide reductases A and B of *Enterococcus faecalis* in oxidative stress
432 and virulence. *Infect Immun* 78:3889–3897.
- 433

434 **Legends to figures**

435

436 Fig. 1. Growth of V583 ery^S with shaking in GM17 medium or supplemented with 1.5mM,
437 1.75mM and 2mM H₂O₂. The arrows indicate the time at which RNA was extracted.

438

439 Fig. 2. Functional classification of genes with statistically significant increases (■) and
440 decreases (□) in mRNA level upon 4 h exposure to hydrogen peroxide (total of 213 genes).
441 The total number of genes associated with increased/decreased mRNA abundance as the
442 result of H₂O₂ exposure in each functional class is shown.

443

444 Fig. 3. Network analysis of genes shown to be associated with significant transcript changes
445 (built using STRING v9.1 [13]).

446

447 Table 1. Oligonucleotide primers used for real-time quantitative PCR in this study.

0188L: TCGCAGCTTGTGGTAATACG	0188R: ACTCGCAATAATCCGTTTGG
0191L: GGAACACCGCAAGAAGTGAT	0191R: GCCGCTGTGTCTGACTATGA
0475L: TCGAATTGCGCTGACAGATA	0475R: TAGCGCGTTCTCCAACCTTT
0567L: CGCTATGCACTGAATGTGCT	0567R: ACCTCCCGGTAAAAAGTGCT
0570L: GCTGGTGTTGGGAAAACCTA	0570R: GGACGATCATGAGGTTTCGAT
0575L: AGGCTTTGGAAAAGGTCGAT	0575R: AAATCCGCATTTTGAGCAAG
0630L: AGTCAACCACGTGGGGATTA	0630R: TTCCACCAATTTTCATCACGA
0666L: TCTTGAGCAAGCGGGTATCT	0666R: AGCGCCATTTGCGTATTATC
1120L: TTAAAGTGATGGGCGTTTCC	1120R: TGCGAAGGATAAGCAGAAGC
1617L: ACCCGAAATCGACGTAACAG	1617R: GGCGCGGACTTTCTTCTAA
1620L: TGAAGGTAAGCCGGTATTGG	1620R: GTCCACGTTTTCTGAATCG
1621L: CAACCTGGACAGTTCGTTTGT	1621R: TGTGATTGCTACGTTTCAGG
1624L: AAATTGGCACCGATTATTGC	1624R: ATTGTATGACCCGCACCTTC
1627L: CACGTCCTGAAACGACAGAA	1627R: CAGGATCTACGGCATTTTTCA
1629L: ATCCATCGCTGGTTCAGAAA	1629R: AGCTGCTTGACCACTGTGAA
1633L: TTAGGCGCACTGGGTTATTT	1633R: TAACAGCTGCGTTTGTGTTGC
1635L: CAGGGATGGCGTTTAATGTT	1635R: TTGCATTCATTCGTCCATGT
1638L: ATAAAACACAGGCGGTGGAA	1638R: GCCGCTGTGACGTTTAATG
1669L: GCCCAAGTCGTTTTGGTAGA	1669R: TGATTTACGATCCGCTAGGC
3085L: AACCGATAGTGCAAGGTTGG	3085R: CCGCTTAGCGCAAAAATAAA

448

449

450
451

Table 2. Genes of *Enterococcus faecalis* V583 ery^S associated with increased transcript abundance as the result of exposure to H₂O₂.

Locus	Descriptions (Gene symbol)	M*	SD
Biosynthesis and metabolism			
EF0693	1-phosphofructokinase (<i>fruK-1</i>)	1.1	0.3
EF1222	adenine deaminase (<i>ade</i>)	1.4	0.2
EF1511	mandelate racemase/muconate lactonizing enzyme family protein	1.1	0.4
EF2999	allantoinase, putative	1.1	0.1
Transport and binding protein			
EF0188	iron compound ABC transporter, substrate-binding protein	2.0	0.6
EF0191	ferrichrome ABC transporter, ATP-binding protein	1.1	0.3
EF0192	ferrichrome ABC transporter, permease protein	1.0	0.1
EF0193	ferrichrome ABC transporter, permease protein (<i>fhuG</i>)	1.0	0.3
EF0475	ferrous iron transport protein A (<i>feoA</i>)	1.3	0.0
EF0476	ferrous iron transport protein B (<i>feoB</i>)	1.6	0.1
EF0567	potassium-transporting ATPase, subunit A (<i>kdpA</i>)	1.2	0.4
EF0568	potassium-transporting ATPase, subunit B (<i>kdpB</i>)	1.2	0.4
EF0569	potassium-transporting ATPase, subunit C (<i>kdpC</i>)	1.2	0.1
EF0575	cationic ABC transporter, ATP-binding protein	1.0	0.2
EF0576	cation ABC transporter, permease protein	1.1	0.1
EF0694	PTS system, fructose-specific family, IIBC components	1.0	0.6
EF1053	ABC transporter, ATP-binding protein	1.0	0.5
EF1117	amino acid ABC transporter, permease protein	2.2	0.4
EF1118	amino acid ABC transporter, permease protein	2.3	0.2
EF1119	amino acid ABC transporter, amino acid-binding protein	2.0	0.1
EF1120	amino acid ABC transporter, ATP-binding protein	1.8	0.1
EF1220	spermidine/putrescine ABC transporter, ATP-binding protein	1.0	0.2
EF3082	iron compound ABC transporter, substrate-binding protein	3.1	0.8
EF3083	iron compound ABC transporter, ATP-binding protein	3.2	0.6
EF3084	iron compound ABC transporter, permease protein	3.2	0.7
EF3085	iron compound ABC transporter, permease protein	3.3	0.6
Regulation and signal			
EF0570	sensor histidine kinase (<i>kdpD</i>)	1.0	0.1
EF0578	helix-turn-helix protein, iron-dependent repressor family	1.3	0.1
Cell envelope and cell division			
EF0887	glycosyl transferase, group 2 family protein	1.5	0.1
EF3314	cell wall surface anchor family protein	1.5	0.1
Stress and Virulence			
EF2068	multidrug resistance protein, putative	1.2	0.1
Unknown			
EF0477	hypothetical protein	1.7	0.8
EF0886	hypothetical protein	1.4	0.2
EF0888	conserved hypothetical protein	1.0	0.1
EF0889	conserved hypothetical protein	1.2	0.4
EF1223	chlorohydrolase family protein	1.2	0.6
EF1512	conserved hypothetical protein	1.1	0.3
EF1808	agaS protein	1.1	0.4
EF2067	conserved hypothetical protein TIGR00481	1.1	0.0

*M is the mean Log₂(fold ratio); SD is the standard deviation.

452
453

Table 3. Genes of *Enterococcus faecalis* V583 ery^S associated with transcript reductions as the result of exposure to H₂O₂.

Locus	Descriptions (Gene symbol)	M*	SD
Biosynthesis and metabolism			
EF0098	L-serine dehydratase, iron-sulfur-dependent, beta subunit (<i>sdhB-1</i>)	-2.4	0.2
EF0099	L-serine dehydratase, iron-sulfur-dependent, alpha subunit (<i>sdhA-1</i>)	-2.7	0.6
EF0630	glyoxalase family protein	-1.0	0.1
EF0867	glyoxalase family protein	-1.2	0.1
EF1358	glycerol dehydrogenase, putative	-1.6	0.5
EF1617	conserved hypothetical protein (<i>EutQ</i>)	-2.6	0.5
EF1618	ethanolamine utilization protein (<i>EutH</i>)	-2.7	0.6
EF1619	carbon dioxide concentrating mechanism protein CcmL, putative (<i>EutN</i>)	-2.8	0.3
EF1620	hypothetical protein (<i>EutX</i>)	-2.9	0.5
EF1621	conserved hypothetical protein (<i>EutY</i>)	-2.8	0.5
EF1622	conserved domain protein (<i>EutZ</i>)	-3.0	0.3
EF1623	microcompartment protein (<i>EutK</i>)	-2.6	0.4
EF1624	aldehyde dehydrogenase, putative (<i>EutE</i>)	-3.1	0.1
EF1625	microcompartment protein family (<i>EutM</i>)	-3.0	0.0
EF1626	ethanolamine utilization protein (<i>EutL</i>)	-3.1	0.1
EF1627	ethanolamine ammonia-lyase small subunit (<i>EutC</i>)	-3.3	0.1
EF1629	ethanolamine ammonia-lyase large subunit (<i>EutB</i>)	-3.4	0.1
EF1632	sensor histidine kinase (<i>HK</i>)	-1.1	0.1
EF1633	response regulator (<i>RR</i>)	-1.1	0.1
EF1634	propanediol utilization protein PduU (<i>EutS</i>)	-1.1	0.1
EF1635	propanol dehydrogenase PduQ, putative (<i>EutG</i>)	-2.1	0.0
EF1637	ATP:cob(I)alamin adenosyltransferase, putative (<i>EutT</i>)	-1.7	0.0
EF1638	propanediol utilization protein PduV (<i>eutP</i>)	-1.3	0.1
EF1669	glyoxylase family protein	-1.0	0.1
Protein biosynthesis and fate			
EF0100	seryl-tRNA synthetase (<i>serS-1</i>)	-3.4	0.1
EF0650	lipoate-protein ligase A (<i>lplA-1</i>)	-1.4	0.1
DNA metabolism			
EF2114	adenine methyltransferase, putative	-1.7	0.1
Transport and binding protein			
EF0636	Na ⁺ H ⁺ antiporter (<i>nhaC-2</i>)	-1.1	0.2
EF0986	cation transporter	-1.0	0.1
EF2623	cadmium-translocating P-type ATPase (<i>cadA</i>)	-2.4	0.4
Regulation and signal			
EF0097	regulatory protein pfoR, putative	-2.0	0.3
EF0110	transcriptional regulator, ArsR family	-1.2	0.1
EF0143	transcriptional regulator, CroCI family	-1.2	0.5
EF0923	transcriptional regulator, LysR family	-1.2	0.2
EF1668	transcriptional regulator, MarR family	-1.4	0.1
EF2141	transcriptional regulator, CroCI family	-1.3	0.1
Cell envelope and cell division			
EF0139	FtsKSpolIIIIE family protein	-1.1	0.0
EF0153	cell wall surface anchor family protein	-1.1	0.2

Stress and virulence		
EF0149	aggregation substance, putative	-1.4 0.2
Mobile and extrachromosomal element functions		
EF0158	conjugal transfer protein, putative	-1.3 0.4
EF2096	tail protein	-1.5 0.1
Unknown		
EF0083	hypothetical protein	-1.1 0.6
EF0109	ThiJPfpl family protein	-1.1 0.1
EF0131	conserved domain protein	-1.5 0.2
EF0132	hypothetical protein	-1.4 0.2
EF0134	hypothetical protein	-1.1 0.1
EF0135	conserved hypothetical protein	-1.0 0.1
EF0136	hypothetical protein	-1.1 0.1
EF0138	conserved domain protein	-1.1 0.0
EF0142	conserved hypothetical protein	-1.0 0.3
EF0144	conserved domain protein	-1.2 0.3
EF0151	hypothetical protein	-1.2 0.4
EF0152	hypothetical protein	-1.0 0.6
EF0154	conserved hypothetical protein	-1.7 0.1
EF0155	conserved hypothetical protein	-1.6 0.4
EF0415	conserved hypothetical protein	-1.4 0.4
EF0519	hypothetical protein	-1.0 0.1
EF0971	conserved hypothetical protein	-1.1 0.1
EF1327	BadFBadGBcrABcrD ATPase family protein	-1.4 0.5
EF1359	conserved hypothetical protein	-1.3 0.3
EF1360	dihydroxyacetone kinase family protein	-1.2 0.4
EF1361	dihydroxyacetone kinase family protein	-1.4 0.2
EF1362	conserved domain protein	-1.1 0.1
EF2014	coenzyme F420 hydrogenase domain protein	-1.0 0.1
EF2065	conserved hypothetical protein	-1.0 0.1
EF2094	hypothetical protein	-1.1 0.1
EF2095	hypothetical protein	-1.1 0.3
EF2097	hypothetical protein	-1.5 0.1
EF2098	hypothetical protein	-1.4 0.2
EF2099	hypothetical protein	-1.6 0.1
EF2100	hypothetical protein	-1.5 0.1
EF2101	hypothetical protein	-1.6 0.2
EF2102	hypothetical protein	-1.7 0.4
EF2103	hypothetical protein	-1.6 0.3
EF2104	hypothetical protein	-1.7 0.1
EF2105	hypothetical protein	-1.7 0.1
EF2106	conserved domain protein	-1.6 0.1
EF2107	hypothetical protein	-1.7 0.1
EF2108	hypothetical protein	-1.6 0.3
EF2109	conserved domain protein	-1.8 0.0
EF2110	hypothetical protein	-1.6 0.0
EF2111	hypothetical protein	-1.7 0.1
EF2112	hypothetical protein	-1.7 0.1
EF2113	conserved hypothetical protein	-1.6 0.3
EF2120	conserved hypothetical protein	-1.0 0.6
EF2125	hypothetical protein	-1.0 0.3
EF2127	hypothetical protein	-1.0 0.2

EF2133	hypothetical protein	-1.0	0.3
EF2134	hypothetical protein	-1.0	0.6
EF2137	hypothetical protein	-1.2	0.3
EF2140	conserved hypothetical protein	-1.2	0.2
EF2606	conserved hypothetical protein	-1.0	0.1
EF2792	conserved hypothetical protein	-1.0	0.0

456 *M is the mean Log₂(fold ratio); SD is the standard deviation.

457

458

459 Table 4. Comparison of gene expression analyzed by real-time quantitative PCR and
 460 Microarray.

<i>Locus</i>	<i>(Gene symbol)</i> Function	$H_2O_2/GM17$	
		RT qPCR	Microarray
<i>ef0188</i>	iron compound ABC transporter, substrate-binding protein	9.24±1.95	3.94±1.20
<i>ef0191</i>	ferrichrome ABC transporter, ATP-binding protein	3.33±0.70	2.20±0.44
<i>ef0475</i>	(feoA) ferrous iron transport protein A	3.16±0.99	2.48±0.03
<i>ef0567</i>	(kdpA) potassium-transporting ATPase, subunit A	2.72±1.27	2.27±0.34
<i>ef0570</i>	(kdpD) sensor histidine kinase KdpD	1.67±0.38	1.99±0.12
<i>ef0575</i>	cationic ABC transporter, ATP-binding protein	2.37±0.77	1.92±0.25
<i>ef1120</i>	amino acid ABC transporter, ATP-binding protein	6.44±0.25	3.52±0.03
<i>ef3085</i>	iron compound ABC transporter, permease protein	18.1±7.18	11.2±3.81
<i>ef0630</i>	glyoxalase family protein	-3.92±0.91	-2.00±0.14
<i>ef0867</i>	glyoxalase family protein	-2.67±0.40	-2.33±0.12
<i>ef1617</i>	(eutQ) conserved hypothetical protein	-8.16±0.48	-6.10±1.41
<i>ef1620</i>	(eutX) hypothetical protein	-8.62±0.50	-7.50±1.41
<i>ef1621</i>	(eutY) conserved hypothetical protein	-3.41±0.95	-7.00±1.41
<i>ef1624</i>	(eutE) aldehyde dehydrogenase, putative	-9.18±0.60	-8.60±1.07
<i>ef1627</i>	(eutC) ethanolamine ammonia-lyase small subunit	-10.1±0.46	-9.80±1.07
<i>ef1629</i>	(eutB) ethanolamine ammonia-lyase large subunit	-7.76±0.45	-10.6±1.07
<i>ef1633</i>	(RR17) response regulator	-4.13±0.24	-2.10±1.07
<i>ef1635</i>	(eutG) propanol dehydrogenase PduQ, putative	-11.7±0.85	-4.30±1.07
<i>ef1638</i>	(eutP) propanediol utilization protein PduV	-2.82±0.26	-2.50±1.07
<i>ef1669</i>	glyoxylase family protein	-2.90±1.15	-2.04±0.04

461 Values are given as means of the fold ratios ± standard deviation (SD).

462

463 Table 5. Expression of all the genes encoding glyoxalase family protein analyzed by
 464 Microarray.

<i>Locus</i>	Function	Level of gene expression				Ratio
		GM17	H ₂ O ₂	GM17	H ₂ O ₂	
<i>ef0358</i>	glyoxalase family protein	330.7	164.6	408.5	224.2	-1.92±0.10
<i>ef0939</i>	methylglyoxal synthase	1627.9	961	1806.1	909.7	-1.84±0.15
<i>ef0630</i>	glyoxalase family protein	1152.1	541.5	945.8	508.4	-2.00±0.14
<i>ef0656</i>	glyoxalase family protein	138.8	112.8	236.1	178.6	-1.28±0.04
<i>ef0666</i>	glyoxalase family protein	2717	1834	2571.2	1811.8	-1.45±0.03
<i>ef0745</i>	glyoxalase family protein	1003.8	974.3	1216.9	1119.1	-1.06±0.03
<i>ef0867</i>	glyoxalase family protein	253.7	114.8	292.7	119.8	-2.33±0.12
<i>ef1669</i>	glyoxylase family protein	260.3	125.4	314.8	161	-2.04±0.04
<i>ef2591</i>	glyoxalase family protein	3065.4	3012.3	3317.9	2821.4	-1.10±0.08
<i>ef3092</i>	glyoxalase family protein	1302.2	1004.4	1238.1	937.8	-1.31±0.01

465 Values are given as means of the fold ratios ± standard deviation (SD).

466

467

468 Table 6. Expression of the genes known to be involved in enterococcal adaptation to oxidative
 469 stress.

<i>Locus</i>	Protein (<i>Gene</i>)	Expression of genes		Ratio H ₂ O ₂ /GM17
		GM17	GM17 +H ₂ O ₂	
Oxidative stress proteins				
<i>ef0453</i>	Organic hydroperoxide resistance protein (<i>ohr</i>)	3712.9	3449.6	-1.07
<i>ef0463</i>	Superoxide dismutase (<i>sodA</i>)	4697.8	4342.7	-1.08
<i>ef0606</i>	DNA-binding protein (<i>dps</i>)	384.5	370	-1.04
<i>ef1211</i>	NADH peroxidase (<i>npr</i>)	4228.5	3577.3	-1.18
<i>ef1338</i>	Thioredoxine reductase (<i>trx</i>)	3292.4	2649.6	-1.24
<i>ef1405</i>	Thioredoxine reductase (<i>trx</i>)	1787.7	1772.6	-1.01
<i>ef1586</i>	NADH oxidase (<i>nox</i>)	5842.3	6058.5	1.04
<i>ef1597</i>	Catalase (<i>katA</i>)	2734.8	2806.2	1.03
<i>ef1681</i>	Peptide methionine-S-sulfoxide reductase (<i>msrA</i>)	2144	1908.5	1.12
<i>ef2738</i>	Peroxiredoxine reductase (<i>ahpF</i>)	3974	3984.6	1.00
<i>ef2739</i>	Alkyl hydroperoxide reductase (<i>ahpC</i>)	4536.4	4583	1.01
<i>ef3164</i>	Peptide methionine-S-sulfoxide reductase (<i>msrB</i>)	2449	1478.1	-1.65
<i>ef3233</i>	DNA-binding protein (<i>dps</i>)	4553.2	4202.9	-1.08
<i>ef3270</i>	Glutathione reductase (<i>gor</i>)	2707.7	2879.4	1.06
Oxidative stress related transcriptional regulators				
<i>ef1525</i>	(<i>fur</i>)	2925.4	2397.8	-1.22
<i>ef1585</i>	(<i>perR</i>)	2741.6	2304.2	-1.19
<i>ef2063</i>	(<i>soxS</i>)	1847.9	1969.6	1.07
<i>ef2417</i>	(<i>zur</i>)	1017.1	804.2	-1.26
<i>ef2958</i>	(<i>hypR</i>)	245.1	256.9	1.05

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484




Fig. 1.




Fig. 2.

