



SPATIO-TEMPORAL POISSON PROCESSES FOR VISITS TO SMALL SETS

Françoise Pène, Benoît Saussol

► To cite this version:

Françoise Pène, Benoît Saussol. SPATIO-TEMPORAL POISSON PROCESSES FOR VISITS TO SMALL SETS. Israel Journal of Mathematics, 2020, 240 (2), pp.625-665. 10.1007/s11856-020-2074-0 . hal-01737173

HAL Id: hal-01737173

<https://hal.univ-brest.fr/hal-01737173>

Submitted on 19 Mar 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SPATIO-TEMPORAL POISSON PROCESSES FOR VISITS TO SMALL SETS

FRANÇOISE PÈNE AND BENOÎT SAUSSOL

ABSTRACT. For many measure preserving dynamical systems (Ω, T, m) the successive hitting times to a small set is well approximated by a Poisson process on the real line. In this work we define a new process obtained from recording not only the successive times n of visits to a set A , but also the position $T^n(x)$ in A of the orbit, in the limit where $m(A) \rightarrow 0$.

We obtain a convergence of this process, suitably normalized, to a Poisson point process in time and space under some decorrelation condition. We present several new applications to hyperbolic maps and SRB measures, including the case of a neighborhood of a periodic point, and some billiards such as Sinai billiards, Bunimovich stadium and diamond billiard.

CONTENTS

1. Introduction	2
2. Poisson process under a one-step decorrelation assumption	3
2.1. Results for discrete-time dynamical systems	4
2.2. Application to special flows	5
2.3. Proofs of the general theorems	6
3. A common framework suitable for systems modeled by Gibbs-Markov-Young towers	8
3.1. General result	8
3.2. Successive visits in a small neighbourhood of a generic point	10
4. Applications to billiard maps and flows	10
4.1. Bunimovich billiard	10
4.2. Sinai billiards	12
5. Successive visits in a small neighborhood of an hyperbolic periodic point	15
5.1. General results around a periodic hyperbolic orbit	15
5.2. SRB measure for Anosov maps	18
6. Billiard in a diamond	21
6.1. Notations, recalls and proof of Theorem 6.1	22
6.2. Short returns	23
Appendix A. More proofs	26
References	28

Date: March 18, 2018.

2000 Mathematics Subject Classification. Primary: 37B20.

1. INTRODUCTION

The study of Poincaré recurrence in dynamical systems such as occurrence of rare events, distribution of return time, hitting time and Poisson law has grown to an active field of research, in deep relation with extreme values of stochastic processes; (see [13] and references therein).

Let $(\Omega, \mathcal{F}, \mu, T)$ be a probability preserving dynamical system. For every $A \in \mathcal{F}$, we set τ_A as the first hitting time to A , i.e.

$$\tau_A(x) := \inf\{n \geq 1 : T^n x \in A\}.$$


In many systems the behavior of the successive visits of a typical orbit $(T^n x)_n$ to the sets A_ε , with $\mu(A_\varepsilon) \rightarrow 0+$ is often asymptotic, when suitably normalized, to a Poisson process. Such results were first obtained by Doeblin [6] for the Gauss map, Pitskel [17] considered the case of Markov chains. The most recent developments concern non uniformly hyperbolic dynamical systems, for example [19, 3, 8, 10, 11] just to mention a few of them.

An important issue of our work is that we take into account not only the times of successive visits to the set, but also the position of the successive visits in A_ε within each return. This study was first motivated by a question asked to us by D. Szász and I.P. Tóth for diamond billiards, that we address in Section 6. Beyond its own interest, Poisson limit theorems for such spatio-temporal processes have been recently use to prove convergence to Lévy stable processes in dynamical systems; See [20] and subsequent works such as [14]. Let us indicate that, at the same time and independently of the present work, analogous processes have been investigated in [7].

We thus consider these events in time and space


$$\{(n, T^n x) : n \geq 1, T^n x \in A_\varepsilon\} \subset [0, +\infty) \times \Omega,$$

that we will normalize both in time and space, as follows.



The successive visit times have order $1/\mu(A_\varepsilon)$, which gives the normalization in time. For the space, we use a family of normalization functions $H_\varepsilon : A_\varepsilon \rightarrow V$. A typical choice when Ω is Euclidean would be to take for A_ε an ε -ball and H_ε a zoom which sends A_ε to size one. Another choice of extremal processes flavor would be to consider A_ε as a rare event, and H_ε would be the strength of the event. We will then consider the family of point processes $(\mathcal{N}_\varepsilon)_\varepsilon$ on $[0, +\infty) \times V$ given by

$$\mathcal{N}_\varepsilon(x) := \sum_{n \geq 1 : T^n(x) \in A_\varepsilon} \delta_{(n\mu(A_\varepsilon), H_\varepsilon(T^n(x)))}. \quad (1)$$



For any measurable subset B of $[0, +\infty) \times V$,

$$\mathcal{N}_\varepsilon(x)(B) = \#\{n \geq 1 : T^n(x) \in A_\varepsilon, (n\mu(A_\varepsilon), H_\varepsilon(T^n(x))) \in B\}.$$

We will simply write $\mathcal{N}_\varepsilon(B)$ for the measurable function $\mathcal{N}_\varepsilon(B) : x \mapsto \mathcal{N}_\varepsilon(x)(B)$.

The main result of the paper provides general conditions under which the point process \mathcal{N}_ε is well estimated by a Poisson point process \mathcal{P}_ε . This virtually contains all spatial information given by the space coordinate H_ε , and the continuous mapping theorem could in principle be applied to recover many properties related to recurrences in A_ε . We then present several applications, with different maps, flows, and sets A_ε .

The structure of the paper is as follows:

In Section 2 we present a general result which gives a convergence to a spatio-temporal point process in a discrete time dynamical system, under some probabilistic one-step decorrelation condition. Then we provide a method to transfer this result to continuous time flows. All these general results are proved in Subsection 2.3.

In Section 3 we introduce a framework, adapted for systems modeled by a Young tower, under which one can apply the general results. In Subsection 3.2 we check these conditions in the case of hitting to balls in a Riemmanian manifold.

In Section 4 we present several applications of the common framework mentioned above to two different types of billiards: the Sinai billiards with finite horizon, the Bunimovich stadium billiards.

In Section 5 we study the case of balls centered around a periodic point in a uniformly hyperbolic system; as a byproduct we recover a compound Poisson distribution for the temporal process.

Section 6 consists in a fine study of visits in the successive visits in the vicinity of the corner in a diamond shaped billiard.

2. POISSON PROCESS UNDER A ONE-STEP DECORRELATION ASSUMPTION

2.1. Results for discrete-time dynamical systems. Let $(\Omega, \mathcal{F}, \mu, T)$ be a probability preserving dynamical system. Let $(A_\varepsilon)_\varepsilon$ be a family of measurable subsets of Ω with $\mu(A_\varepsilon) \rightarrow 0+$ as $\varepsilon \rightarrow 0$. Let V be a locally compact metric space endowed with its Borel σ -algebra \mathcal{V} . Let $(H_\varepsilon)_\varepsilon$ be a family of measurable functions $H_\varepsilon : A_\varepsilon \rightarrow V$. We set $E := [0, +\infty) \times V$ and we endow it with its Borel σ -algebra $\mathcal{E} = \mathcal{B}([0, +\infty)) \otimes \mathcal{V}$. We also consider the family of measures $(m_\varepsilon)_\varepsilon$ on (V, \mathcal{V}) defined by

$$m_\varepsilon := \mu(H_\varepsilon^{-1}(\cdot)|A_\varepsilon) \quad (2)$$

and \mathcal{W} a family stable by finite unions and intersections of relatively compact open subsets of V , that generates the σ -algebra \mathcal{V} . Let λ be the Lebesgue measure on $[0, \infty)$.

We will approximate the point process defined by (1) by a Poisson point process on E . Given η a σ -finite measure on (E, \mathcal{E}) , recall that a process \mathcal{N} is a Poisson point process on E of intensity η if

- (i) \mathcal{N} is a point process (i.e. $\mathcal{N} = \sum_i \delta_{x_i}$ with x_i E -valued random variables),
- (ii) For every pairwise disjoint Borel sets $B_1, \dots, B_n \subset E$, the random variables $\mathcal{N}(B_1), \dots, \mathcal{N}(B_n)$ are independent Poisson random variables with respective parameters $\eta(B_1), \dots, \eta(B_n)$.

Let $M_p(E)$ be the space of all point measures defined on E , endowed with the topology of vague convergence ; it is metrizable as a complete separable metric space. A family of point processes $(\mathcal{N}_\varepsilon)_\varepsilon$ converges in distribution to \mathcal{N} if for any bounded continuous function $f : M_p(E) \rightarrow \mathbb{R}$ the following convergence holds true

$$\mathbb{E}(f(\mathcal{N}_\varepsilon)) \rightarrow \mathbb{E}(f(\mathcal{N})), \quad \text{as } \varepsilon \rightarrow 0. \quad (3)$$

For a collection \mathcal{A} of measurable subsets of Ω , we define the following quantity:

$$\Delta(\mathcal{A}) := \sup_{A \in \mathcal{A}, B \in \sigma(\cup_{n=1}^\infty T^{-n}\mathcal{A})} |\mu(A \cap B) - \mu(A)\mu(B)|. \quad (4)$$

Our main general result is the following one.

Theorem 2.1. *We assume that*

- (i) *for any finite subset \mathcal{W}_0 of \mathcal{W} we have $\Delta(H_\varepsilon^{-1}\mathcal{W}_0) = o(\mu(A_\varepsilon))$,*
- (ii) *there exists a measure m on (V, \mathcal{V}) such that for every $F \in \mathcal{W}$, $m(\partial F) = 0$ and $\lim_{\varepsilon \rightarrow 0} \mu(H_\varepsilon^{-1}(F)|A_\varepsilon)$ converges to $m(F)$.*

Then the family of point processes $(\mathcal{N}_\varepsilon)_\varepsilon$ converges strongly¹ in distribution to a Poisson process \mathcal{P} of intensity $\lambda \times m$.

In particular, for every relatively compact open $B \subset E$ such that $(\lambda \times m)(\partial B) = 0$, $(\mathcal{N}_\varepsilon(B))_\varepsilon$ converges in distribution to a Poisson random variable with parameter $(\lambda \times m)(B)$.

We emphasize that Theorem 2.1 remains valid when ε is restricted to a subsequence $\varepsilon_k \rightarrow 0$, in the assumptions and the conclusion.

¹i.e. with respect to any probability measure absolutely continuous w.r.t. μ

Condition (ii) is equivalent to the fact that the family of measures $(m_\varepsilon)_\varepsilon$ converges vaguely² to m . This is sometimes too strong, especially when the m_ε are not absolutely continuous. Nevertheless, without this hypothesis, our point process \mathcal{N}_ε may remain well approximated by a Poisson process \mathcal{P}_ε with varying intensities $\lambda \times m_\varepsilon$, in a sense that can be made precise.

Theorem 2.2. *We assume that for any vague limit point m of $(m_\varepsilon)_\varepsilon$ and for any sequence $\mathcal{E} = (\varepsilon_k)_k$ converging to 0 achieving the above limit, there exists a family $\mathcal{W}_\mathcal{E}$ of relatively compact open subsets of V , stable by finite unions and intersections, that generates the σ -algebra \mathcal{V} and that*

- (i) *for any finite subset \mathcal{W}_0 of $\mathcal{W}_\mathcal{E}$ we have $\Delta(H_{\varepsilon_k}^{-1}\mathcal{W}_0) = o(\mu(A_{\varepsilon_k}))$,*
- (ii) *for every $F \in \mathcal{W}_\mathcal{E}$, $m(\partial F) = 0$.*

Then the family of point processes $(\mathcal{N}_\varepsilon)_\varepsilon$ is approximated strongly in distribution by a family of Poisson processes $(\mathcal{P}_\varepsilon)_\varepsilon$ of intensities $\lambda \times m_\varepsilon$, in the sense that for any $\nu \ll \mu$ and for every continuous and bounded $f: M_p(E) \rightarrow \mathbb{R}$

$$\mathbb{E}_\nu(f(\mathcal{N}_\varepsilon)) - \mathbb{E}(f(\mathcal{P}_\varepsilon)) \rightarrow 0. \quad (5)$$

Proof of Theorem 2.2. Suppose that (5) does not hold for some f . Then there exists $\vartheta > 0$ and a sequence $\varepsilon_k \rightarrow 0$ such that for all k ,

$$|\mathbb{E}_\nu(f(\mathcal{N}_{\varepsilon_k})) - \mathbb{E}(f(\mathcal{P}_{\varepsilon_k}))| > \vartheta. \quad (6)$$

Up to taking a subsequence if necessary, we may assume that $(m_{\varepsilon_k})_k$ converges to some m . Applying Theorem 2.1 with the sequence $(\varepsilon_k)_k$ we get that $(\mathcal{N}_{\varepsilon_k})_k$ converges to a Poisson point process of intensity $\lambda \times m$, and $(\mathcal{P}_{\varepsilon_k})_k$ as well, which contradicts (6). \square

This proof shows that the possible non convergence of the measures m_ε is not a serious problem. **In the rest of the paper, we will assume without loss of generality - to simplify the exposition of our general results - that the intensity measures involved in the results converge to a unique limit.** Clearly, if it is not the case, one can always reduce the problem to that case by passing through a subsequence.

2.2. Application to special flows. In this section we show how to pass from the discrete time setting to the continuous time one. Given $(\Omega, \mathcal{F}, \mu, T)$ and an integrable function $\tau: \Omega \rightarrow (0, +\infty)$, the special flow over $(\Omega, \mathcal{F}, \mu, T)$ with roof function $\tau: \Omega \rightarrow (0, +\infty)$ is the flow $(\mathcal{M}, \mathcal{T}, \nu, (Y_t)_t)$ defined as follows:

$$\mathcal{M} := \{(x, t) \in \Omega \times [0, +\infty) : t < \tau(x)\},$$

\mathcal{T} is the trace in \mathcal{M} of the product σ -algebra $\mathcal{F} \otimes \mathcal{B}([0, +\infty))$

$$\nu := \left(\frac{\mu \times \lambda}{\int_\Omega \tau d\mu} \right)_{|\mathcal{T}},$$

²This means that for any continuous test function φ with compact support the integrals $m_\varepsilon(\varphi)$ converge to $m(\varphi)$. In particular m may not be a probability because of a loss of mass at infinity.

where λ is the Lebesgue measure on $[0, +\infty)$ and $Y_s(x, t) = (x, t + s)$ with the identification $(x, \tau(x)) \equiv (T(x), 0)$, i.e.

$$Y_s(x, t) = \left(T^{n_s(x, t)}, t + s - \sum_{k=0}^{n_s(x, t)-1} \tau \circ T^k(x) \right),$$

with $n_s(x, t) := \sup\{n : \sum_{k=0}^{n-1} \tau \circ T^k(x) \leq t + s\}$ the number of visits of the orbit $(Y_u(x, t))_{u \in (0, s)}$ to $\Omega \times \{0\}$ before time t . We will write

$$\forall x \in \Omega, \quad \forall n \in \mathbb{N}, \quad S_n \tau(x) := \sum_{k=0}^{n-1} \tau \circ T^k(x) \quad \text{and} \quad \bar{\tau} := \int_{\Omega} \tau(x) d\mu(x).$$

We define also the canonical projection $\Pi : \mathcal{M} \rightarrow \Omega$ by $\Pi(x, t) = x$.

Let $(\mathcal{A}_\varepsilon)_\varepsilon$ be a sequence of subsets of \mathcal{M} . We are interested in a process that records the times where at least one entrance into \mathcal{A}_ε occurs between two consecutive returns to the base. This only depends on the projection $A_\varepsilon := \Pi \mathcal{A}_\varepsilon$. Let V be a locally compact metric space endowed with its Borel σ -algebra \mathcal{V} . Let $(H_\varepsilon)_\varepsilon$ be a family of measurable functions from A_ε to V . This leads us to the definition

$$\mathfrak{N}_\varepsilon(y) := \sum_{t > 0 : Y_t(y) \in A_\varepsilon \times \{0\}} \delta_{(t\mu(A_\varepsilon)/\bar{\tau}, H_\varepsilon(\Pi Y_t(y)))}.$$

Theorem 2.3. *Assume that $(\mathcal{N}_\varepsilon)_\varepsilon$, defined on (Ω, μ, T) by (1) with $A_\varepsilon := \Pi \mathcal{A}_\varepsilon$ and H_ε given above, converges in distribution, with respect to some probability measure $\tilde{\mu} \ll \mu$, to a Poisson point process of intensity $\lambda \times m$, where m is some measure on (V, \mathcal{V}) .*

Then the family of point processes $(\mathfrak{N}_\varepsilon)_\varepsilon$ converges strongly in distribution to a Poisson process \mathcal{P} of intensity $\lambda \times m$.

2.3. Proofs of the general theorems. In this section we prove Theorems 2.1 and 2.3.

To show that the family of point processes $(\mathcal{N}_\varepsilon)_\varepsilon$ on $\mathbb{R}^+ \times V = E$ converges (with respect to some probability measure \mathbb{P}) to a Poisson point process with a σ -finite intensity η , we can apply Kallenberg's criterion (Proposition 3.22 in [18]). It suffices to prove that for some system \mathcal{R} stable by finite intersection and union of relatively compact open subsets, that generates the σ -algebra $\mathcal{V} \times \mathcal{B}(\mathbb{R}^+)$, the following holds:

For any $R \in \mathcal{R}$,

- (O) $\eta(\partial R) = 0$,
- (A) $\mathbb{E}(\mathcal{N}_\varepsilon(R)) \rightarrow \eta(R)$,
- (B) $\mathbb{P}(\mathcal{N}_\varepsilon(R) = 0) \rightarrow e^{-\eta(R)}$.

The last condition will be obtained from the simple next geometric approximation.

Proposition 2.4. *Let \mathcal{A} be a collection of measurable subsets of Ω . Then for any $r \geq 1$, any positive integers $p_1, \dots, p_r, q_1, \dots, q_r$ such that $p_i + q_i < p_{i+1}$ for any $i = 1, \dots, r-1$, and for any sets $A_1, \dots, A_r \in \mathcal{A}$ we have*

$$\left| \mu(\forall i, \tau_{A_i} \circ T^{p_i} > q_i) - \prod_{i=1}^r (1 - \mu(A_i))^{q_i} \right| \leq \sum_{i=1}^r q_i \Delta(\mathcal{A}),$$

with Δ defined in (4).

Proof. We will write $a = b \pm c$ to say that $|a - b| \leq c$. Note that for any integer $q_1 \geq 1$ we have

$$\{\tau_{A_1} > q_1\} = T^{-1}\{\tau_{A_1} > q_1 - 1\} - T^{-1}(A_1 \cap \{\tau_{A_1} > q_1 - 1\}).$$

Using the T -invariance of μ , this gives

$$\begin{aligned} & \mu(\forall i, \tau_{A_i} \circ T^{p_i} > q_i) \\ &= \mu(\forall i, \tau_{A_i} \circ T^{p_i - p_1} > q_i) \\ &= \mu(\tau_{A_1} > q_1 - 1; \forall i \geq 2, \tau_{A_i} \circ T^{p_i - p_1 - 1} > q_i) \\ &\quad - \mu(A_1 \cap \{\tau_{A_1} > q_1 - 1; \forall i \geq 2, \tau_{A_i} \circ T^{p_i - p_1 - 1} > q_i\}) \\ &= (1 - \mu(A_1))\mu(\tau_{A_1} > q_1 - 1; \forall i \geq 2, \tau_{A_i} \circ T^{p_i - p_1 + 1} > q_i) \pm \Delta(\mathcal{A}). \end{aligned}$$

By an immediate induction on q_1 we obtain

$$\mu(\forall i \geq 1, \tau_{A_i} \circ T^{p_i} > q_i) = (1 - \mu(A_1))^{q_1} \mu(\forall i \geq 2, \tau_{A_i} \circ T^{p_i - q_1} > q_i) \pm q_1 \Delta(\mathcal{A}).$$

The conclusion follows by an induction on the number r of sets. \square

Proof of Theorem 2.1. The strong convergence in distribution is by Theorem 1 in [23] a direct consequence of the classical convergence in distribution with respect to μ . The later will be proven as announced using Kallenberg's criterion with $\eta := \lambda \times m$.

Let \mathcal{R} be the collection of finite unions of open rectangles $I \times F$, where I is an open bounded interval in $[0, \infty)$ and $F \in \mathcal{W}$.

Let $R \in \mathcal{R}$. We rearrange the subdivision given by the endpoints of the intervals defining R to write $R = R' \cup \bigcup_{i=1}^r (t_i, s_i) \times F_i$, where $t_i < s_i$, $F_i \in \mathcal{W}$ for $i = 1, \dots, r$, $s_i \leq t_{i+1}$ for every $i = 1, \dots, r-1$ and R' is contained in a finite number of vertical strips $\{t\} \times V$. We assume without loss of generality that $R' = \emptyset$, as it will be clear from the sequel that the same arguments would give $\mathbb{E}_\mu[\mathcal{N}_\varepsilon(R')] \rightarrow 0$ and $\mu(\mathcal{N}_\varepsilon(R') = 0) \rightarrow 1$.

Note that for all $i = 1, \dots, r$, $m_\varepsilon(F_i) \rightarrow m(F_i)$ as $\varepsilon \rightarrow 0$ by Hypothesis (ii) and the Portemanteau theorem.

Condition (A) follows from the definition and the linearity of the expectation. Indeed,

$$\mathbb{E}_\mu[\mathcal{N}_\varepsilon(R)] = \sum_{i=1}^r \sum_{n=\lfloor \frac{t_i}{\mu(A_\varepsilon)} \rfloor + 1}^{\lceil \frac{s_i}{\mu(A_\varepsilon)} \rceil - 1} \mu(H_\varepsilon^{-1}(F_i)) \sim \sum_i (s_i - t_i) m_\varepsilon(F_i) \sim (\lambda \times m)(R).$$

For Condition (B), set $p_i = \lfloor t_i / \mu(A_\varepsilon) \rfloor$ and $q_i = \lceil s_i / \mu(A_\varepsilon) \rceil - 1 - p_i$ and $A_{\varepsilon,i} = H_\varepsilon^{-1} F_i$. Observe that $\mathcal{N}_\varepsilon(R) = 0$ is equivalent to $\forall i, \tau_{A_{\varepsilon,i}} \circ T^{p_i} > q_i$.

By Proposition 2.4 and due to assumption (i), we have

$$\begin{aligned}
\mu(\mathcal{N}_\varepsilon(R) = 0) &= \prod_i (1 - \mu(A_{\varepsilon,i}))^{q_i} \pm \sum_{i=1}^r q_i \Delta(\{A_{\varepsilon,i}\}) \\
&= \prod_i (1 - \mu(A_\varepsilon) m_\varepsilon(F_i))^{q_i} + o(1) \\
&= \exp\left(\sum_i q_i \mu(A_\varepsilon) m_\varepsilon(F_i)\right) + o(1) \\
&= \exp((\lambda \times m)(R)) + o(1).
\end{aligned}$$

□

Proof of Theorem 2.3. Due to Theorem 1 of [23], the assumption holds also with $\tilde{\mu}$ s.t. $\frac{d\tilde{\mu}}{d\mu} = \frac{\tau}{\bar{\tau}}$, and it suffices to prove the convergence in distribution of $(\mathcal{N}_\varepsilon)_\varepsilon$ with respect to ν .

By hypotheses, $(\mathcal{N}_\varepsilon)_\varepsilon$ converges in distribution wrt $\Pi_*\nu$ (which coincide with the probability measure on Ω with density $\tau/\bar{\tau}$ wrt μ) to a Poisson point process \mathcal{P} of intensity $\lambda \times m$, i.e. $(\mathcal{N}_\varepsilon \circ \Pi)_\varepsilon$ converges in distribution, with respect to ν , to \mathcal{P} .

Note that for $y = (x, s) \in \mathcal{M}$

$$\mathfrak{N}_\varepsilon(y) = (\psi_{\varepsilon,y})_*(\mathcal{N}_\varepsilon(\Pi(y)))$$

with

$$\psi_{\varepsilon,y}(t, z) = ((S_{\lfloor t/\mu(A_\varepsilon) \rfloor} \tau(\Pi y) - s) \mu(A_\varepsilon) / \bar{\tau}, z)$$

We conclude using the facts that $\mu(A_\varepsilon) \rightarrow 0$ and that $S_n \tau / (n \bar{\tau}) \rightarrow 1$ ν -almost surely. □

3. A COMMON FRAMEWORK SUITABLE FOR SYSTEMS MODELED BY GIBBS-MARKOV-YOUNG TOWERS

3.1. General result. The authors have studied in [16] the case of a temporal Poisson point process, corresponding to $V = \{0\}$ and $H_\varepsilon \equiv 0$ (Theorem 3.5 therein appears as Theorem 2.1 of the present paper in this specific context). In the above mentioned article, the general context was the case of dynamical systems modeled by a Gibbs Markov Young tower as studied in [1].

Hypothesis 3.1. Let $\alpha, \beta > 0$. Let Ω be a metric space endowed with a Borel probability measure μ and a μ -preserving transformation T . Assume that there exists a sequence of finite partitions $(\mathcal{Q}_k)_{k \geq 0}$ of an extension $(\tilde{\Omega}, \tilde{\mu}, \tilde{T})$ of (Ω, μ, T) by $\tilde{\Pi} : \tilde{\Omega} \rightarrow \Omega$ such that one of the two following assumptions holds:

- (I) either $\sup_{Q \in \mathcal{Q}_k} \text{diam}(\tilde{\Pi}(Q)) \leq Ck^{-\alpha}$,
- (II) or $\sup_{Q \in \mathcal{Q}_{2k}} \text{diam}(\tilde{\Pi} \tilde{T}^k Q) \leq Ck^{-\alpha}$.

Assume moreover that, there exists $C' > 0$ such that, for every k, n with $n \geq 2k$, for any $A \in \sigma(\mathcal{Q}_k)$ and for any $B \in \sigma(\bigcup_{m \geq 0} \mathcal{Q}_m)$,

$$\left| \text{Cov}_{\tilde{\mu}}(\mathbf{1}_A, \mathbf{1}_B \circ \tilde{T}^n) \right| \leq C' n^{-\beta} \tilde{\mu}(A).$$

Case (I) is appropriate for non-invertible systems, whereas Case (II) is appropriate for invertible systems. Wide classes of examples of dynamical systems satisfying these assumptions can be found in [1] (see also [21, 22]).

Proposition 3.2. *Let $(\Omega, \mathcal{F}, \mu, T)$ satisfying Hypothesis 3.1. With the notations of the beginning of Section 2.1, assume that there exists $p_\varepsilon = o(\mu(A_\varepsilon)^{-1})$ such that*

- (i) $\mu(\tau_{A_\varepsilon} \leq p_\varepsilon | A_\varepsilon) = o(1)$,
- (ii) $\mu((\partial A_\varepsilon)^{\lfloor C 4^\alpha p_\varepsilon^{-\alpha} \rfloor}) = o(\mu(A_\varepsilon))$
- (iii) $\mu(H_\varepsilon^{-1}(\cdot) | A_\varepsilon)$ converges vaguely to some measure m ,
- (iv) for all $F \in \mathcal{W}$, $m(\partial F) = 0$ and $\mu((\partial(H_\varepsilon^{-1}F))^{\lfloor C 4^\alpha p_\varepsilon^{-\alpha} \rfloor} | A_\varepsilon) = o(1)$.

Then the assumptions of Theorem 2.1 are satisfied.

We postpone the proof to the appendix. When assumption (iii) is missing, one has to change (iv) accordingly, going through a subsequence $\mathcal{E} = (\varepsilon_k)$, a family $\mathcal{W}_\mathcal{E}$ and a limit point m to get that the assumptions of Theorem 2.2 are satisfied.

Remark 3.3. • Notice that the Assumption (i) is always satisfied when the normalized first return time $\mu(A_\varepsilon)\tau_{A_\varepsilon}$ to A_ε is asymptotically exponentially distributed with parameter one, in particular when the temporal process (corresponding to $V = \{0\}$) converges in distribution to a Poisson process of intensity λ .
 • It may happen that one only has a nonuniform control of the diameters instead of Hypothesis 3.1 (I) or (II). Indeed it is possible to avoid these hypotheses. It suffices to replace (ii) with

$$\mu(\partial A_\varepsilon^{\lfloor k_\varepsilon \rfloor}) = o(\mu(A_\varepsilon)),$$

where in the non invertible case (I) $\partial A_\varepsilon^{\lfloor k_\varepsilon \rfloor} = \cup \tilde{\Pi}(Q)$, the union begin on those $Q \in \mathcal{Q}_{k_\varepsilon}$ such that $\text{diam } \tilde{\Pi}(Q) > Ck_\varepsilon^{-\alpha}$ and $\partial A_\varepsilon \cap \tilde{\Pi}(Q) \neq \emptyset$, $k_\varepsilon = \lfloor p_\varepsilon/2 \rfloor$; in the invertible case (II) $\partial A_\varepsilon^{\lfloor k_\varepsilon \rfloor} = \cup \tilde{\Pi}(T^{k_\varepsilon}Q)$, the union begin on those $Q \in \mathcal{Q}_{2k_\varepsilon}$ such that $\text{diam } \tilde{\Pi}(T^{k_\varepsilon}Q) > Ck_\varepsilon^{-\alpha}$ and $\partial A_\varepsilon \cap \tilde{\Pi}(T^{k_\varepsilon}Q) \neq \emptyset$, $k_\varepsilon = \lfloor p_\varepsilon/4 \rfloor$; The modification of the proof of Proposition 3.2 is immediate.

The following result is helpful to get assumption (iv) in many cases.

Proposition 3.4. *Assume that V is an open subset of \mathbb{R}^d , for some $d > 0$, that H_ε are \mathfrak{h} -Hölder continuous maps with respective Hölder constant $C_\mathfrak{h}(H_\varepsilon)$, that there exists $\eta_\varepsilon \rightarrow 0+$ such that $C_\mathfrak{h}(H_\varepsilon)\eta_\varepsilon^\mathfrak{h} \rightarrow 0$ and that $m_\varepsilon \rightarrow m$ (where m is a finite measure on V). Then there exists a family \mathcal{W} of relatively compact open subsets of B , stable by finite unions and intersections, generating the Borel σ -algebra of V , such that*

- (i) $m(\partial F) = 0$ for any $F \in \mathcal{W}$,
- (ii) for any $F \in \mathcal{W}$,

$$\mu((\partial H_\varepsilon^{-1}F)^{\lfloor \eta_\varepsilon \rfloor} | A_\varepsilon) = o(1).$$

Proof. (i) Let $\pi_j : \mathbb{R}^d \rightarrow \mathbb{R}$ be the j -th canonical projection (i.e. $\pi_j((x_i)_i) = x_j$). The set $\mathcal{G}^j := \{a \in \mathbb{R} : (\pi_j)_* m(a) = 0\}$ is dense in \mathbb{R} , since its complement is at most countable.

We then define \mathcal{W} as the collection of open rectangles $\prod_{j=1}^d (a_j; b_j) \subset B$, with $a_j, b_j \in \mathcal{G}^j$. By construction $m(\partial F) = 0$ for any $F \in \mathcal{W}$. The density of the \mathcal{G}^j 's implies that \mathcal{W} generates the Borel σ -algebra.

(ii) For any $F \in \mathcal{W}$ we have the inclusion

$$(\partial H_\varepsilon^{-1} F)^{[\eta_\varepsilon]} \subset H_\varepsilon^{-1} \partial F^{[C_h(H_\varepsilon) \eta_\varepsilon^h]}.$$

Hence, for every $\varepsilon_0 > 0$,

$$\mu((\partial H_\varepsilon^{-1} F)^{[\eta_\varepsilon]} | A_\varepsilon) \leq m_\varepsilon(\partial F^{[C_h(H_\varepsilon) \eta_\varepsilon^h]}) \leq m_\varepsilon(\partial F^{[M_{\varepsilon_0}]}) ,$$

for any $0 < \varepsilon < \varepsilon_0$, with $M_{\varepsilon_0} := \sup_{\varepsilon \in (0, \varepsilon_0)} C_h(H_\varepsilon) \eta_\varepsilon^h$. Therefore

$$\lim_{\varepsilon \rightarrow 0} \mu((\partial H_\varepsilon^{-1} F)^{[\eta_\varepsilon]} | A_\varepsilon) \leq m(\partial F^{[M_{\varepsilon_0}]}),$$

which goes to 0 as $\varepsilon_0 \rightarrow 0$ since $m(\partial F) = 0$. \square

3.2. Successive visits in a small neighbourhood of a generic point.

The purpose of the next result is to give examples for which Theorem 2.1 applies for returns in small balls, in the same context as in [16].

Theorem 3.5. *Assume that Ω is a d -Riemannian manifold and that $\dim_H \mu = \lim_{\varepsilon \rightarrow 0} \frac{\log \mu(B(x_0, \varepsilon))}{\log \varepsilon}$ for μ -almost every $x_0 \in \Omega$. Assume Hypothesis 3.1 with $\alpha > \dim_H \mu$.*

Then for μ -almost every $x_0 \in \Omega$ such that

$$\exists \delta \in (1, \alpha \dim_H \mu), \quad \mu(B(x_0, \varepsilon) \setminus B(x_0, \varepsilon - \varepsilon^\delta)) = o(\mu(B(x_0, \varepsilon))), \quad (7)$$

the family of point processes

$$\mathcal{N}_\varepsilon(x) = \sum_{n: T^n(x) \in B(x_0, \varepsilon)} \delta_{(n\mu(B(x_0, \varepsilon), \varepsilon^{-1} \exp_{x_0}^{-1}(T^n x))}$$

is strongly approximated in distribution by a Poisson process \mathcal{P}_ε of intensity $\lambda \times m_\varepsilon^{x_0}$ where $m_\varepsilon^{x_0} = \mu(H_\varepsilon^{-1}(\cdot) | B(x_0, \varepsilon))$.

Proof of Theorem 3.5. We apply Proposition 3.2, assuming without loss of generality that the measures $m_\varepsilon^{x_0}$ are converging.

Fix $\sigma \in (\delta/\alpha, \dim_H \mu)$ and set $p_\varepsilon = \varepsilon^{-\sigma}$. The assumptions imply that the temporal return times process converges in distribution to a Poisson process of parameter one (See [16]). By Remark 3.3 this shows that Assumption (i) of Proposition 3.2 is true. Assumption (ii) follows from (7). Finally, the family \mathcal{W} comes from Proposition 3.4 above, which proves also the last assumption (iv) and thus the theorem. \square

4. APPLICATIONS TO BILLIARD MAPS AND FLOWS

4.1. Bunimovich billiard. The Bunimovich billiard is an example of weakly hyperbolic system (with polynomial decay of the covariance of Hölder functions).

Let $\ell > 0$. We consider the planar domain Q union of the rectangle $[-\ell/2; \ell/2] \times [-1, 1]$ and of the two planar discs of radius 1 centered at $(\pm \ell/2, 0)$. We consider a point particle moving with unit speed in Q , going straight on between two reflections off ∂Q and reflecting with respect to the classical Descartes law of reflection (incident angle=reflected angle). The billiard system (Ω, μ, T) describes the evolution at reflected times of a point

particle moving in the domain Q as described above. We define the set Ω of reflected vectors as follows


$$\Omega := \{\partial Q \times S^1 : \langle \vec{n}_q, \vec{v} \rangle \geq 0\},$$

where \vec{n}_q is the unit normal vector normal to ∂Q , directed inside Q , at q . Such a reflected vector (q, \vec{v}) can be represented by $x = (r, \varphi) \in \mathbb{R}/(2(\pi + \ell)\mathbb{Z})$, r corresponding to the counterclockwise curvilinear abscissa of q on ∂Q (starting from a fixed point on ∂Q) and φ measuring the angle between \vec{n}_q and \vec{v} . The transformation T maps a reflected vector to the reflected vector corresponding to the next reflection time. This transformation preserves the measure μ given (in coordinates) by $d\mu(r, \varphi) = h(r, \varphi) dr d\varphi$ where $h(r, \varphi) = \frac{\cos \varphi}{2|\partial Q|}$. We endow Ω with the supremum metric $d((r, \varphi), (r', \varphi')) = \max(|r - r'|, |\varphi - \varphi'|)$.

Theorem 4.1. *For μ -almost every $x_0 = (r_0, \varphi_0) \in \Omega$, the family of point processes*

$$\sum_{n \geq 1 : d(T^n(x), x_0) < \varepsilon} \delta_{\left(n\varepsilon^2 h(x_0), \frac{T^n(x) - x_0}{\varepsilon}\right)}$$

converges in distribution (with x distributed with respect to any probability measure absolutely continuous with respect to the Lebesgue measure on Ω) to a Poisson Point Process with intensity $\lambda \times \lambda_2$, where here λ_2 is the 2-dimensional normalized Lebesgue measure on $(-1, 1)^2$.



Proof. The convergence comes directly from Theorem 3.5, case (II) with $\alpha = 1$, $\zeta = 1$ and $\dim_H \mu = 2$, (due to [16], namely in Section 9 therein) and from the fact that $\mu(H_\varepsilon^{-1}(\cdot)|A_\varepsilon)$ converges in distribution to the normalized Lebesgue measure on $B_{|\cdot|_\infty}(0, 1)$. To identify the intensity we observe that $\mu(B(x_0, \varepsilon)) \sim \cos \varphi_0 \varepsilon^2 / (2|\partial Q|)$ as $\varepsilon \rightarrow 0+$. \square

We now consider the billiard flow $(\mathcal{M}, \nu, (Y_t)_t)$ with

$$\mathcal{M} = \{(q, \vec{v}) \in Q \times S^1 : q \in \partial Q \Rightarrow \langle \vec{n}_q, \vec{v} \rangle \geq 0\},$$

where $Y_t(q, \vec{v}) = (q', \vec{v}')$ if a particle that was at time 0 at position q with speed \vec{v} will be at time t at position q' with speed \vec{v}' and where ν is the normalized Lebesgue measure on \mathcal{M} .

An important well known fact is that the billiard flow system $(\mathcal{M}, \nu, (Y_t)_t)$ can be represented as the special flow over the billiard map system (Ω, μ, T) with roof function $\tau : \Omega \rightarrow (0, +\infty)$ given by $\tau(x) := \inf\{t > 0 : Y_t(x) \in \partial Q \times S^1\}$. This enables us to get the following result.

Corollary 4.2. *For μ -almost every $x_0 = (r_0, \varphi_0) \in \Omega$, the family of point processes*

$$\sum_{t>0 : Y_t(y) \in B_\Omega(x_0, \varepsilon)} \delta_{\left(\frac{t\varepsilon^2 \cos \varphi_0}{2\pi \text{Area}(Q)}, \frac{Y_t(y) - x_0}{\varepsilon}\right)}$$

converges in distribution (with respect to any probability measure absolutely continuous with respect to the Lebesgue measure on \mathcal{M}) to a Poisson Point Process with intensity $\lambda \times \lambda_2$, where here λ_2 is the 2-dimensional normalized Lebesgue measure on $[-1, 1]^2$ and where $B_\Omega(x_0, \varepsilon)$ means the ball in Ω .

Proof. Set $\bar{\tau} := \int_\Omega \tau d\mu$. Due to Theorems 4.1 and 2.3, the point process

$$\sum_{t>0 : Y_t(y) \in B_\Omega(x_0, \varepsilon)} \delta_{\left(\frac{t\varepsilon^2 \cos \varphi_0}{2\bar{\tau}|\partial Q|}, \frac{Y_t(y) - x_0}{\varepsilon}\right)}$$

converges in distribution to a Poisson Point Process with intensity $\lambda \times \lambda_2$. Moreover $2\pi \text{Area}(Q) = \int_\Omega \tau \cos \varphi dr d\varphi = 2|\partial Q|\bar{\tau}$. \square

4.2. Sinai billiards. Let $I \in \mathbb{N}^*$ and O_1, \dots, O_I be open convex subsets of \mathbb{T}^2 with C^3 -smooth boundary of positive curvature, and pairwise disjoint closures. We then set $Q = \mathbb{T}^2 \setminus \bigcup_{i=1}^I O_i$. As for the Bunimovich billiard, we consider a point particle moving in Q , with unit speed and elastic reflections off ∂Q . This model is called the Sinai billiard. We assume moreover that the horizon is finite, i.e. that the time between two reflections is uniformly bounded.

For this choice of Q , we consider now the billiard map (Ω, μ, T) and the billiard flow $(\mathcal{M}, \nu, (Y_t)_t)$ defined as for the Bunimovich billiard in Subsection 4.1.

Under this finite horizon assumption, the Sinai billiard has much stronger hyperbolic properties than the Bunimovich billiard (with namely an exponential decay of the covariance of Hölder functions), but nevertheless, compared to Anosov map, its study is complicated by the presence of discontinuities.

Theorem 4.3. *For μ -almost every $x_0 = (q_0, \vec{v}_0) \in \Omega$ (represented by (r_0, φ_0)),*

- **[Return times in a neighborhood of x_0]** *The conclusions of Theorem 4.1 and of its Corollary 4.2 hold also true.*
- **[Return times in a neighborhood of the position q_0 of x_0]** *The family of point processes*

$$\sum_{n \geq 1 : T^n(x) \in B_{\partial Q}(q_0, \varepsilon) \times S^1} \delta_{\left(\frac{2\varepsilon n}{|\partial Q|}, \frac{r - r_0}{\varepsilon}, \varphi\right)}$$

(where x is represented by (r, φ)) converges in distribution (with respect to any probability measure absolutely continuous with respect to the Lebesgue measure on Ω) to a Poisson Point Process with intensity $\lambda \times m_0$, where here m_0 is the probability measure on $[-1, 1] \times [-\pi/2; \pi/2]$ with density $(r, \varphi) \mapsto \frac{\cos(\varphi)}{4}$.

- *The family of point processes*

$$\sum_{t>0 : Y_t(y) \in B_Q(q_0, \varepsilon) \times S^1} \delta_{\left(\frac{2\varepsilon t}{\pi \text{Area}(Q)}, \frac{r - r_0}{\varepsilon}, \varphi\right)}$$

converges in distribution (with respect to any probability measure absolutely continuous with respect to the Lebesgue measure on \mathcal{M}) to a Poisson Point Process with intensity $\lambda \times m_0$, with m_0 as above.

Due to [21], the Sinai billiard (Ω, μ, T) satisfies Hypothesis 3.1-(II) with any $\alpha > 0$ and any $\beta > 0$.

Proof. The first item follows from Theorem 3.5 as in Theorem 4.1 and Corollary 4.2.

The third item follows from the second one as Corollary 4.2 comes from Theorem 4.1.

Let us prove the second item. Let $x_0 = (q_0, \varphi_0) \in \Omega$. We set $A_\varepsilon(x_0) := [q_0 - \varepsilon, q_0 + \varepsilon] \times [\pi/2, \pi/2]$ and $H_\varepsilon : (q, \varphi) \mapsto (\varepsilon^{-1}(q - q_0), \varphi)$. Note first that $\mu(A_\varepsilon) = 2\varepsilon/|\partial Q|$ and that $\mu(H_\varepsilon^{-1}(\cdot)|A_\varepsilon)$ converges vaguely to m_0 .

We follow verbatim the proof of Theorem 3.5 which invokes Proposition 3.2, except for its assumption (i):

The convergence of the temporal process as in the proof of Theorem 3.5, associated to the position of the billiard particle, is not present in the literature in these terms. However, a slight adaptation of [15, Lemma 6.4-(iii)] gives that for any $\sigma < 1$ and μ -a.e. $x_0 \in \partial Q \times]-\frac{\pi}{2}; \frac{\pi}{2}[$ we have

$$\mu(\tau_{A_\varepsilon} \leq \varepsilon^{-\sigma} | A_\varepsilon) = o(1).$$

That is (i) of Proposition 3.2 holds with $p_\varepsilon = \varepsilon^{-\sigma}$. \square

Let us write $\Pi_Q : \mathcal{M} \rightarrow Q$ and $\Pi_V : \mathcal{M} \rightarrow S^1$ for the two canonical projections, which correspond respectively to the position and to the speed. Using results established in [15], we also state a result of convergence to a spatio-temporal Poisson point process for entrance times in balls for the flows. We endow \mathcal{M} with the metric d given by $d((q, \vec{v}), (q', \vec{v}')) = \max(d_0(q, q'), |\angle(\vec{v}, \vec{v}')|)$, where d_0 is the euclidean metric in Q and where $\angle(\cdot, \cdot)$ is the angular measure of the angle.


Theorem 4.4. *For ν -a.e. $y_0 = (q_0, \vec{v}_0) \in \mathcal{M}$,*

- *the family of point processes*

$$\sum_{t : (Y_s(y))_s \text{ enters } B(y_0, \varepsilon) \text{ at time } t} \delta \left(\frac{2\varepsilon^2 t}{\pi \text{Area}(Q)}, \frac{\Pi_Q(Y_t(y)) - q_0}{\varepsilon}, \frac{\angle(\vec{v}_0, \Pi_V(Y_t(y)))}{\varepsilon} \right)$$

converges in distribution (when y is distributed with respect to any probability measure absolutely continuous with respect to the Lebesgue measure on \mathcal{M}) to a Poisson Point Process with intensity $\lambda \times \tilde{m}_1$, where \tilde{m}_1 is the probability measure of density $(p, \vec{u}) \mapsto \frac{1}{4} \langle \tilde{n}_p, \vec{v}_0 \rangle^+$ on $S^1 \times [-1, 1]$ (with $\langle \cdot, \cdot \rangle^+$ the positive part of the scalar product in \mathbb{R}^2 and \tilde{n}_p the inward normal vector to S^1 at p).³


³In the limit, the authorized normalized positions are the positions located on a semi-circle (corresponding to positions at which the vector \vec{v}_0 enters the ball) and the normalized variation of speed is uniform in $[-1, 1]$



- the family of point processes

$$\sum_{t : (Y_s(y))_s \text{ enters } B(q, \epsilon) \times S^1 \text{ at time } t} \delta \left(\frac{2\pi\epsilon t}{\text{Area}(Q)}, \frac{\Pi_Q(Y_t(y)) - q_0}{\epsilon}, \Pi_V(Y_t(y)) \right)$$

converges in distribution (when y is distributed with respect to any probability measure absolutely continuous with respect to the Lebesgue measure on \mathcal{M}) to a Poisson Point Process with intensity $\lambda \times \tilde{m}_0$ where \tilde{m}_0 is the probability measure with density $(p, \vec{u}) \mapsto \frac{1}{4\pi} \langle \tilde{n}_p, \vec{u} \rangle^+$ on $S^1 \times S^1$.⁴



Proof. We apply Theorem 2.3 to go from the discrete time to the continuous time. Let $\mathcal{A}_\epsilon := B(x, \epsilon)$ (resp. $\mathcal{A}_\epsilon := B(q, \epsilon) \times S^1$). Return times to these sets have already been studied in [15]. We set $A_\epsilon := \Pi \mathcal{A}_\epsilon$ and study the discrete time process associated to these sets. To this end, we apply Proposition 3.2 after checking its assumptions.

- We know that $\mu(A_\epsilon) = 2\epsilon^2/|\partial Q|$ due to [15, Lemma 5.1] (resp. $\mu(A_\epsilon) = 2\pi\epsilon/|\partial Q|$ due to [15, Lemma 5.1]). So $d = 2$ (resp. $d = 1$). Let $\sigma < d$ and $\delta > 1$.
- Note that $\mu((\partial A_\epsilon)^{[\epsilon^\delta]}) = o(\mu(A_\epsilon))$. Due to [15, Theorem 3.3] (resp. [15, Lemma 6.4], $\mu(\tau_{A_\epsilon} \leq \epsilon^{-\sigma} | A_\epsilon) = o(1)$.
- We define $\mathcal{B}_\epsilon := \{(q, \vec{v}) \in \partial B(x_0, \epsilon) \times S^1 : \langle \tilde{n}_q, \vec{v} \rangle \geq 0\}$. We endow it with the measure $\tilde{\mu}$ given by $d\tilde{\mu}(q, \vec{v}) = \cos \varphi dr d\varphi$ with $\varphi = \angle(\tilde{n}_q, \vec{v})$ and r the curvilinear abscissa of q on $\partial B(x_0, \epsilon)$.
- Let $\tau_{A_\epsilon}(y) := \inf\{t > 0 : Y_t(y) \in A_\epsilon\}$.
- We define $H_\epsilon : A_\epsilon \rightarrow S^1 \times [-1, 1]$ which maps $x = (q, \vec{v}) \in A_\epsilon$ to $\epsilon^{-1}(\Pi_Q(Y_{\tau_{A_\epsilon}}(x)) - q_0, \angle(\vec{v}_0, \vec{v}))$ (resp. $H_\epsilon : A_\epsilon \rightarrow S^1 \times S^1$ which maps $x = (q, \vec{v}) \in A_\epsilon$ to $(\epsilon^{-1}\Pi_Q(Y_{\tau_{A_\epsilon}}(x)) - q_0, \vec{v})$).
- Note that the image measure of $\mu(\cdot | A_\epsilon)$ by $x \mapsto Y_{\tau_{A_\epsilon}}(x)$ corresponds to $\tilde{\mu}(\cdot | Y_{\tau_{A_\epsilon}}(A_\epsilon))$. The set $Y_{\tau_{A_\epsilon}}(A_\epsilon)$ consists of points of \mathcal{B}_ϵ such that $\angle(\vec{v}_0, v) < \epsilon$ (resp. $Y_{\tau_{A_\epsilon}}(A_\epsilon) = \mathcal{B}_\epsilon$).

⁴In the limit the authorized vectors are the unit vectors entering the ball.

- Hence $\mu(H_\varepsilon^{-1}(\cdot)|A_\varepsilon)$ is the image measure of $\mu(\cdot|Y_{\tau_{A_\varepsilon}}(A_\varepsilon))$ by $(q, \vec{v}) \mapsto \varepsilon^{-1}(q - q_0, \langle \vec{v}_0, \vec{v} \rangle)$ (resp. $\mu(H_\varepsilon^{-1}(\cdot)|A_\varepsilon)$ is the image measure of $\mu(\cdot|\mathcal{B}_\varepsilon)$ by $(q, \vec{v}) \mapsto (\varepsilon^{-1}q, \vec{v})$).

Hence we obtain the convergence in distribution of these families of measures.

- For the construction of \mathcal{W} we use Proposition 3.4.

□

5. SUCCESSIVE VISITS IN A SMALL NEIGHBORHOOD OF AN HYPERBOLIC PERIODIC POINT

5.1. General results around a periodic hyperbolic orbit. We consider the case of a periodic point x_0 of smallest period p (i.e. p is the smallest $n > 0$ such that $T^n x_0 = x_0$).

By periodicity, returns to $B(x_0, \varepsilon)$ appear in clusters, so that we cannot hope that the return process is represented by a simple Poisson process. However, the occurrence of clusters should be well separated and have a chance to be represented by a simple Poisson Process.

Thus we define A_ε as the set of points of $B(x_0, \varepsilon)$ leaving $B(x_0, \varepsilon)$ for a time at least q_0 , i.e.

$$A_\varepsilon := B(x_0, \varepsilon) \setminus \bigcup_{j=1}^{q_0} T^{-jp} B(x_0, \varepsilon)$$

and consider \mathcal{N}_ε defined by (1) with this choice of A_ε . This definition of A_ε essentially records the last passage among a series of hitting to the ball. We emphasize that in general, one has to consider $q_0 > 1$ to avoid clustering of occurrences of A_ε due to finite time effects⁵.

Lemma 5.1. *Assume x_0 is a hyperbolic fixed point of T and that T is $C^{1+\alpha}$ in a neighborhood U of the orbit $x_0, \dots, T^{p-1}x_0$. Then there exist an integer q_0 and a $a > 0$ such that for any $\varepsilon > 0$ sufficiently small, for any $n = 1, \dots, \lfloor a \log 1/\varepsilon \rfloor$, $A_\varepsilon \cap T^{-n} A_\varepsilon = \emptyset$, where $A_\varepsilon := B(x_0, \varepsilon) \setminus \bigcup_{j=1}^{q_0} T^{-jp} B(x_0, \varepsilon)$.*

For $\varepsilon > 0$ small enough, we define, as in Theorem 3.5, $H_\varepsilon : B(x_0, \varepsilon) \mapsto B(0, 1)$ by $H_\varepsilon := \varepsilon^{-1} \exp^{-1}$. We are interested in the behaviour of the point processes $(\tilde{\mathcal{N}}_\varepsilon)_\varepsilon$ defined by

$$\tilde{\mathcal{N}}_\varepsilon(x) := \sum_{n : T^n(x) \in B(x_0, \varepsilon)} \delta_{(n\mu(B(x_0, \varepsilon)), H_\varepsilon(T^n(x)))}. \quad (8)$$

⁵Indeed, consider the determinant one hyperbolic matrix $M = \begin{pmatrix} -0.2 & 1.8 \\ 0.6 & -0.4 \end{pmatrix}$. The vector $v = \begin{pmatrix} 0.5 \\ 0.7 \end{pmatrix}$ belongs to the unit ball B_1 , $Mv \notin B_1$, $M^2v \in B_1$ and $M^3v \notin B_1$. Assume that T preserves the Lebesgue measure μ and has a fixed point x_0 such that $D_{x_0}T = M$. Let $A_\varepsilon = B(x_0, \varepsilon) \setminus T^{-1}B(x_0, \varepsilon)^c$. One easily shows that the inequality

$$\mu(A_\varepsilon \cap T^{-2}A_\varepsilon) \geq \mu(B(x_0, \varepsilon) \cap T^{-1}B(x_0, \varepsilon)^c \cap T^{-2}B(x_0, \varepsilon) \cap T^{-3}B(x_0, \varepsilon)^c),$$

contradicts the assumption that $\Delta(\{A_\varepsilon\}) = o(\mu(A_\varepsilon))$.

The extremal index defined by

$$\theta_\varepsilon = \frac{\mu(A_\varepsilon)}{\mu(B(x_0, \varepsilon))}$$

relates the above process to

$$\tilde{\mathcal{N}}_\varepsilon(x) := \sum_{n : T^n(x) \in B(x_0, \varepsilon) \setminus \{x_0\}} \delta_{(n\mu(A_\varepsilon), H_\varepsilon(T^n(x)))} \quad (9)$$

in an obvious way. Indeed, $\tilde{\mathcal{N}}_\varepsilon = \Theta_\varepsilon(\tilde{\mathcal{N}}_\varepsilon)$ where

$$\Theta_\varepsilon \left(\sum_n \delta_{(t_n, x_n)} \right) = \sum_n \delta_{(\theta_\varepsilon^{-1} t_n, x_n)}. \quad (10)$$

We see these processes as point processes on $[0, +\infty) \times \dot{B}(0, 1)$ (where $\dot{B}(0, 1) = B(0, 1) \setminus \{0\}$ is the open punctured ball).

Assume that Ω is a d -Riemannian manifold. A p -periodic point of T is said to be hyperbolic if T^p defines a C^1 diffeomorphism between two neighborhoods of x_0 and if DT_{x_0} admits no eigenvalue of modulus 1. We write $E_{x_0}^u$ for the spectral space associated to eigenvalues of modulus strictly larger than 1.

Theorem 5.2. *Assume that T^{-1} is well defined on a small neighborhood of x_0 and that $(\mathcal{N}_\varepsilon)_\varepsilon$ converges in distribution to a Poisson point process \mathcal{P} with intensity $\lambda \times m$, then the sequence of point processes $(\tilde{\mathcal{N}}_\varepsilon)_\varepsilon$ converges in distribution to the point process $\mathcal{N} = \Psi(\mathcal{P})$ on $[0, +\infty) \times \dot{B}(0, 1)$, with*

$$\Psi \left(\sum_n \delta_{(t_n, x_n)} \right) := \sum_{n : x_n \neq 0} \sum_{k=0}^{\ell_{x_n}} \delta_{(t_n, DT_{x_0}^{-kp}(x_n))},$$

where $\ell_y := \inf\{k \geq 0 : DT_{x_0}^{-kp}(y) \in B(0, 1) \setminus \bigcup_{j=1}^{q_0} DT_{x_0}^{jp} B(0, 1)\}$.

Proof. Note that m has support in $B(0, 1) \setminus \bigcup_{j=1}^{q_0} DT_{x_0}^{jp} B(0, 1)$.

Observe that, for every ε small enough, $\tilde{\mathcal{N}}_\varepsilon(x)$ is the image measure of $\mathcal{N}_\varepsilon(x)$ by

$$\Psi_\varepsilon : \sum_n \delta_{(t_n, x_n)} \mapsto \sum_n \sum_{k=0}^{\ell_{x_n, \varepsilon}} \delta_{(t_n - kp\mu(A_\varepsilon), H_\varepsilon T^{-kp} H_\varepsilon^{-1}(x_n))},$$

with

$$\ell_{y, \varepsilon} := \inf\{k \geq 0 : H_\varepsilon T^{-kp}(H_\varepsilon^{-1}y) \in B(0, 1) \setminus \bigcup_{j=1}^{q_0} T^{jp} B(0, 1)\}.$$

Observe that, for every $x \in B(0, 1)$,

$$\lim_{\varepsilon \rightarrow 0} \ell_{x, \varepsilon} = \ell_x,$$

and that, for every k , every $t > 0$ and every $x \in B(0, 1)$,

$$\lim_{\varepsilon \rightarrow 0} (t - kp\mu(A_\varepsilon), H_\varepsilon T^{-kp} H_\varepsilon^{-1}(x)) = (t, DT_{x_0}^{-kp}(x)).$$

Let us consider a set $R = \cup_i]r_i, s_i[\times F_i$ such that $\mathbb{E}[\mathcal{N}(\partial R)] = 0$ where $0 \leq r_i < s_i \leq r_{i+1}$ and where F_i are open precompact subsets of $\dot{B}(0, 1)$.

Note that, since the closure of $\cup_i F_i$ does not contain 0, there exists $K > 0$ and $\varepsilon_0 > 0$ (depending on the F_i 's) such that

$$\sup_{\varepsilon \in (0, \varepsilon_0), y \in \cup_i F_i} \inf\{k \geq 0 : H_\varepsilon^{-1} T^{kp} H_\varepsilon(y) \in A_\varepsilon\} < K$$

and

$$\sup_{\varepsilon \in (0, \varepsilon_0), y \in \cup_i F_i} \inf\{k \geq 0 : H_\varepsilon^{-1} T^{-kp} H_\varepsilon(x) \in B(0, 1) \setminus \bigcup_{j=1}^{q_0} T^{jp} B(0, 1)\} < K$$

so that

$$\forall y \in \bigcup_i F_i, \left[\exists x \in A_\varepsilon, \exists k \in \{0, \dots, \ell_{x, \varepsilon}\}, y = H_\varepsilon T^{-kp}(x) \right] \Rightarrow \ell_{x, \varepsilon} \leq 2K.$$

By definition of $\ell_{x, \varepsilon}$, for every $y \in \cup_i F_i$, every $x \in A_\varepsilon$, every $k \in \{0, \dots, \ell_{x, \varepsilon}\}$ such that $y = H_\varepsilon T^{-kp}(x)$, we have:

$$x = T^{\tau_{A_\varepsilon}^{(0)}}(y)$$

with $\tau_A^{(0)}(y) := \inf\{k \geq 0 : T^k(y) \in A\}$. Due to lemma 5.1, there exists $\varepsilon_1 \in (0, \varepsilon_0)$ such that, for every $\varepsilon \in (0, \varepsilon_1)$,

$$\forall y \in \bigcup_i F_i, \quad \{T^k(y), k = 0, \dots, K\} \cap A_\varepsilon = \{T^{\tau_{A_\varepsilon}^{(0)}}(y)\}.$$

Therefore, for every $\varepsilon \in (0, \varepsilon_1)$

$$\tilde{\mathcal{N}}_\varepsilon(R) = \tilde{\Psi}_{\varepsilon, K}(\mathcal{N}_\varepsilon)(R) = \mathcal{N}_\varepsilon \left(\bigcup_{k=0}^K \varphi_{\varepsilon, k}^{-1}(R) \right),$$

with

$$\tilde{\Psi}_{\varepsilon, K} : \sum_n \delta_{(t_n, x_n)} \mapsto \sum_n \sum_{k=0}^K \delta_{(t_n - kp\mu(A_\varepsilon), H_\varepsilon T^{-kp} H_\varepsilon^{-1}(x_n))}$$

and with $\varphi_{\varepsilon, k} : (t, x) \mapsto (t - kp\mu(A_\varepsilon), H_\varepsilon T^{-kp} H_\varepsilon^{-1}(x))$. Arguing analogously for \mathcal{N} and \mathcal{P} instead of $\tilde{\mathcal{N}}_\varepsilon$ and \mathcal{N}_ε , we obtain

$$\mathcal{N}(R) = \tilde{\Psi}_K(\mathcal{P})(R) = \mathcal{P} \left(\bigcup_{k=0}^K \varphi_k^{-1}(R) \right)$$

and

$$(\lambda \times m) \left(\bigcup_{k=0}^K \varphi_k^{-1}(\partial R) \right) = \mathbb{E}[\mathcal{N}(\partial R)] = 0,$$

with

$$\tilde{\Psi}_K : \sum_n \delta_{(t, x)} \mapsto \sum_n \sum_{k=0}^K \delta_{(t_n, H_\varepsilon T^{-kp} H_\varepsilon^{-1}(x_n))}.$$

and

$$\varphi_k : (t, x) \mapsto (t\mu(A_\varepsilon), DT_{x_0}^{-kp}(x)).$$

Since $(\mathcal{N}_\varepsilon)_\varepsilon$ converges in distribution to \mathcal{P} , we conclude that $\left(\mathcal{N}_\varepsilon \left(\bigcup_{k=0}^K \varphi_k^{-1}(R) \right)\right)_\varepsilon$ converges in distribution to $\mathcal{P} \left(\bigcup_{k=0}^K \varphi_k(R) \right)$. Moreover

$$\left| \mathcal{N}_\varepsilon \left(\bigcup_{k=0}^K \varphi_k^{-1}(R) \right) - \mathcal{N}_\varepsilon \left(\bigcup_{k=0}^K \varphi_{\varepsilon,k}(R) \right) \right| \leq \mathcal{N}_\varepsilon \left(\bigcup_{k=0}^K \varphi_k^{-1}(\partial R^{[\eta_\varepsilon]}) \right)$$

with $\lim_{\varepsilon \rightarrow 0} \eta_\varepsilon = 0$. Since $(\mathcal{N}_\varepsilon)_\varepsilon$ converges in distribution to \mathcal{P} and since $\mathcal{N}(\partial R) = 0$, we conclude that

$$\left(\mathcal{N}_\varepsilon \left(\bigcup_{k=0}^K \varphi_k^{-1}(R) \right) - \mathcal{N}_\varepsilon \left(\bigcup_{k=0}^K \varphi_{\varepsilon,k}(R) \right) \right)_\varepsilon$$

converges in distribution to 0. Therefore $(\tilde{\mathcal{N}}_\varepsilon(R))_\varepsilon$ converges in distribution to $\mathcal{N}(R)$. \square

Theorem 5.2 is designed for invertible systems, near a hyperbolic periodic point, and does not apply to expanding maps. Indeed, in such a non-invertible situation, one has to define the set A_ε with the first passage in the ball $B(x_0, \varepsilon)$ and not the last. More precisely, one has to set

$$A_\varepsilon = T^{-(q_0+1)}B(x_0, \varepsilon) \setminus \bigcup_{j=1}^{q_0} T^{-j}B(x_0, \varepsilon).$$

We leave to the reader the generalization of Theorem 5.2 and the result of the next section to this case.

5.2. SRB measure for Anosov maps. We now consider a C^2 Anosov map T on the d -dimensional riemaniann manifold Ω . We assume that the measure μ is the SRB measure of the system [12], and that x_0 is a periodic point of T of smallest period p .

Theorem 5.3. *We assume that $\mu(B(x_0, 2\varepsilon))\varepsilon^{b_0} = o(\mu(B(x_0, \varepsilon)))$ for some $b_0 > 0$ sufficiently small⁶. Then*

- (a) *The point process \mathcal{N}_ε for entrances in A_ε is asymptotically Poisson \mathcal{P}_ε , of intensity $\lambda \times m_\varepsilon$*
- (b) *The point process $\tilde{\mathcal{N}}_\varepsilon$ for entrances in $B(x_0, \varepsilon)$ is asymptotically $\Psi(\mathcal{P}_\varepsilon)$.*
- (c) *The point process $\tilde{\tilde{\mathcal{N}}}_\varepsilon$ for entrances in $B(x_0, \varepsilon)$ is asymptotically $\Theta_\varepsilon(\Psi(\mathcal{P}_\varepsilon))$.*
- (d) *The return time point process*

$$\mathcal{T}_\varepsilon := \sum_{n: T^n x \in B(x_0, \varepsilon)} \delta_{n\mu(B(x_0, \varepsilon))}$$

is asymptotically the compound Poisson point process $\pi\Theta_\varepsilon(\Psi(\mathcal{P}_\varepsilon))$, where π is the projection on the time axis.

The compound Poisson distribution (d) has already been established for few dynamical systems [9, 10, 8, 2], typically with strong assumptions on the dimension (1 or 1+1), the measure (non singular) and the shape of the balls (e.g. products of stable and unstable balls). We point out that

⁶Indeed this happens when for example the pointwise dimension of μ at x_0 exists and is bounded away from 0 and ∞ .

our result is valid for balls $B(x_0, \varepsilon)$ in the original Riemmanian metric and with a possibly singular measure; Note that the convergence of the extremal exponent θ_ε is not expected with singular measures.

Proof. Due to the proof of Theorem 2.2, we assume that $(m_\varepsilon)_\varepsilon$ converges to some m . (b) will follow from (a) by Theorem 5.2. The fact that (b) implies (c), which implies (d) comes from the definitions.

Let's prove (a) by applying Proposition 3.2. It is well known that our system satisfies Hypothesis 3.1-(II) for any $\alpha, \beta > 0$. Our assumption on $(m_\varepsilon)_\varepsilon$ ensures Assumption (iii) of Proposition 3.2.

Choose b_0, b such that $0 < b_0 < b < ad_u \log \lambda^{-1}$, with a given by Lemma 5.1. Let $p_\varepsilon = \varepsilon^{-\sigma}$ with $\sigma = b - b_0$. By Lemma 5.1

$$\mu(A_\varepsilon \cap \{\tau_{A_\varepsilon} \leq p_\varepsilon\}) \leq \sum_{n=\lfloor a \log 1/\varepsilon \rfloor + 1}^{p_\varepsilon} \mu(A_\varepsilon \cap T^{-n}A_\varepsilon).$$

By Lemma 5.4 (with $c = 1$) this sum is bounded by

$$p_\varepsilon \varepsilon^b \mu(B(x_0, 2\varepsilon)) = o(\mu(B(x_0, \varepsilon)))$$

by assumption. Hence assumption (i) of Proposition 3.2 holds by Lemma 5.6. (ii) comes from Lemma 5.7 and (iv) follows as in the proof of Theorem 3.5. \square

Lemma 5.4. *For any $a, b, c > 0$ such that $a \log \lambda + b/d_u + c < 1$, for ε small enough, for any $n \geq a \log 1/\varepsilon$ we have*

$$\mu(A_\varepsilon \cap T^{-n}A_\varepsilon) \leq \varepsilon^b \mu(B(x_0, \varepsilon + \varepsilon^c)).$$

Proof. Let $\kappa > 0$ small and consider a partition (or a cover with finite multiplicity) of Ω by pieces of unstable cubes (or balls) $W \in \mathcal{W}$ of size $\varepsilon^{1-\kappa}$. Let \mathcal{V} be the set of the $V = T^{-n}W$, $W \in \mathcal{W}$.

We can disintegrate the measure with respect to \mathcal{V} such that for any set Z

$$\mu(Z) = \int_{\mathcal{V}} \mu(Z|V) d\mu(V). \quad (11)$$

Since each $W = T^n V$ is a small piece of a smooth manifold, its intersection with $B(x_0, \varepsilon)$ consists at most in an unstable ball of radius $C\varepsilon$. Therefore the proportion of the unstable volume of $W \cap B(x_0, \varepsilon)$ in W is bounded by $C\varepsilon^{\kappa d_u}$, where d_u is the unstable dimension. By distortion, we get that $\mu(T^{-n}B(x_0, \varepsilon)|V) \leq C\varepsilon^{\kappa d_u}$.

Using (11) we get

$$\begin{aligned} \mu(B(x_0, \varepsilon) \cap T^{-n}B(x_0, \varepsilon)) &= \int_{\mathcal{V}} \mu(B(x_0, \varepsilon) \cap T^{-n}B(x_0, \varepsilon)|V) d\mu(V) \\ &\leq \int_{\mathcal{V}} \mu(T^{-n}B(x_0, \varepsilon)|V) 1_{V \cap B(x_0, \varepsilon) \neq \emptyset} d\mu(V) \\ &\leq C\varepsilon^{\kappa d_u} \mu(B(x_0, \varepsilon + \varepsilon^c)), \end{aligned}$$

since $\text{diam } V \leq C\lambda^n \varepsilon^{1-\kappa} \leq \varepsilon^c$. \square

Lemma 5.5. *There exists a constant $\delta_0 > 0$ independent of ε such that*

$$\mu(B(x_0, \frac{7}{8}\varepsilon)) \leq (1 - \delta_0) \mu(B(x_0, \varepsilon)).$$

Proof. We fix a measurable partition \mathcal{V} of unstable manifolds such that the d_u -dimensional Lebesgue measure of V is bounded from below by a constant, and $x_0 \notin \overline{\cup_V \partial V}$, such that the disintegration (11) holds true.

Let $\rho = \frac{7}{8}$. Suppose that $V \in \mathcal{V}$ intersects the ball $B(x_0, \rho\varepsilon)$. Then V intersects also the sphere $S(x_0, \frac{\rho+1}{2}\varepsilon)$ in a point say x , and the ball $B(x, \frac{1-\rho}{2}\varepsilon) \cap V$ is contained in $B(x_0, \varepsilon) \setminus B(x_0, \rho\varepsilon)$. Since $\mu(\cdot|V)$ is equivalent to the d_u -dimensional Lebesgue measure, there exists $\delta_0 > 0$ such that

$$\mu(B(x_0, \varepsilon) \setminus B(x_0, \rho\varepsilon)|V) \geq \mu(B(x, \frac{1-\rho}{2}\varepsilon)|V) \geq \delta_0 \mu(B(x_0, \varepsilon)|V)$$

and the lemma follows by integration. \square

Lemma 5.6. *The extremal index is bounded away from zero: $\theta_\varepsilon \geq \frac{\delta_0}{q_0+1} > 0$.*

Proof. We first observe, using equation (17) that

$$B(x_0, \varepsilon) \cap \{\tau_{B(x_0, \varepsilon)} \circ T^{q_0} \leq q_0\} \subset T^{-q} B(x_0, \frac{7}{8}\varepsilon).$$

for some integer q . Denote for simplicity $B = B(x_0, \varepsilon)$. By Lemma 5.5 this gives

$$\mu(B \cap \{\tau_B \circ T^{q_0} > q_0\}) \geq \delta_0 \mu(B).$$

Recall that

$$\mu(A_\varepsilon) = \theta_\varepsilon \mu(B).$$

We have

$$B \cap \{\tau_B \circ T^{q_0} > q_0\} \subset \bigcup_{j=0}^{q_0} T^{-j} A_\varepsilon$$

hence

$$\mu(B \cap \{\tau_B \circ T^{q_0} > q_0\}) \leq (q_0 + 1) \theta_\varepsilon \mu(B).$$

This implies that $\delta_0 \leq (q_0 + 1) \theta_\varepsilon$. \square

Lemma 5.7. *Suppose that T is $C^{1+\alpha}$ and that $\alpha > 1 - 1/\bar{d}_\mu(x_0)$. Then*

$$\mu(B(x_0, \varepsilon) \setminus B(x_0, \varepsilon - \varepsilon^\delta)) = o(\mu(B(x_0, \varepsilon)))$$

for any δ such that $\bar{d}_\mu(x_0) < \frac{\delta+1}{2} < \frac{1}{1-\alpha}$.

Proof. We fix a measurable partition \mathcal{V} of unstable manifolds such that the d_u -dimensional Lebesgue measure of V is bounded from below by a constant, and $x_0 \notin \overline{\cup_V \partial V}$, such that the disintegration (11) holds true.

Up to applying the exponential map at x_0 we assume that $B(x_0, \varepsilon_0)$ is the ball $B(0, \varepsilon_0)$ of \mathbb{R}^d .

Let $\varepsilon \in (0, \varepsilon_0)$. Let $V \in \mathcal{V}$. Let $y \in V \cap B(0, \varepsilon) \setminus B(0, \varepsilon - \varepsilon^\delta)$. Up to a rotation we suppose that locally V is the graph of a $C^{1+\alpha}$ map φ from $U \subset \mathbb{R}^{d_u}$ to \mathbb{R}^{d-d_u} , with $d_x \varphi = 0$ where $x \in U$ is such that $y = (x, \varphi(x))$. We have

$$(\varepsilon - \varepsilon^\delta)^2 \leq |x|^2 + |\varphi(x)|^2 \leq \varepsilon^2. \quad (12)$$

Let $h \in \mathbb{R}^{d_u}$ such that $x+h \in U$ and $(x+h, \varphi(x+h)) \in B(0, \varepsilon) \setminus B(0, \varepsilon - \varepsilon^\delta)$.

We have

$$(\varepsilon - \varepsilon^\delta)^2 \leq |x+h|^2 + |\varphi(x+h)|^2 \leq \varepsilon^2. \quad (13)$$

Subtracting (12) to (13) we get, setting $a(h) = |\varphi(x+h)|^2 - |\varphi(x)|^2$, since $\varepsilon^{2\delta} \leq \varepsilon^{1+\delta}$,

$$3\varepsilon^{1+\delta} \geq ||x+h|^2 - |x|^2 + a(h)| = ||h|^2 + 2x \cdot h + a(h)|.$$

Assume that $|h| > 6\varepsilon^{\frac{1+\delta}{2}}$. We fix a unit vector $u \in \mathbb{R}^{d_u}$ and seek for the solutions $h = tu$ of the above equation, which becomes

$$|t^2 + 2x \cdot ut + a(tu)| \leq 3\varepsilon^{1+\delta}.$$

Therefore, dividing by $|t| > 6\varepsilon^{\frac{1+\delta}{2}}$ we end up with

$$\left| t + 2x \cdot u + \frac{a(tu)}{t} \right| \leq \frac{1}{2}\varepsilon^{\frac{1+\delta}{2}}.$$

Let t, t' be two solutions. We obtain by subtraction

$$\left| t' - t + \frac{a(t'u)}{t'} - \frac{a(tu)}{t} \right| \leq \varepsilon^{\frac{1+\delta}{2}}.$$

Note that the function $g_u(t) := \frac{a(tu)}{t}$ is C^1 in the range of t 's, and

$$g'_u(t) = \frac{1}{t}(2(d_{x+tu}\varphi u) \cdot \varphi(x+tu)) - \frac{a(tu)}{t^2}.$$

Since φ is $C^{1+\alpha}$ we have $|d_{x+tu}\varphi u| = O(|t|^\alpha)$. In addition,

$$a(tu) = (|\varphi(x+tu)| - |\varphi(x)|)(|\varphi(x+tu)| + |\varphi(x)|) = O(\varepsilon|t|^{1+\alpha}).$$

This implies that $|g'_u(t)| = O(\varepsilon|t|^{\alpha-1}) = O(\varepsilon^{1+(\alpha-1)\frac{1+\delta}{2}}) = o(1)$, hence

$$|(t-t')(1+o(1))| \leq \varepsilon^{\frac{1+\delta}{2}}.$$

Using radial integration this gives that the d_u -dimensional Lebesgue measure of those h such that $(x+h, \varphi(x+h)) \in B(0, \varepsilon) \setminus B(0, \varepsilon - \varepsilon^\delta)$ is $O(\varepsilon^{\frac{1+\delta}{2}})$. Hence $\mu(B(x_0, \varepsilon) \setminus B(x_0, \varepsilon - \varepsilon^\delta)|V) = O(\varepsilon^{\frac{1+\delta}{2}})$, and the result follows by integration using (11). \square

6. BILLIARD IN A DIAMOND


We consider a diamond shaped billiard, with no cusp. The billiard table Q is a bounded closed part of \mathbb{R}^2 delimited by 4 convex obstacles $(\Gamma_i)_{i \in \mathbb{Z}/4\mathbb{Z}}$ (with C^3 -smooth boundary, with positive curvature) placed in such a way that, for every $i \in \mathbb{Z}/4\mathbb{Z}$, $\partial\Gamma_i$ meets $\partial\Gamma_{i+1}$ transversely at some point called corner C_i , but has no common point with $\partial\Gamma_j$ for $j \neq i-1, i+1$. In our representation of this billiard table, $C_1 = (0, 0)$ is on the left side of Q and the inner bisector at the corner C_1 is horizontal. We consider again the billiard flow $(\mathcal{M}, \nu, (Y_t)_t)$ and the billiard map (Ω, μ, T) in the domain Q .

For any $\varepsilon > 0$, we put a virtual vertical barrier I_ε of length ε joining a point $a_\varepsilon^{(1)} \in \partial\Gamma_1$ to a point $a_\varepsilon^{(2)} \in \partial\Gamma_2$. and we are interested in the times at which the billiard flow enters the corner by crossing the barrier I_ε . So that we define

$$\mathcal{A}_\varepsilon := I_\varepsilon \times S_-^1, \quad \text{with} \quad S_-^1 := \{\vec{v} = (v_1, v_2) \in S^1 : v_1 < 0\}.$$

We take $V := \mathbb{R} \times S_-^1$ and

$$\mathcal{H}_\varepsilon : (q = (q_1, q_2), \vec{v}) \mapsto \left(\frac{q_2}{\varepsilon}, \vec{v} \right).$$



Theorem 6.1. *The point process*

$$\sum_{t>0: Y_t \in \mathcal{A}_\varepsilon} \delta\left(\frac{\varepsilon t}{\pi \text{Area}(Q)}, \mathcal{H}_\varepsilon(Y_t(\cdot))\right)$$

converges in distribution to a Poisson point process on $[-1/2, 1/2] \times S_-^1$ with density $\lambda \times m_0$, with m_0 the probability measure of density proportional to $(\bar{q}, \vec{v}) \mapsto |v_1|$ with $\vec{v} = (v_1, v_2)$.

6.1. Notations, recalls and proof of Theorem 6.1. Let us recall some useful facts and notations.

Due to the transversality of $\partial\Gamma_1$ and $\partial\Gamma_2$ at C_1 , there exist $0 < \theta_1 < \theta_2$ such that, for $\varepsilon > 0$ small enough, the distance on ∂Q between C_1 and $a_\varepsilon^{(i)}$ is between $\theta_1\varepsilon$ and $\theta_2\varepsilon$.

Here Ω is the set of reflected unit vectors based on $\partial Q \setminus \{C_1, \dots, C_4\}$. We parametrize Ω by $\bigcup_{i \in \mathbb{Z}/4\mathbb{Z}} \{i\} \times]0, \text{length}(\partial\Gamma_i \cap Q)[\times [-\frac{\pi}{2}; \frac{\pi}{2}]$. A reflected vector (q, \vec{v}) is represented by (i, r, φ) if $q \in \partial\Gamma_i$ at distance r (on $Q \cap \partial\Gamma_i$) of C_{i-1} and if φ is the angular measure in $[-\pi/2, \pi/2]$ of $(\vec{n}(q), \vec{v})$ where $\vec{n}(q)$ is the normal vector to ∂Q at q .

For any C^1 -curve γ in Ω , we write $\ell(\gamma)$ for the euclidean length in the (r, φ) coordinates of γ . If moreover γ is given in coordinates by $\varphi = \phi(r)$, then we also write $p(\gamma) := \int_\gamma \cos(\phi(r)) dr$. We define the time until the next reflection in the future by

$$\tau^+(q, \vec{v}) := \min\{s > 0 : q + s\vec{v} \in \partial Q\}.$$

We also define $\tau^- : \Omega \rightarrow (0, +\infty)$ for the time until the last reflection in the past (corresponding to $\tau^- = \tau^+ \circ T^{-1}$ when T^{-1} is well defined) by

$$\tau^-(q, \vec{v}) := \min\{s > 0 : q + s\vec{v}^- \in Q\},$$

with \vec{v}^- be the reflected vector with respect to the normal to ∂Q at q , i.e. \vec{v}^- is the unit vector satisfying the following angular equality $\angle(\vec{n}(q), \vec{v}) = \angle(\vec{v}^-, \vec{n}(q))$. It will be useful to define $R_0 := \{\varphi = \pm\pi/2\}$, $\mathcal{C}_+ = \{(q, \vec{v}) \in \Omega : q + \tau^+(q, \vec{v})\vec{v} \in \{C_1, \dots, C_4\}\}$ and $\mathcal{C}_- = \{(q, \vec{v}) \in \Omega : q + \tau^-(q, \vec{v})\vec{v}^- \in \{C_1, \dots, C_4\}\}$. Observe that, for every $k \geq 1$, T^k defines a C^1 -diffeomorphism from $\Omega \setminus \mathcal{S}_{-k}$ to $\Omega \setminus \mathcal{S}_k$ with $\mathcal{S}_{-k} := T^{-k}R_0 \cup \bigcup_{m=0}^{k-1} T^{-m}(R_0 \cup \mathcal{C}_+)$ and $\mathcal{S}_k := T^kR_0 \cup \bigcup_{m=0}^{k-1} T^m(R_0 \cup \mathcal{C}_-)$. As for the other billiard models, we set $\pi_Q : \Omega \rightarrow Q$ for the canonical projection.

Despite the absence of the so called complexity bound in billiards with corners, De Simoi and Toth have shown in [5] that some expansion condition

holds, from which the growth lemma [4, Theorem 5.52] follows. It says that for any weakly homogeneous unstable curve W one has

$$m_W(r_n < \delta) \leq c\theta^n \delta + c\delta m_W(W) \quad (14)$$

where m_W is the one dimensional Lebesgue measure on W , and $r_n(x)$ denotes the distance (on $T^n W$) of $T^n(x)$ to the boundary of the homogeneous piece of $T^n W$ containing x .

In particular, for those systems one can build a Young tower with exponential parameters, from which it follows that Hypothesis 3.1-(II) is satisfied for every $\alpha, \beta > 0$.

Proof of Theorem 6.1. Let $A_\varepsilon \subset \Omega$ be the set of all possible configurations at the reflection time just after the particle crosses the virtual barrier I_ε from the right side. Note that $\mu(A_\varepsilon) = \frac{1}{2|\partial Q|} \int_{[-\varepsilon/2, \varepsilon/2] \times [-\pi/2, \pi/2]} \cos \varphi \, dr \, d\varphi = \frac{\varepsilon}{|\partial Q|}$.

Due to Theorem 2.3, it is enough to prove that $(\mathcal{N}_\varepsilon)_\varepsilon$ converges to a Poisson point process with density $\lambda \times m_0$ for the above choice of A_ε and for $H_\varepsilon : A_\varepsilon \rightarrow \mathbb{R} \times S^1_-$, such that given by $\mathcal{H}_\varepsilon = H_\varepsilon \circ T \circ \Pi$ with Π the projection defined in Subsection 2.2. To this end we apply Proposition 3.2.

The fact that Assumption (i) of Proposition 3.2 is satisfied for $p_\varepsilon = \varepsilon^{-\sigma}$ comes from Proposition 6.2. We take $\alpha > 3/\sigma$. Assumption (ii) comes from the fact that the boundary of each connected component of ∂A_ε is made of a part of R_0 and of a C^1 -increasing curve $r = R(\varphi)$ with $R'(\varphi) = 1/(\kappa(r) + \frac{\cos(\varphi)}{\tau^-(r, \varphi)}) \leq 1/\min \kappa$ corresponding to reflected vectors in the corner on the leftside of I_ε coming from $\{a_\varepsilon^{(1)}, a_\varepsilon^{(2)}\}$ and to $T(R_0 \cap A_\varepsilon)$. The image measure of $\mu(\cdot|A_\varepsilon)$ by H_ε is proportional to $\cos(\varphi) dr d\varphi$ where r is the position on $[-1/2, 1/2]$ and φ the angle (in $[-\pi/2, \pi/2]$) between the vector $(-1, 0)$ and the incident vector (i.e. the speed vector at the time when the particle crosses I_ε). We take for \mathcal{W} the set of rectangles of the form $(a, b) \times (c, d)$ in the above (r, φ) coordinates. Outside the strips $A_\varepsilon \cap \{(r, \varphi) \mid |r - a_\varepsilon^{(i)}| < \varepsilon^2\}$, H_ε is $K\varepsilon^{-2}$ -Lipschitz (indeed the jacobian is in $O(\varepsilon^{-1}/\cos \varphi) \leq c\varepsilon^{-2}$) and so, using the argument of the proof of Proposition 3.4-(ii), we conclude that (iv) is satisfied since $\mu(A_\varepsilon) = O(\varepsilon)$. \square

6.2. Short returns. The aim of this subsection is the following result.

Proposition 6.2. *There exists $\sigma > 0$ such that $\mu(\tau_{A_\varepsilon} < \varepsilon^{-\sigma} | A_\varepsilon) = o(1)$.*

To this end, we will recall useful facts and introduce some notations. Let $\tau_0 := \min_{(i,j)} \min_{j \neq i, i+1} \text{dist}(C_i, \Gamma_j)/10$.

Definition 6.3. *We say that a curve γ of Ω satisfies assumption (C) if it is given by $\varphi = \phi(r)$ with ϕ C^1 -smooth, increasing and such that $\min \kappa \leq \phi' \leq \max \kappa + \frac{1}{\tau_0}$.*

We recall the following facts.

- There exist $C_0, C_1 > 0$ and $\lambda_1 > 1$ such that, for every γ satisfying Assumption (C) and every integer m such that $\gamma \cap \mathcal{S}_{-m} = \emptyset$, $T^m \gamma$ is a C^1 -smooth curve satisfying $C_1 p(T^m \gamma) \geq \lambda_1^m p(\gamma)$ and $\ell(\gamma) \leq C_0 \sqrt{p(T\gamma)}$.

- There exist $C_2 > 0$ and $\lambda_2 > \lambda_1^{1/2}$ such that, for every integer m , the number of connected components of $\Omega \setminus \mathcal{S}_{-m}$ is less than $C_2 \lambda_2^m$. Moreover \mathcal{S}_{-m} is made of curves $\varphi = \phi(r)$ with ϕ C^1 -smooth and strictly decreasing.
- If $\gamma \subset \Omega \setminus \mathcal{S}_{-1}$ is given by $\varphi = \phi(r)$ or $r = \mathfrak{r}(\varphi)$ with ϕ or \mathfrak{r} increasing and C^1 smooth, then $T\gamma$ is C^1 , is given by $\varphi = \phi_1(r)$ with $\min \kappa \leq \phi'_1 \leq \max \kappa + \frac{1}{\min_\gamma \tau^+}$. Moreover $\int_{T\gamma} d\varphi \geq \int_\gamma d\varphi$.
- There exists m_0 such that, for every $x \in \Omega \setminus \bigcup_{k=0}^{m_0-1} T^{-k} \mathcal{C}_+$, there exists $k \in \{1, \dots, m_0\}$ such that $\tau^+(T^{k-1}) > \tau_0$.

Let $A_\varepsilon \subset \Omega$ be the set of possible configurations of a particle at the reflection time just after the particle reaches the virtual barrier I_ε from the right side. We observe that there exists $K_0 > 0$ and $\varepsilon_0 > 0$ such that, for every $\varepsilon \in (0, \varepsilon_0)$, for every q between C_1 and $a_\varepsilon^{(i)}$, the set of \vec{v} such that $(q, \vec{v}) \in A_\varepsilon$ has Lebesgue measure at least K_0 .

Lemma 6.4 (Very quick returns). *There exists $K_1 > 0$ such that,*

$$\forall s \geq 1, \quad \mu(T^{-s}(A_\varepsilon)|A_\varepsilon) \leq K_1(\lambda_2/\lambda_1^{\frac{1}{2}})^s \varepsilon^{\frac{1}{2}}.$$

Proof. Let $q \in \pi_Q(A_\varepsilon)$. Let γ_1 be a connected component of $\pi_Q^{-1}(\{q\}) \setminus \mathcal{S}_s$. We define $\gamma := \gamma_1 \cap A_\varepsilon \cap T^{-s}A_\varepsilon$. Let m be the smallest positive integer such that $\min_\gamma \tau \circ T^{m-1} > \tau_0$. By definition of A_ε and of m_0 , $m < \min(m_0, s)$. Hence $T^m\gamma$ satisfies Assumption (C) and

$$\ell(\gamma) \leq \int_\gamma d\varphi \leq \int_{T^m\gamma} d\varphi \leq \ell(T^m\gamma) \leq C_0 \sqrt{p(T^m\gamma)}.$$

Moreover, since $\gamma \cap S_{-s} = \emptyset$, we also have

$$p(T^m\gamma) \leq C_1 \lambda_1^{m-s} p(T^s\gamma).$$

But, since $T^s\gamma$ is an increasing curve contained in A_ε , we conclude that $p(T^s\gamma) \leq \theta\varepsilon$. Hence

$$\ell(\gamma) \leq C_0 \sqrt{C_1 \lambda_1^{m-s} \theta \varepsilon}.$$

By using the fact that $\pi_Q^{-1}(\{q\}) \setminus \mathcal{S}_s$ contains at most $C_2 \lambda_2^s$ connected components and by integrating on $\pi_Q(A_\varepsilon)$, we obtain

$$\mu(T^{-s}(A_\varepsilon) \cap A_\varepsilon) \leq \frac{C_0 \sqrt{\theta C_1} C_2 \lambda_1^{\frac{m}{2}} (\lambda_2/\lambda_1^{\frac{1}{2}})^s \varepsilon^{\frac{1}{2}}}{2 \text{length}(\partial Q)} \text{length}(\pi_Q(A_\varepsilon)).$$

We conclude by using the fact that $\mu(A_\varepsilon) \geq (1 - \sin K_0) \text{length}(\pi_Q(A_\varepsilon))$ and by setting $K_1 := \frac{C_0 \sqrt{\theta C_1} C_2 \lambda_1^{\frac{m_0}{2}}}{1 - \sin K_0}$. \square

Lemma 6.5 (quick returns). *For any $a > 0$, there exists $s_a > 0$ such that*

$$\sum_{n=-a \log \varepsilon}^{\varepsilon^{-s_a}} \mu(A_\varepsilon \cap T^{-n} A_\varepsilon) = o(\mu(A_\varepsilon)).$$

Proof. We take a measurable partition \mathcal{V} of $\Omega \setminus \mathcal{S}_1$ by the unstable curves $\frac{d\varphi}{dr} = \kappa(r)$. By disintegration there exist a probability measure $\tilde{\mu}$ on \mathcal{V} , and a constant $c < \infty$ such that for any measurable set B we have

$$c^{-1}\mu(B) \leq \int_{\mathcal{V}} m_W(B) d\tilde{\mu}(W) =: \mu_0(B).$$

We define \tilde{A}_ε as the set of $T^{-j}x$ where $x \in A_\varepsilon$ and j is the minimal integer such that $T^{-\ell}x$ does not belong to the sides adjacent to the corner. Any corner sequence is bounded by some constant m depending only on the billiard table, thus $A_\varepsilon \subset \cup_{\ell=1}^m T^\ell \tilde{A}_\varepsilon$. It follows by invariance that

$$\mu(A_\varepsilon \cap T^{-n}A_\varepsilon) \leq m^2 \max_{|\ell| \leq m} \mu(\tilde{A}_\varepsilon \cap T^{-n-\ell} \tilde{A}_\varepsilon).$$

Therefore it suffices to control $\mu(\tilde{A}_\varepsilon \cap T^{-n} \tilde{A}_\varepsilon)$.

Note that \tilde{A}_ε is at a long distance from the corner, hence there are finitely many decreasing curves φ_j , $j = 1..m_0$ and a constant c such that $\tilde{A}_\varepsilon \subset \tilde{V}_\varepsilon$ where

$$\tilde{V}_\varepsilon = \cup_j \{(r, \varphi) : \varphi_j(r) - c\varepsilon < \varphi < \varphi_j(r) + c\varepsilon\}$$

where $(r, \varphi_j(r))$ represents a vector pointing exactly to the corner provided its two adjacent obstacles are removed. In particular $\frac{d\varphi_j(r)}{dr} \leq -\kappa$. We denote the k th homogeneity strip⁷ by \mathbb{H}_k for $k \neq 0$ and set $\mathbb{H}_0 = \cup_{|k| < k_0} \mathbb{H}_k$ for some fixed k_0 . Set $s := \min(-a \log \theta, 1)/3$. Let $k_\varepsilon = \varepsilon^{-s}$ and $H^\varepsilon = \cup_{|k| \leq k_\varepsilon} \mathbb{H}_k$. For any $W \in \mathcal{V}$ we set $W^\varepsilon = W \cap \tilde{V}_\varepsilon$, $W_k = W \cap \mathbb{H}_k$ and $W_k^\varepsilon = W_k \cap \tilde{V}_\varepsilon$. Each W_k^ε is a weakly homogeneous unstable curve.

We cut each curve W_k^ε into small pieces $W_{k,i}^\varepsilon$ such that each $T^j W_{k,i}^\varepsilon$, $j = 0, \dots, n$ is contained in a homogeneity strip and a connected component of $\Omega \setminus \mathcal{S}_1$. For $x \in W_{k,i}^\varepsilon$ we denote by $r_n(x)$ the distance (in $T^n W$) of $T^n(x)$ to the boundary of $T^n W_{k,i}^\varepsilon$.

By definition of W^ε

$$\begin{aligned} m_W(\tilde{A}_\varepsilon \cap T^{-n} \tilde{A}_\varepsilon) &\leq m_{W^\varepsilon}(T^{-n} \tilde{V}_\varepsilon) \\ &\leq m_{W^\varepsilon}(\mathbb{H}_\varepsilon^c) + \sum_{|k| \leq k_\varepsilon} m_{W_k^\varepsilon}(\{r_n \geq \varepsilon^{1-s}\} \cap T^{-n} \tilde{V}_\varepsilon) + m_{W_k^\varepsilon}(r_n < \varepsilon^{1-s}). \end{aligned}$$

The first term inside the sum is bounded by the sum $\sum_i m_{W_{k,i}^\varepsilon}(T^{-n} \tilde{V}_\varepsilon)$ over those i 's such that $T^n W_{k,i}^\varepsilon$ is of size larger than ε^{1-s} . In particular $m_{T^n W_{k,i}^\varepsilon}(T^n W_{k,i}^\varepsilon) \geq \varepsilon^{1-s}$. On the other hand, by transversality

$$m_{T^n W_{k,i}^\varepsilon}(\tilde{V}_\varepsilon) \leq c\varepsilon.$$

By distortion (See Lemma 5.27 in [4]) we obtain

$$m_{W_{k,i}^\varepsilon}(T^{-n} \tilde{V}_\varepsilon) \leq c\varepsilon^s m_{W_{k,i}^\varepsilon}(W_{k,i}^\varepsilon).$$

Summing up over these i gives the first term inside the sum is bounded by

$$m_{W_k^\varepsilon}(\{r_n \geq \varepsilon^{1-s}\} \cap T^{-n} \tilde{V}_\varepsilon) \leq c\varepsilon^s m_{W_k^\varepsilon}(W_k^\varepsilon).$$

On the other hand, the growth lemma (14) implies that

$$m_{W_k^\varepsilon}(r_n < \varepsilon^{1-s}) \leq c\theta^n \varepsilon^{1-s} + c\varepsilon^{1-s} m_{W_k^\varepsilon}(W_k^\varepsilon).$$

⁷See [4] for notations and definitions.

A final summation over k gives

$$m_W(\tilde{A}_\varepsilon \cap T^{-n} \tilde{A}_\varepsilon) \leq m_W(\tilde{V}_\varepsilon \cap \mathbb{H}_\varepsilon^c) + c(\varepsilon^s + \varepsilon^{1-s})m_W(\tilde{V}_\varepsilon) + ck_\varepsilon \theta^n \varepsilon^{1-s}.$$

Integrating over $W \in \mathcal{V}$ gives

$$\mu(\tilde{A}_\varepsilon \cap T^{-n} \tilde{A}_\varepsilon) \leq \mu_0(\tilde{V}_\varepsilon \cap \mathbb{H}_\varepsilon^c) + O(\varepsilon^{1+s/3}) = O(\mu(A_\varepsilon) \varepsilon^{s/3}),$$

where we used the fact that μ_0 is equivalent to Lebesgue and $\tilde{V}_\varepsilon \cap \mathbb{H}_\varepsilon^c$ is contained in the union of at most m_0 rectangles of width $O(\varepsilon)$ and height $k_\varepsilon^{-2} = \varepsilon^{2s}$. We take $s_a = s/6$. \square

Proof of Proposition 6.2. Choose $a = 1/(4 \log(\lambda_2/\lambda_1^{1/2}))$. Observe that, due to Lemma 6.4, we have

$$\sum_{s=1}^{-a \log \varepsilon} \mu(T^{-s} A_\varepsilon | A_\varepsilon) \leq \frac{K_1}{\lambda_2/\lambda_1^{1/2} - 1} (\lambda_2/\lambda_1^{1/2})^{-a \log \varepsilon} \varepsilon^{1/2} \leq \frac{K_1}{\lambda_2/\lambda_1^{1/2} - 1} \varepsilon^{1/4}.$$

This combined with Lemma 6.5 leads to

$$\sum_{n=1}^{\varepsilon^{-sa}} \mu(T^{-n} A_\varepsilon | A_\varepsilon) = o(1).$$

\square

APPENDIX A. MORE PROOFS

Proof of Proposition 3.2. Item (ii) of Theorem 2.1 being satisfied by assumption by \mathcal{W} , it remains to prove Item (i).

Let $\mathcal{G}_0 = \{G_1, \dots, G_L\}$ be a finite subcollection of $H_\varepsilon^{-1}\mathcal{W}$. Let $t > 0$. Let $A \in \sigma(\mathcal{G}_0)$ and $B \in \sigma(\bigcup_{n=1}^{N_{\varepsilon,t}} T^{-n} \mathcal{G}_0)$, with $N_{\varepsilon,t} := \lfloor t/\mu(A_\varepsilon) \rfloor$.

Set $X_j := (1_{T^{-j}(G_1)}, \dots, 1_{T^{-j}(G_L)}) \in \mathbb{R}^L$. Note that $\mathbf{1}_B = g(X_1, \dots, X_{N_{\varepsilon,t}})$ for some $g: (\{0, 1\}^L)^{N_{\varepsilon,t}} \rightarrow \{0, 1\}$. Note that if ε is small enough, $N_{\varepsilon,t} > p_\varepsilon$ by assumption.

Let $B_1 = \{g(0, \dots, 0, X_1, \dots, X_{N_{\varepsilon,t}-p_\varepsilon}) = 1\}$ so that $|\mathbf{1}_B - \mathbf{1}_{B_1} \circ f^{p_\varepsilon}| \leq \mathbf{1}_{\{\tau_{A_\varepsilon} \leq p_\varepsilon\}}$. Note that

$$|\mu(B \cap A | A_\varepsilon) - \mu(B_1 \cap A | A_\varepsilon)| \leq \mu(\tau_{A_\varepsilon} \leq p_\varepsilon | A_\varepsilon) = o(1).$$

Moreover

$$|\mu(B) - \mu(B_1)| \leq \mu(\tau_{A_\varepsilon} \leq p_\varepsilon) \leq p_\varepsilon \mu(A_\varepsilon) = o(1).$$

Set $K := \lfloor p_\varepsilon/4 \rfloor$.

Under (I), set $M = 0$, $\mathcal{P}_K = \mathcal{Q}_K$.

Under (II), set $M = K$, $\mathcal{P}_K = \mathcal{Q}_{2K}$.

It remains to show that

$$|\mu(B_1 \cap A) - \mu(B_1)\mu(A)| = o(\mu(A_\varepsilon)), \quad (15)$$

i.e.

$$|\text{Cov}_{\tilde{\mu}}(\mathbf{1}_{\tilde{T}^{-M}, \tilde{\Pi}^{-1}B_1}, \mathbf{1}_{\tilde{T}^{-M}, \tilde{\Pi}^{-1}A})| = o(\tilde{\mu}(\tilde{T}^{-M} \tilde{\Pi}^{-1} A_\varepsilon)).$$

We approximate $\tilde{T}^{-M}\tilde{\Pi}^{-1}A$ (resp. $\tilde{T}^{-M}\tilde{\Pi}^{-1}A_\varepsilon$) by the union of the atoms of \mathcal{P}_K intersecting this set. We write \tilde{A} (resp. \tilde{A}_ε) for this union. Note that $\tilde{T}^{-M}\tilde{\Pi}^{-1}A \subset \tilde{A}$ (resp. $\tilde{T}^{-M}\tilde{\Pi}^{-1}A_\varepsilon \subset \tilde{A}_\varepsilon$) and that

$$\tilde{A} \setminus \tilde{T}^{-M}\tilde{\Pi}^{-1}A \subset A' := \bigcup_{Q \in Q_K: \tilde{\Pi}\tilde{T}^M Q \cap A \neq \emptyset, \tilde{\Pi}\tilde{T}^M Q \setminus B_1 \neq \emptyset} Q$$

and so (due to the assumption on the diameters of the atoms of \mathcal{P}_K),

$$\begin{aligned} \tilde{\mu}(A') &\leq \tilde{\mu} \left(\bigcup_{Q \in Q_K: \tilde{\Pi}\tilde{T}^M Q \subset (\partial A)^{[CK^{-\alpha}]}} Q \right) \\ &\leq \mu((\partial A)^{[CK^{-\alpha}]}) = o(\mu(A_\varepsilon)). \end{aligned}$$

Analogously

$$\tilde{A}_\varepsilon \setminus \tilde{T}^{-M}\tilde{\Pi}^{-1}A_\varepsilon \subset A'_\varepsilon := \bigcup_{Q \in Q_K: \tilde{\Pi}\tilde{T}^M Q \cap A_\varepsilon \neq \emptyset, \tilde{\Pi}\tilde{T}^M Q \setminus A_\varepsilon \neq \emptyset} Q$$

and

$$\tilde{\mu}(A'_\varepsilon) = o(\mu(A_\varepsilon)).$$

Finally we approximate $\tilde{T}^{-M}\tilde{\Pi}^{-1}B_1$ by a union \tilde{B} of atoms of $\sigma(\bigcup_{\ell=1}^{N_{\varepsilon,t}-p_\varepsilon} \mathcal{P}_K)$ such that $\tilde{T}^{-M}\tilde{\Pi}^{-1}B_1 \subset \tilde{B}_1$ and

$$\tilde{B}_1 \setminus \tilde{T}^{-M}\tilde{\Pi}^{-1}B_1 \subset B'_1 := \bigcup_{\ell=1}^{N_{\varepsilon,t}-p_\varepsilon} F^{-\ell} \left(\bigcup_{Q \in \mathcal{P}_K: \tilde{\Pi}\tilde{T}^M Q \subset \bigcup_{k=1}^L (\partial G_k)^{[CK^{-\alpha}]}} Q \right),$$

and

$$\tilde{\mu}(B'_1) = (N_{\varepsilon,t} - p_\varepsilon) o(\mu(A_\varepsilon)).$$

Therefore

$$\begin{aligned} &|Cov_{\tilde{\mu}}(\mathbf{1}_{\tilde{\Pi}^{-1}A}, \mathbf{1}_{\tilde{\Pi}^{-1}B_1}) - Cov_{\tilde{\mu}}(\mathbf{1}_{\tilde{A}}, \mathbf{1}_{\tilde{B}_1})| \\ &\leq |Cov_{\tilde{\mu}}(\mathbf{1}_{\tilde{A}} - \mathbf{1}_{\tilde{T}^{-M}\tilde{\Pi}A}, \mathbf{1}_{\tilde{B}_1}) + Cov_{\tilde{\mu}}(\mathbf{1}_{\tilde{T}^{-M}\tilde{\Pi}A}, \mathbf{1}_{\tilde{B}_1} - \mathbf{1}_{\tilde{T}^{-M}\tilde{\Pi}B_1})| \\ &\leq |Cov_{\tilde{\mu}}(\mathbf{1}_{A'}, \mathbf{1}_{\tilde{B}})| + |Cov_{\tilde{\mu}}(\mathbf{1}_{\tilde{A}}, \mathbf{1}_{B'_1})| \\ &\quad + 2\tilde{\mu}(A')\tilde{\mu}(\tilde{B}_1) + 2\tilde{\mu}(\tilde{A})\tilde{\mu}(B'_1) \\ &\leq C'_0(\mu(A_\varepsilon)p^{-\beta} + o(\mu(A_\varepsilon))(N - p)\mu(A_\varepsilon)) = o(\mu(A_\varepsilon)). \end{aligned}$$

□

Proof of Lemma 5.1. Since the analysis is local, taking the exponential map at x_0 if necessary we may assume that $U \subset \mathbb{R}^d$. Let L_0 be the Lipschitz norm of T . We first observe that if $a > 0$ is sufficiently small, then for any $\varepsilon > 0$ small and $n \leq a \log 1/\varepsilon$, $\|x - x_0\| < \varepsilon$ and $\|T^n x - x_0\| < \varepsilon$ implies that

$$\|T^n x_0 - x_0\| \leq \|T^n x - T^n x_0\| + \|T^n x - x_0\| \leq L_0^n \varepsilon + \varepsilon,$$

and thus n is a multiple of p . Hence without loss of generality we assume that $p = 1$.

Let q be the integer given by Lemma A.1 for the hyperbolic matrix $D_{x_0}T$.

Since the $C^{1+\alpha}$ norm of T^n is growing at most exponentially fast, changing the value of $a > 0$ if necessary, there exists $c, L \geq L_0$ such that for any $x \in B(x_0, \varepsilon)$,

$$\|T^n x - T^n x_0 - D_{x_0} T^n(x - x_0)\| \leq cL^n \varepsilon^{1+\alpha} \leq \varepsilon/2, \quad (16)$$

for any $1 \leq n \leq a \log 1/\varepsilon$.

Let $2q \leq n \leq a \log(1/\varepsilon)$ and suppose that $x \in B(x_0, \varepsilon) \cap T^{-n}B(x_0, \varepsilon)$. Using (16) we obtain $\|D_{x_0} T^n(x - x_0)\| \leq \|T^n x - x_0\| + \varepsilon/2 < \frac{3}{2}\varepsilon$. Thus Lemma A.1 gives $\|D_{x_0} T^q(x - x_0)\| \leq \frac{3}{8}\varepsilon$. Using again (16) we get

$$\|T^q x - T^q x_0\| \leq \|D_{x_0} T^q(x - x_0)\| + \varepsilon/2 < \frac{7}{8}\varepsilon.$$

Hence

$$B(x_0, \varepsilon) \cap T^{-n}B(x_0, \varepsilon) \subset T^{-q}B(x_0, \frac{7}{8}\varepsilon). \quad (17)$$

Set $q_0 = 2q$. When $n \geq q_0$ this proves that

$$A_\varepsilon \cap T^{-n}A_\varepsilon \subset A_\varepsilon \cap T^{-q}B(x_0, \varepsilon) = \emptyset,$$

while when $n < q_0$ the intersection is empty by definition. \square

Lemma A.1. *Let A be hyperbolic matrix. There exists an integer q such that for any vector v , any $n \geq 2q$, $\|A^q v\| \leq \max(\|v\|, \|A^n v\|)/4$.*

We leave the proof to the reader.

REFERENCES

- [1] J. F. Alves and D. Azevedo, Statistical properties of diffeomorphisms with weak invariant manifolds. Preprint ArXiv:1310.2754.
- [2] M. Carney, M. Nicol and H.K. Zhang, Compound Poisson law for hitting times to periodic orbits in two-dimensional hyperbolic systems. J. Stat. Phys. **169**-4 (2017) 804–823.
- [3] J.-R. Chazottes and P. Collet Poisson approximation for the number of visits to balls in nonuniformly hyperbolic dynamical systems, Erg. Th. Dynam. Sys. **33**-1 (2013) 49–80.
- [4] N. Chernov and R. Markarian, Chaotic billiards. Mathematical Surveys and Monographs, 127. American Mathematical Society, Providence, RI, (2006) xii+316 pp.
- [5] J. De Simoi, I.P. Toth, An Expansion Estimate for Dispersing Planar Billiards with Corner Points, Ann. H. Poincaré **15** (2014) 1223–1243
- [6] W. Doeblin, Remarques sur la thorie mtrique des fractions continues (French), Compositio Math. **7** (1940) 353–371.
- [7] A. C. Moreira Freitas, J. M. Freitas and M. Magalhães, Complete convergence and records for dynamically generated stochastic processes, Preprint arXiv:1707.07071.
- [8] J. Freitas, A. Freitas and M. Todd, The compound Poisson limit ruling periodic extreme behaviour of non-uniformly hyperbolic dynamics, Communications in Mathematical Physics **321** (2013) 483–527
- [9] N. Haydn and M. Todd, Ergodic Theory Dynam. Systems **36**-5 (2016) 1616–1643
- [10] N. Haydn, S. Vaienti, The compound Poisson distribution and return times in dynamical systems, Probability Theory and related fields **144**-3/4 (2009) 517–542
- [11] N. haydn and K. Wasilewska, Limiting distribution and error terms for the number of visits to balls in non-uniformly hyperbolic dynamical systems, Preprint arXiv:1402.2990
- [12] A. Katok and B. Hasselblatt, Introduction to the modern theory of dynamical systems, Cambridge Univ. Press, 1995.

- [13] V. Lucarini *et al*, Extremes and recurrence in dynamical systems. Pure and Applied Mathematics (Hoboken). John Wiley & Sons, Inc., Hoboken, NJ, 2016. xi+295 pp.
- [14] I. Melbourne and R. Zweimüller, Weak convergence to stable Lvy processes for nonuniformly hyperbolic dynamical systems, Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, **51**-2 (2015) 545–556.
- [15] F. Pène and B. Saussol, Back to balls in billiards, Comm. Math. Phys. **293**-3 (2010) 837–866
- [16] F. Pène and B. Saussol, Poisson law for some non-uniformly hyperbolic dynamical systems with polynomial rate of mixing. Ergodic Theory Dynam. Systems **36**-8 (2016) 2602–2626.
- [17] B. Pitskel, Poisson limit theorem for Markov chains, Ergodic Theory & Dynam. Systems **11**-3 (1001) 501–513.
- [18] S. Resnick, Extreme values, regular variation, and point processes, Springer Series in Operation research and financial engineering, Springer 2008.
- [19] B. Saussol, S. Troubetzkoy and S. Vaienti, Recurrence, dimensions and Lyapunov exponents, Journal of Statistical Physics **106** (2002) 623–634
- [20] M. Tyran-Kamińska, Weak convergence to Lévy stable processes in dynamical systems, Stochastics and Dynamics **10**(02) (2010) 263–289.
- [21] L.-S. Young, Statistical properties of dynamical systems with some hyperbolicity, Ann. of Math. **147** (1998) 585–650.
- [22] L.-S. Young, Recurrence times and rates of mixing, Israel J. Math. **110** (1999) 153–188.
- [23] R. Zweimüller, Mixing limit theorems for ergodic transformations, Journal of Theoretical Probability, **20** (2007), 1059–1071.

1)UNIVERSITÉ DE BREST, LABORATOIRE DE MATHÉMATIQUES DE BRETAGNE ATLANTIQUE, CNRS UMR 6205, BREST, FRANCE, 2)FRANÇOISE PÈNE IS SUPPORTED BY THE IUF.

E-mail address: francoise.pene@univ-brest.fr

E-mail address: benoit.saussol@univ-brest.fr