



Predicting the energy budget of the scallop *Argopecten purpuratus* in an oxygen-limiting environment

Arturo Aguirre-Velarde, Laure Pecquerie, Fred Jean, Gérard Thouzeau,
Jonathan Flye-Sainte-Marie

► To cite this version:

Arturo Aguirre-Velarde, Laure Pecquerie, Fred Jean, Gérard Thouzeau, Jonathan Flye-Sainte-Marie.
Predicting the energy budget of the scallop *Argopecten purpuratus* in an oxygen-limiting environment.
Journal of Sea Research (JSR), 2019, 143, pp.254-261. 10.1016/j.seares.2018.09.011 . hal-02114544

HAL Id: hal-02114544

<https://hal.univ-brest.fr/hal-02114544>

Submitted on 29 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Predicting the energy budget of the scallop *Argopecten purpuratus* in an oxygen-limiting environment

Arturo Aguirre-Velarde^{a,b,*}, Laure Pecquerie^b, Frédéric Jean^b, Gérard Thouzeau^b, Jonathan Flye-Sainte-Marie^b

^aLaboratorio de Ecofisiología Acuática, IMARPE, Esquina Gamarra y General Valle S/N Chucuito Callao, Peru

^bLEMAR, Université Bretagne Occidentale, CNRS, IRD, Ifremer, Plouzané, France

Abstract

1 Low concentrations of oxygen determine marine species distribution and abundance along the Peruvian
2 coast with consequences for human activity such as fishing and aquaculture. In order to assess bioener-
3 getic consequences of oxygen limitation on the Peruvian scallop *Argopecten purpuratus*, we first devel-
4 oped a Dynamic Energy Budget (DEB) model of growth and reproduction calibrated on field experimen-
5 tal datasets. Then, we included oxygen availability as an additional forcing variable using a simple rule
6 based on the ability of the scallop to regulate oxygen consumption. The resulting model was tested using
7 growth/reproduction monitoring and simultaneous high frequency environmental records in Paracas Bay
8 (Peru) at two different depths: suspended in the water column and on the sea bottom. Simulations indicated
9 that scallops growth and reproduction was not food-limited. The negative observed effects of hypoxia on
10 growth and reproduction could be explained by a decrease in both assimilation and reserve mobilization.
11 However, hypoxic conditions in summer were not sufficient to explain the observed losses in somatic tissue
12 weights and the disruption of reproduction. The latter two patterns were better simulated when assuming in-
13 creased somatic maintenance costs due to the presence of H₂S during milky turquoise water discolouration
14 events observed during summer.

Keywords:

DEB theory, bioenergetics, hypoxia, hydrogen sulfide, growth, reproduction, Peru

*Email: aaguirre@imarpe.gob.pe

1. Introduction

The abundant primary production in Peruvian coasts sustains a large biomass of primary and secondary consumers (e.g. small fish and bivalves). Nevertheless, as recently emphasized by Breitburg et al. (2018), these highly productive areas are the place of a paradox: they support some of the world's most prolific fisheries but are also associated with very strong oxygen-minimum zones (OMZ). Degradation processes of the large amount of settled organic matter consume the oxygen in deep waters and on the bottom (Gewin, 2010) generating a vast OMZ between 50 and 1000 m depth (Helly and Levin, 2004). Organic matter degradation in the absence of oxygen also produces toxic hydrogen sulfide (H_2S). Such giant H_2S plume has been reported off Peru (Schunck et al., 2013). However, OMZ impacts on littoral areas remains poorly known. High frequency environmental monitoring in Peruvian littoral bays recently revealed the frequent occurrence of strong oxygen limitations in very shallow areas (Aguirre-Velarde et al., 2016). These phenomena were also sporadically associated with the presence of milky, presumably sulfide-rich, waters. The impact of such events on resources and economic activities in these littoral bays (artisanal fisheries and aquaculture) is still poorly understood such and requires improving knowledge for an ecosystem-based sustainable management.

Unlike vagile organisms such as fishes that are, to some extent, able to avoid unfavorable conditions, sessile or semi-vagile organisms such as bivalves have to deal with local environmental constraints such as oxygen limitation. Although they have developed adaptations to cope with such events (Abele-Oeschger and Oeschger, 1995; Diaz and Rosenberg, 1995; Levin, 2003), negative effects of hypoxia/anoxia on metabolism, physiology, growth, reproduction and survival have been described (Herreid, 1980; Grieshaber et al., 1994; Burnett and Stickle, 2001; Levin et al., 2009). A decrease of food consumption, increase in ventilation and heart beat rates, and the production of potentially toxic metabolites due to the activation of anaerobic pathways, may all affect the energetic budget of organisms subjected to oxygen-limiting conditions. The peruvian scallop is found on seabeds frequently affected by hypoxic/anoxic events and exhibits an important ability to cope with such events including to regulate its respiration and feeding rates (Aguirre-Velarde et al., 2016, 2018). Although low effects of daily cycles of hypoxia on *A. purpuratus* energy budget were found by Aguirre-Velarde et al. (2018), stronger effects of hypoxic/anoxic events, combined with the occurrence of milky waters were observed in the field (Paracas bay, Peru; Aguirre-Velarde, 2016).

Oxygen limitations naturally occur in Peru, but the frequency of hypoxic events globally tends to increase as a result of industrialization and climate change (Levin et al., 2009; Rabalais et al., 2010). Declining

oxygen in global ocean and coastal waters is considered as one of the most important changes occurring in marine ecosystems (see review in Breitburg et al., 2018). This study aims to better understand and predict the effects of ocean deoxygenation on marine resources. The growth and reproduction responses of the Peruvian Scallop, *A. purpuratus*, under a highly variable environment, frequently limited in oxygen were modelled within the framework of Dynamic Energy Budget (DEB) theory (Kooijman, 2010). We first developed a standard DEB model for the benthic phase of *A. purpuratus*. We then added oxygen availability as a forcing variable of the model. The toxic effect due to the presence of milky waters was also taken into account by increasing somatic maintenance costs during these events. The resulting model was tested against high frequency observations of scallops exposed to hypoxic conditions in a culture area in Paracas Bay.

2. Material and methods

2.1. Field data and observed patterns

The dataset used in this study came from a survey undertaken in Paracas Bay (13°49'35" S, 76°17'43" W), between August 28, 2012 and March 10, 2013 (late austral winter to summer) on a 5-m depth scallop aquaculture bed. Two size groups of scallops (group 1: 60-70mm, mean = 66.8±CI 0.97 mm; group 2: 30-40mm, mean = 36.5± CI 1.2 mm) were exposed to two treatments. Half of both groups (780 ind.s each) was reared in the water column in suspended cages (2m above the seabed) while the another half was reared in cages placed on the bottom (30 ind. per cage). Each group/treatment was sampled weekly by removal of one cage for measurement of shell height and dry weight (soma and gonads). Environmental conditions were simultaneously monitored at both depths using autonomous data-loggers. Temperature and oxygen saturation were recorded hourly whereas fluorescence of chlorophyll-a was recorded hourly for a 24h period once a week. The occurrence of milky water events was visually monitored and recorded daily.

A small ($\approx 0.7^{\circ}\text{C}$) but significant difference was recorded between depths for the mean temperature (Wilcoxon test, $p < 0.001$), it is small. Prolonged periods of severe hypoxia/anoxia were recorded during summer. Oxygen saturation was significantly lower near bottom than in the suspended treatment (Wilcoxon test, $p < 0.001$), with median values of 25.6% and 39.4%, respectively. Chlorophyll-a concentration (proxy of trophic resource) did not present a marked seasonality or significant differences between depths. Milky water events were recorded in 2013 on Jan. 03, Jan. 28, Feb. 07 and Mar. 04 and were accompanied by anoxic conditions (Fig. 1).

74 Interestingly, although temperature and chlorophyll-a concentrations were close between culture con-
 75 ditions, important differences were observed for growth and gonadal investment (Fig. 5, data points) with
 76 higher growth in the suspended treatment compared to the bottom. In addition, in early summer (mid-
 77 December) when oxygen limitation was higher and when milky water events were more frequent, shell
 78 growth stopped in bottom-cultured scallops and was highly reduced in suspended-culture scallops after
 79 mid-January.




Figure 1: Environmental conditions monitored between August 2012 and March 2013 in Paracas Bay at 3 m depth (level of the suspended cages; blue lines) and on the bottom (red lines). (a): temperature, (b): oxygen saturation, (c): mean chlorophyll-a concentration estimated by fluorescence. The dotted vertical black lines indicate the occurrence of milky water discolouration events.

80 2.2. Peruvian scallop bioenergetics

81 2.2.1. DEB model and additional assumptions

82 A DEB model allows to quantify the energy acquisition and allocation within an individual through the
 83 dynamics of four state variables: structural volume (V), reserve (E) and reproduction buffer (E_R) and ma-
 84 turity (E_H). A conceptual scheme is given in Fig 2. Briefly, the theory assumes that energy uptake from the

85 environment is incorporated into a reserve pool. Energy mobilized from reserve is used for maintenance,
 86 growth, development and reproduction according to the so-called κ rule: a fixed proportion κ of the energy
 87 from reserve is allocated to maintenance and growth. The remaining fraction $1 - \kappa$ is spent on development
 88 (increase in maturity, juvenile stage) and reproduction (adult stage). Maintenance processes have priority
 89 over growth, development and reproduction. Life stage transition occurs at the maturity thresholds E_H^b
 90 (birth), E_H^j (metamorphosis) and E_H^p (puberty). A summary of the equations is given in Table 1. The uptake
 91 rate (i.e. assimilation rate; \dot{p}_A) is taken to be proportional to structural surface area of the organism. Dy-
 92 namics of the reserve can be computed as the difference between assimilation (\dot{p}_A) and energy mobilization
 93 (\dot{p}_C). Structural maintenance rate (\dot{p}_M) depends on structural volume in ectotherms. Maturity maintenance
 94 increases with maturity level (proportional to E_H) until puberty, i.e. once E_H^p is reached. In adults, energy
 95 for reproduction is allocated to a reproduction buffer and emptied (at least partially) at each spawning event.
 96 All physiological rates are assumed to depend on temperature in the same way and to follow an Arrhenius
 97 relationship (see Table 1). Further details regarding DEB theory can be found in Kooijman (2010), Sousa
 98 et al. (2010) and Nisbet et al. (2012). As for many bivalve species, we considered a model with acceleration
 99 between birth and metamorphosis ("abj") to better represent the larval and adult phases.




Figure 2: Conceptual scheme of the DEB model applied to the Peruvian scallop. Continuous arrows corresponds to the standard DEB model. The effects of hypoxia and toxicity related to the occurrence of milky turquoise waters are represented by dotted arrows.

100 When the mobilization rate is not sufficient to cover maintenance costs ($\kappa\dot{p}_C < \dot{p}_M$ and/or $(1-\kappa)\dot{p}_C < \dot{p}_J$)
 101 maintenance costs are covered in priority from reproduction buffer and, if not sufficient, from structure. To
 102 avoid an additional parameter, we assumed that the efficiency of remobilization of reproduction buffer from

Table 1: Summary of the equations of the standard Dynamic Energy Budget model used in this study. Parameters values, units and definitions are provided in Table 4.

Equation	Definition
$c(T) = \exp\left(\frac{T_A}{T_1} - \frac{T_A}{T}\right)$	Temperature correction factor (Arrhenius equation)
$f = \frac{X}{X + K}$	Holling type II scaled functional response to food density
$\dot{p}_A = \{\dot{p}_{Am}\} f V^{2/3} c(T)$	Assimilation rate
$\dot{p}_C = \frac{[E]}{[E_G] + \kappa[E]} ([E_G] \dot{v} V^{2/3} + \dot{p}_M) c(T)$	Reserve mobilization rate
$\dot{p}_M = [\dot{p}_M] V c(T)$	Maintenance rate
$\dot{p}_J = \dot{k}_J E_H c(T)$	Maturity maintenance rate
$\frac{dE}{dt} = \dot{p}_A - \dot{p}_C$	Reserve dynamics
$\frac{dV}{dt} = \frac{\kappa \dot{p}_C - \dot{p}_M}{[E_G]}$	Structural growth
$\frac{dE_H}{dt} = (1 - \kappa) \dot{p}_C - \dot{p}_J \quad \text{if } E_H < E_H^p$	Maturity dynamics (equals 0, in adults)
$\frac{dE_R}{dt} = ((1 - \kappa) \dot{p}_C - \dot{p}_J) \kappa_R \quad \text{if } E_H \geq E_H^p$	Reproduction buffer dynamics (equals 0 in juveniles)

structure equals κ_R .

Histological studies showed that the Peruvian scallop exhibits an important gametogenetic activity and is a "partial spawner", only releasing ripe gametes (Brown and Guerra, 1980; Avendaño and Le Pennec, 1996). In Paracas Bay, the reproductive activity occurs throughout the year (Wolff, 1988) with partial spawnings each ≈ 28 days (gametogenesis-spawning) synchronised with the lunar cycle (Cueto-Vega, 2016). Monitoring of gonado-somatic index indicated that 40% of the gonad weight is released at each spawning event (Cueto-Vega, 2016). Thus we introduced the following simple rules for handling the reproduction buffer: (1) spawning events were simulated every 28 days after the 5th of August to synchronise with the observed spawning cycles and (2) at each spawning event, 40% of the gonad content (taken as E_R) was

released.

Table 2: Equations allowing to compute the observables from the standard DEB model state variables and energy fluxes. Parameter values, units and definitions are given in Table 4. ^a: $\dot{p}_G = \kappa \dot{p}_C - \dot{p}_M$ stands for the growth flux, \dot{p}_D the dissipation flux that corresponds to the sum of \dot{p}_M , \dot{p}_J and $(1 - \kappa)\dot{p}_C - \dot{p}_J$ in juveniles, η_{O*} are the coefficients that couple oxygen fluxes to energy fluxes (\dot{p}_*). See supplementary material 1 of Thomas et al. (2018, this issue) for a detailed description of the computation of respiration rate and associated parameters definitions and units.

Equations	Units	Definition
$L_w = \frac{V^{(1/3)}}{\delta_M}$	cm	Shell height
$W_d = d_{Vd} V + \frac{E}{\rho_E}$	g_{dw}	Somatic dry weight
$W_w = \frac{W_d}{w}$	g_{ww}	Somatic wet weight
$W_{Rw} = \frac{E_R}{\rho_E w}$	g_{ww}	Gonad wet weight
$W_{d0} = \frac{E_0}{\rho_E}$	g_{dw}	Egg dry weight
$N = \frac{0.6 * E_R}{E_0}$	#	Fecundity (60% gonad = female)
$\frac{\dot{J}_O}{W_d} = \frac{\eta_{OA} \dot{p}_A + \eta_{OD} \dot{p}_D + \eta_{OG} \dot{p}_G}{W_d}$	mol O ₂ g ⁻¹ d ⁻¹	Dry weight-specific respiration rate ^a

2.2.2. Impacts of environmental stressors

As reviewed by Thomas et al. (2018, this issue), the feeding process is commonly impaired by oxygen limitation (Wu, 2002), probably due to the high oxygen demand of digestive/assimilation processes (see Kramer, 1987; Willows, 1992). We thus assumed that the assimilation rate (\dot{p}_A) was affected by low oxygen saturations according to Aguirre-Velarde et al. (2018). Under hypoxic conditions, energy production through aerobic metabolic pathways is impaired and less efficient anaerobic pathways are activated. As hypothesized in Aguirre-Velarde et al. (2018), we assumed that an organism under hypoxic conditions is limited in its ability to "mobilize" energy required for metabolic activities (maintenance, growth, and reproduction). This effect was accounted by modifying \dot{p}_C as a function of oxygen saturation. For parsimony purpose, the same oxygen correction function C_{DO} was applied on the assimilation rate \dot{p}_A and the reserve mobilization rate \dot{p}_C (Fig. 2; Eq. 2). Facing hypoxia, many organisms follow a so-called oxyregulator type

124 law (e.g. see Herreid, 1980) as illustrated in Fig. 3. They exhibit a two-phase response: 1) Respiration rate
 125 is maintained above a critical saturation point (S_{O_2c} , in %), this phase is called oxyregulation, 2) Below this
 126 point, they are not able anymore to maintain their oxygen consumption, which decreases linearly with the
 127 oxygen saturation of the surrounding water S_{O_2} ; this phase being called oxyconformity. *A. purpuratus* has
 128 been shown to exhibit such a two-phase response to hypoxia (Aguirre-Velarde et al., 2016). Accordingly,
 129 C_{DO} was modelled as single parameter two-phase linear response (eq. 3 and illustrated in Fig 3).

$$\dot{p}_A = c(T) C_{DO} \{\dot{p}_{Am}\} f V^{2/3} \quad (1)$$

$$\dot{p}_C = c(T) C_{DO} \left(\frac{[E]}{[E_G] + \kappa[E]} \dot{v} [E_G] V^{2/3} + \dot{p}_M \right) \quad (2)$$

$$\begin{aligned} \text{with } C_{DO} &= \frac{S_{O_2}}{S_{O_2c}} \quad \text{if } S_{O_2} < S_{O_2c} \\ &= 1 \quad \text{otherwise} \end{aligned} \quad (3)$$

130 where C_{DO} varies between 1 and 0. Under hypoxic conditions, below the organism regulatory capacities
 131 (i.e S_{O_2c}), the simultaneous decrease in the assimilation and mobilization fluxes restrict both reserve inputs
 132 and outputs proportionally to C_{DO} .




Figure 3: Illustration of the one parameter two-phase oxygen correction function used, based on the oxyregulator type law (Herreid, 1980): above S_{O_2c} fluxes are maintained (oxyregulation) but they decrease linearly with oxygen saturation below S_{O_2c} (oxyconformity).

133 In addition, scallop mortality events have been associated to milky waters discolourations, accompanied
 134 by high concentrations of toxic H_2S (Cabello et al., 2002; González-Hunt, 2010). In bivalves, exposition
 135 to sulfides may result in oxidative damage (Joyner-Matos et al., 2010) and production of oxydative stress
 136 proteins (Joyner-Matos et al., 2006) suggesting energy investment in cell repair. For simplicity sake, the

Table 3: Comparison between "zero-variate" observed data used to estimate model parameters and values predicted by the DEB model. References of observed data and relative errors, calculated as $\frac{\text{predicted}-\text{observed}}{\text{observed}}$, are specified.

Observable	Unit	Observed value	Predicted value	Relative error	Reference
Egg dry weight (W_{d0})	g	3.182e-08	3.102e-08	0.0251	Egg diameter (= 0.0066 cm) (pers. obs.)
Age at birth (a_b) at 18°C	d	4	0.4	0.900	Bellolio et al. (1994)
Age at puberty (a_p) at 16°C	d	100	61.3	0.386	Estim. from Mendo et al. (1989)
Life span	d	1825	1802	0.013	Stotz and Gonzalez (1997)
Shell height at birth	cm	0.01	0.015	0.538	Bellolio et al. (1994)
Shell height at puberty	cm	3	3.3	0.100	Mendo et al. (1989)
Ultimate shell height	cm	11	12.69	0.153	Wolff (1987)
Respiration rate* at 16°C	mgO ₂ h ⁻¹ g ⁻¹	0.66	1.2	0.818	Aguirre-Velarde et al. (2016)
Respiration rate* at 25°C	mgO ₂ h ⁻¹ g ⁻¹	1.13	1.98	0.752	Aguirre-Velarde et al. (2016)
Length as a function of time	d - cm	figure not shown		0.02635	Aguirre-Velarde et al. (2016)
Somatic wet weight as a function of time	d - g	figure not shown		0.08987	Aguirre-Velarde et al. (2016)
Gonad wet weight as a function of time	d - g	figure not shown		0.3913	Aguirre-Velarde et al. (2016)
Fecundity as a function of length	cm - #	figure not shown		0.3466	Aguirre-Velarde et al. (2016)

* Normoxic conditions, for 3-cm shell height individuals

effect of the occurrence of milky waters was modeled by multiplying maintenance costs (\dot{p}_M) by a constant factor for three consecutive days (average observed duration of these events). Following Muller et al. (2010), we assumed that the damages induced by H₂S increased maintenance costs. During the simulation process, this factor was gradually increased to fit the model to growth/reproduction observations.

2.3. Parameter estimation and model simulations

The parameters of the standard DEB model for *A. purpuratus* were estimated following the AmP procedure (Marques et al., 2018, Matlab routines available at <https://github.com/add-my-pet>). We used both "zero-variate" data (set of single-valued trait observations) and "uni-variate data" (a dependent variable as a function of an independent variable). Zero-variate data are shown in Table 3. Uni-variate data came from growth and reproduction observations from group 2 suspended scallops and environmental monitoring performed in Paracas Bay during the late winter to spring 2012, a period during which hypoxia events have low incidence.

Temperature, chlorophyll-*a* and oxygen saturation time series obtained during the monitoring in Paracas Bay (Fig. 1) were linearly interpolated when needed to complete hourly data and used as forcing variables for simulations. Rules for the conversions of model theoretical variables into observable variables are given in Table 2.

3. Results

3.1. Parameter estimation

The DEB parameter set obtained for *A. purpuratus* is shown in Table 4. Zero-variate data, including both life history traits and physiological rates were predicted (Tab. 3) with mean relative error of 0.148. However, age at puberty was underestimated by 38% and the maximum reproduction rate was 63% lower than the observed value. When including hypoxia and milky waters stress, the best match between simulations and observations was obtained for S_{O_2c} value of 40% combined with an increase of \dot{p}_M by a factor 6 when milky waters occur.

3.2. Simulations

The estimated half-saturation constant (K) resulted in a simulated scaled functional response (f) that remained close to 1 in both conditions (average $f = 0.954$ on the bottom and 0.956 in the suspended treatment) thus suggesting low food limitation in scallops' growth and reproduction.

The obtained C_{DO} times series (Fig. 4) are in average significantly lower on the bottom than in suspended culture ($p < 0.01$). At both depths, relatively low C_{DO} values (<0.5) occurred between November and March, with a particularly critical period between January and March during which values were close or equal to zero repeatedly, particularly at bottom. Note also that during the first half of February, considerably low C_{DO} values were obtained for consecutive days both at bottom and in suspended culture.

Model simulations including the effect of hypoxia and milky waters (fig. 5) accurately predicted the difference observed in growth and reproduction between culture conditions (Fig. 5 a and c). The model also predicted higher gonad growth and spawning amplitude in suspended treatment rather than at bottom, as observed (Fig. 5 d). The simple formulation for handling reproduction buffer, based on a 28-day spawning cycle, accurately simulated the periodic sharp decreases of gonadal weight (due to spawning events) before the beginning of environmental summer disturbances. As a consequence, simulated total wet weight was close to the observations under both experimental treatments (Fig. 5 d). Model simulations better fitted data for scallops of size group 2 (small initial size) than for those of size group 1 for which the model

Table 4: List of estimated values, symbols, and units of the parameters for the standard *A. purpuratus* DEB model. All rates are expressed for a reference temperature of $T_1 = 293.15$ K (20 °C) and a functional response $f=1$. The values in parenthesis correspond to parameter values after acceleration in an "abj" model.

Parameter	Symbol	Value	Units
<i>Primary parameters:</i>			
Digestion efficiency	κ_X	0.8	–
Maximum surface-area-specific assimilation rate	$\{\dot{p}_{Am}\}$	95.01 (482.82)	$\text{J cm}^{-2} \text{d}^{-1}$
Volume-specific somatic maintenance rate	$[\dot{p}_M]$	73.98	$\text{J cm}^{-3} \text{d}^{-1}$
Volume-specific cost for structure	$[E_G]$	2401	J cm^{-3}
Energy conductance	\dot{v}	0.056 (0.284)	cm d^{-1}
Fraction of utilized reserve for growth + som. maintenance	κ	0.6414	–
Maturity maintenance rate coefficient	\dot{k}_J	0.002	d^{-1}
Maturity threshold at birth	E_H^b	0.000176	J
Maturity threshold at metamorphosis	E_H^j	0.02324	J
Maturity threshold at puberty	E_H^p	2525	J
Fraction of the reproduction buffer fixed into eggs	κ_R	0.95	–
<i>Auxiliary and compound parameters:</i>			
Arrhenius temperature	T_A	4746	K
Shape coefficient	δ_M	0.33	–
Dry to wet weight ratio	w	0.09	–
Structure density (dry weight)	d_{Vd}	0.09	$\text{g}_{ww} \text{cm}^{-3}$
Energy density of reserve	ρ_E	23013	J g_{dw}^{-1}
Half-saturation coefficient of scaled functional response	K	0.5	$\mu\text{gChla L}^{-1}$
Maximum reserve density	$[E_m]$	$\frac{\dot{p}_{Am}}{\dot{v}}$	J cm^{-3}
Scaled reserve density	e	$\frac{[E]}{[E_m]}$	–

178 overestimated somatic and gonadal weights during the September - December 2012 period. The model
179 captured the somatic weight loss observed between January and March when the milky events occurred




Figure 4: Temporal evolution of the oxygen correction function C_{DO} obtained from environmental monitoring in Paracas bay (Fig. 1b) and taking into account S_{O_2c} of 40%. Both time series were used for the four model simulations presented in Fig. 5 (two depths and two different initial sizes).

(Fig. 5 c). At the same period, the model also predicted the decline observed sharp in the reproductive activity.

4. Discussion

4.1. Estimation of model parameters for *A. purpuratus*

The overall fit between data and predictions for the calibration procedure is good (Mean relative error MRE = 0.4) given the number of data sets we used. The adjustment for the larval phase in particular needs improvement, but our objectives was to focus on the effect of hypoxia during the juvenile and adult stages and the fit for these stages is well satisfying. Predicted age at puberty (55.8 d) was lower than observed (100 d). It is possible that our value (age for 3-cm shell height, pers. obs.) is slightly overestimated since first maturity for the Peruvian scallop has been reported at smaller sizes (1.3 and 2.5cm, Disalvo et al., 1984; Mendo et al., 1989, respectively). In addition, these observations are based on the presence of a differentiated gonad, that occurs after onset of puberty (i.e. first allocation of energy to reproduction; Kooijman, 2010). This underestimation might also be linked to the fact that only life history traits data of the benthic phase of *A. purpuratus* were used for parameters estimations; including larval trait data might improve this pattern.

The Peruvian scallop is a simultaneous hermaphroditic species, producing both sperm and oocytes. Information about sperm production is rather scarce because it is less limiting in hatchery processes. For this reason, model parametrization only took into account the female gonad. A more precise estimate of reproduction rate must include the production of both oocytes and sperm. *A. purpuratus* has high growth and reproduction rates, captured in the model through the combination of high somatic maintenance costs

200 ($[\dot{p}_M]=73.98$ compared to a typical value of $20 \text{ J cm}^{-3} \text{ d}^{-3}$; Kooijman, 2013) and low investment in struc-
 201 tural growth and maintenance ($\kappa=0.64$ compared to a typical value of 0.8, Kooijman, 2013). This pattern,
 202 allowing fast growth and high reproduction rate, is consistent with Kooijman (2013) "waste to hurry" hy-
 203 pothesis. Lavaud et al. (2014) published a DEB parameter set for *Pecten maximus*, a Pectinidae with similar
 204 maximum length but slower growth. Comparatively, the estimated values of shape coefficient are simi-
 205 lar for both species, but the κ value for *A. purpuratus* was lower while values for structural maintenance
 206 costs ($[\dot{p}_M]$) and energy conductance ($\dot{\nu}$) were higher. This combination of parameters reflects the highly
 207 energetic investment in reproduction and faster growth rate of *A. purpuratus* compared to *P. maximus*.

208 4.2. Effects of Hypoxia

209 DEB models can be useful to explore the consequences of environmental stress factors in order to better
 210 understand the ecological challenges that organisms are facing in their natural environment. The present
 211 work combined with Lavaud et al. (2018, this issue) are the first DEB applications taking into account the
 212 impact of oxygen limitation. Chlorophyll-*a* and temperature, were very similar between the two culture
 213 depths (suspended and bottom) and predicted growth and reproduction without the effect of hypoxia were
 214 thus very close between culture conditions. In both conditions food was rarely limiting. We introduced a
 215 simple rule for the effect of oxygen, based on the notion of critical oxygen saturation threshold ($S_{O_{2c}}$) below
 216 which metabolism is dependent on oxygen saturation (see e.g. Herreid, 1980). This single parameter yet
 217 could capture the observed trends in growth and reproduction between culture depths. On the seabed, where
 218 exposure to hypoxia was frequent and sometimes prolonged, scallop growth and reproduction was highly
 219 impacted. Decreasing assimilation and mobilization fluxes below $S_{O_{2c}}$ restricted both the energy input
 220 and output from reserve towards metabolic functions (maintenance, growth, reproduction). It thus seems
 221 appropriate to use a single law, based on the ability of the organisms to regulate their oxygen consumption,
 222 to constrain both reserve mobilization and assimilation fluxes.

223 A $S_{O_{2c}}$ value of 40% was used for simulations but Aguirre-Velarde et al. (2016) rather found a $S_{O_{2c}}$ of
 224 24% for juveniles. If hypoxia tolerance has been shown to depend on size, whether tolerance increases or
 225 decreases with size would depend on species (e.g. see Dupont-Prinet et al., 2013a,b). In bivalves, there
 226 is evidence that tolerance to hypoxia decreases with size (Wang and Widdows, 1993; Hicks and McMahon,
 227 2005). Recent experiments showed that bigger scallops survived hypoxia (1% saturation) for less time than
 228 smaller ones (Cueto-Vega, R., unpublished data), thus indicating size-dependence of the tolerance to hy-
 229 poxia in *A. purpuratus*. A higher $S_{O_{2c}}$ for bigger individuals is thus expected and is consistent with the idea

that the ability to extract oxygen from the water depends on surface-to-volume ratio. Further experiments are needed to confirm this pattern. In addition, in natural environments, hypoxia is accompanied by other potential environmental stressors (e.g. pH, H₂S, high POM concentrations, etc.) that might interact with the S_{O_2c} value. Further modeling approach in the DEB context could explore the effects of environmental stressors on the elemental balance level. This will require evaluating the relevance in terms of complexity-benefit. Various adaptations (biochemical, physiological, behavioural, etc.) would have to be considered while the selected approach must be sufficiently flexible.

4.3. *Effect of milky waters*

Milky turquoise waters events in Paracas Bay have been associated with the presence of hydrogen sulphide H₂S (Schunck et al., 2013). The metabolic consequences of H₂S and/or elemental sulfur on filter feeders physiology remain poorly studied (see example in Laudien et al., 2002). In the present study, maintenance costs were increased by a factor of six during a fixed period of time (3 days, average duration of milky water events) to mimic its toxic effect. Under these conditions, when reserve mobilization is close to zero (anoxia or severe hypoxia), increased maintenance needs must be paid from the structure and reproduction buffer. The estimated multiplication factor is rather high but the modified $[\dot{p}_M]$ value stays within the range reported by Kooijman (2013) for unstressed organisms. The exposure time to milky waters was relatively short and it is likely that toxic effects of H₂S damage cells/tissues and the reparation of these lesions lasts longer than 3 days. Further experiments are needed to validate the hypothesis that H₂S strongly increases the maintenance costs and to improve the formulation and the calibration of the effect of H₂S in interaction with hypoxia/anoxia.

4.4. *Model limitations*

The Peruvian scallop is naturally exposed to large fluctuations of dissolved oxygen and must present metabolic, physiological and behavioural adaptations to hypoxia. While trends in growth and reproduction are well represented by the model, the underestimation of growth predictions for suspended scallops may show limitations of the model with a simple rule to capture the whole evolutionary adaptations of this species to deal with oxygen limitation. On the other hand, bottom scallops could potentially be exposed to other environmental stressors such as increased CO₂ (decrease in pH) due to respiration. It should not be ruled out either a greater exposure of bottom-growing scallops to H₂S diffusing from the sediment. The cumulative impacts of multiple stressors would often be worse than expected for a single stressor (Crain

et al., 2008) which could explain the observed-predicted divergences, specially for scallops growing on bottom. As emphasized by Montalto et al. (2014), increasing the efforts to better characterize stressors in coastal areas at a relevant temporal resolution is necessary to improve the predictions.

The rule used for the restriction of metabolic fluxes allowed evaluating the global energetic consequences on the organism, but not the contribution of anaerobic metabolism below the regulation capacity. More or less efficient anaerobic metabolic pathways could have an impact on the organism's ability to cope with oxygen limitation (Hochachka and Buck, 1996), while the metabolism of anaerobic endproducts accumulated during hypoxia events could change the maintenance costs (e.g. oxygen debt), thus affecting the energy budget.

Primary production in the Peruvian coastal zone, though presenting a seasonality due to upwelling variability, is important throughout the year and is dominated by diatoms mainly (Rojas de Mendiola, 1981; Bruland et al., 2005). Under these conditions, the *in situ* chlorophyll-*a* monitoring carried out represents well this abundant availability at both water depths (near to bottom and 2m above the seabed). However, in other scenarios (comparison between different bays, latitudes, etc.) where the trophic resource is potentially different (e.g. in composition and/or quality), it is pertinent to include forcing variables that improve the characterization of the available food sources (e.g. Thomas et al., 2011).

Acknowledgements

This work is part of a PhD thesis supported by IRD, within the framework of the LMI DISCOH, and LabexMER (ANR-10-LABX-19-01) and the PICS BISCOT. Additional funding was provided by the International Foundation of Science (IFS, grant number A/5210-1). We would like to acknowledge the participants of the summer school MEMS (Combining Modelling and Experimental approaches for Marine organisms under Stress, Brest, Aug. 29– Sept 2, 2016) for fruitful discussions.

References

- Abele-Oeschger, D., Oeschger, R., 1995. Hypoxia-induced autoxidation of haemoglobin in the benthic invertebrates *Arenicola marina* (Polychaeta) and *Astarte borealis* (Bivalvia) and the possible effects of sulphide. J. Exp. Mar. Bio. Ecol. 187 (1), 63–80.
- Aguirre-Velarde, A., 2016. La bioénergétique du pétoncle péruvien (*Argopecten purpuratus*) dans un contexte environnemental limitant en oxygène. Ph.D. thesis, École doctorale des Sciences de la Mer, Université de Bretagne Occidentale.
- Aguirre-Velarde, A., Jean, F., Thouzeau, G., Flye-Sainte-Marie, J., 2016. Effects of progressive hypoxia on oxygen uptake in juveniles of the Peruvian scallop, *Argopecten purpuratus* (Lamarck, 1819). Aquaculture 451, 385–389.

288 Aguirre-Velarde, A., Jean, F., Thouzeau, G., Flye-Sainte-Marie, J., 2018. Feeding behaviour and growth of the peruvian scallop
 289 (*Argopecten purpuratus*) under daily cyclic hypoxia conditions. Journal of Sea Research 131, 85–94.

290 Avendaño, M., Le Pennec, M., 1996. Contribucion al conocimiento de la biologia reproductiva de *Argopecten purpuratus*
 291 (Lamarck, 1819) en Chile. Estud. Ocean. 15, 1–10.

292 Bellolio, G., Toledo, P., Campos, B., 1994. Morfologia de la concha larval y portlarval del ostion *Argopecten purpuratus* (Lamarck,
 293 1819; bivalvia, pectiniade) en Chile.

294 Breitburg, D., Levin, L. A., Oschlies, A., Grégoire, M., Chavez, F. P., Conley, D. J., Garçon, V., Gilbert, D., Gutiérrez, D., Isensee,
 295 K., Jacinto, G. S., Limburg, K. E., Montes, I., Naqvi, S. W. A., Pitcher, G. C., Rabalais, N. N., Roman, M. R., Rose, K. A.,
 296 Seibel, B. A., Telszewski, M., Yasuhara, M., Zhang, J., 2018. Declining oxygen in the global ocean and coastal waters. Science
 297 359 (6371), 1–11.

298 Brown, D., Guerra, R., 1980. Recuperación gonadal en ostión *Chlamys (Argopecten) purpurata* (Lamarck 1819) luego de evac-
 299 uación de gametos. Arch. Biol. Med. Exp 13 (3), 363–368.

300 Bruland, K. W., Rue, E. L., Smith, G. J., DiTullio, G. R., 2005. Iron, macronutrients and diatom blooms in the Peru upwelling
 301 regime: Brown and blue waters of Peru. Mar. Chem. 93 (2-4), 81–103.

302 Burnett, L. E., Stickle, W. B., 2001. Physiological responses to hypoxia. In: Rabalais, N., Turner, R. (Eds.), Coast. Hypoxia
 303 Consequences Living Resour. Ecosyst. Coast. Estuar. Stud., American g Edition. Wiley Online Library, Washington D.C., pp.
 304 101–114.

305 Cabello, R., Tam, J., Jacinto, M. E., 2002. Procesos naturales y antropogénicos asociados al evento de mortalidad de conchas de
 306 abanico ocurrido en la bahía de Paracas (Pisco, Peru) en junio del 200. Rev. peru. biol. 9 (2), 49–65.

307 Crain, C. M., Kroeker, K., Halpern, B. S., 2008. Interactive and cumulative effects of multiple human stressors in marine systems.
 308 Ecol. Lett. 11 (12), 1304–1315.

309 Cueto-Vega, R., 2016. Influencia de las variables ambientales sobre el indice gonado-somatico de la concha de abanico *Argopecten*
 310 *purpuratus* (L.1819) en la Bahía de Paracas. Thesis, Facultad de Pesqueria, Universidad Nacional Agraria La Molina.

311 Diaz, R. J., Rosenberg, R., 1995. Marine benthic hypoxia: a review of its ecological effect and the behavioural responses of benthic
 312 macrofauna. Oceanogr. Mar. Biol. 33, 245–303.

313 Disalvo, L. H., Alarcon, A., Martinez, E., Uribe, E., 1984. Progress in mass culture of *Chlamys (Argopecten) purpurata* Lamarck
 314 (1819) with notes on its natural history. Rev. Chil. Hist. Nat. 54, 35–45.

315 Dupont-Prinet, A., Pillet, M., Chabot, D., Hansen, T., Tremblay, R., Audet, C., 2013a. Northern shrimp (*Pandalus borealis*) oxygen
 316 consumption and metabolic enzyme activities are severely constrained by hypoxia in the Estuary and Gulf of St. Lawrence. J.
 317 Exp. Mar. Bio. Ecol. 448, 298–307.

318 Dupont-Prinet, A., Vagner, M., Chabot, D., Audet, C., MacLatchey, D., 2013b. Impact of hypoxia on the metabolism of Greenland
 319 halibut (*Reinhardtius hippoglossoides*). Can. J. Fish. Aquat. Sci. 70 (3), 461–469.

320 Gewin, V., 2010. Dead in the water. Nature 466 (August), 812–814.

321 González-Hunt, R. M., 2010. Auge y crisis: la pesquería de la concha de abanico (*Argopecten purpuratus*) en la región Pisco-
 322 Paracas, costa sur del Perú. Espac. y Desarro. 22, 25–51.

323 Grieshaber, M. K., Hardewig, I., Kreutzer, U., Pörtner, H. O., 1994. Physiological and metabolic responses to hypoxia in inverte-
 324 brates. Rev. Physiol. Biochem. Pharmacol. 125, 43–147.

325 Helly, J. J., Levin, L. A., 2004. Global distribution of naturally occurring marine hypoxia on continental margins. Deep Sea Res. I

51 (9), 1159–1168.

Herreid, C. F., 1980. Review hypoxia in invertebrates. *Comp. Biochem. Physiol.* 67A, 311–320.

Hicks, D. W., McMahon, R. F., 2005. Effects of Temperature on Chronic Hypoxia Tolerance in the Non-Indigenous Brown Mussel, *Perna perna* (Bivalvia: Mytilidae) From the Texas Gulf of Mexico. *J. Molluscan Stud.* 71, 401–408.

Hochachka, P. W., Buck, L. T., 1996. Unifying theory of hypoxia tolerance: Molecular/metabolic defense and rescue mechanisms for surviving oxygen lack. *Proc. Natl. Acad. Sci. United States Am. Biochem.* 93, 9493–9498.

Joyner-Matos, J., Downs, C. A., Julian, D., 2006. Increased expression of stress proteins in the surf clam *Donax variabilis* following hydrogen sulfide exposure. *Comp. Biochem. Physiol. - A Mol. Integr. Physiol.* 145 (2), 245–257.

Joyner-Matos, J., Predmore, B. L., Stein, J. R., Leeuwenburgh, C., Julian, D., 2010. Hydrogen sulfide induces oxidative damage to RNA and DNA in a sulfide-tolerant marine invertebrate. *Physiol. Biochem. Zool.* 83 (2), 356–365.

Kooijman, S., 2013. Waste to hurry: dynamic energy budgets explain the need of wasting to fully exploit blooming resources. *Oikos* 122 (3), 348–357.

Kooijman, S. A. L. M., 2010. Dynamic Energy Budget theory for metabolic organisation - third edition, cambridge Edition. Cambridge University Press.

Kramer, D. L., 1987. Dissolved oxygen and fish behavior. *Environ. Biol. Fishes* 18 (2), 81–92.

Laudien, J., Schiedek, D., Brey, T., Po, H., 2002. Survivorship of juvenile surf clams *Donax serra* (Bivalvia, Donacidae) exposed to severe hypoxia and hydrogen sulphide. *J. Exp. Mar. Bio. Ecol.* 271, 9–23.

Lavaud, R., Flye-Sainte-Marie, J., Jean, F., Emmery, A., Strand, O., Kooijman, S. A. L. M., 2014. Feeding and energetics of the great scallop, *Pecten maximus*, through a deb model. *Journal of Sea Research* 94 (0), 5–18.

Lavaud, R., Thomas, Y., Pecquerie, L., Benoit, H., Swain, D., Guyondet, T., Flye-Sainte-Marie, J., Chabot, D., 2018. Modelling the impact of hypoxia on the energy budget of Atlantic cod in two populations in the Gulf of Saint-Lawrence. *Journal of Sea Research*, this issue.

Levin, L., Ekau, W., a.J. Gooday, Jorissen, F., Middelburg, J., Naqvi, S., Neira, C., Rabalais, N., Zhang, J., 2009. Effects of natural and human-induced hypoxia on coastal benthos. *Biogeosciences* (6), 2063–2098.

Levin, L. A., 2003. Oxygen minimum zone benthos: adaptation and community response to hypoxia. *Oceanogr. Mar. Biol.* 41, 1–45.

Marques, G. M., Augustine, S., Lika, K., Pecquerie, L., Domingos, T., Kooijman, S. A. L. M., 2018. The AmP project: Comparing species on the basis of dynamic energy budget parameters. *PLOS Computational Biology* 14 (5), e1006100.

Mendo, J., Yamashiro, C., Rubio, J., Kameya, A., Maldonado, M., Guzman, S., 1989. Evaluation de la poblacion de concha de abanico (*Argopecten purpuratus*) en la bahia de Independencia, Pisco, Peru. 23 de Setiembre - 9 de Octubre de 1987. Tech. rep., IMARPE, Callao.

Montalto, V., Sarà, G., Ruti, P., 2014. Testing the effects of temporal data resolution on predictions of the effects of climate change on bivalves. *Ecol. Modell.* 278, 1–8.

Muller, E. B., Nisbet, R. M., Berkley, H. A., 2010. Sublethal toxicant effects with dynamic energy budget theory: Model formulation. *Ecotoxicology* 19 (1), 48–60.

Nisbet, R. M., Jusup, M., Klanjscek, T., Pecquerie, L., 2012. Integrating dynamic energy budget (DEB) theory with traditional bioenergetic models. *J. Exp. Biol.* 215 (6), 892–902.

Rabalais, N. N., Díaz, R. J., Levin, L. a., Turner, R. E., Gilbert, D., Zhang, J., 2010. Dynamics and distribution of natural and

human-caused hypoxia. *Biogeosciences* 7 (2), 585–619.

Rojas de Mendiola, B., 1981. Seasonal phytoplankton distribution along the Peruvian coast. *Coast. Estuar. Sci.* 1, 348–356.

Schunck, H., Lavik, G., Desai, D. K., Grosskopf, T., Kalvelage, T., Loscher, C. R., Paulmier, a., Contreras, S., Siegel, H., Holtap-
pels, M., Rosenstiel, P., Schilhabel, M. B., Graco, M., Schmitz, R. a., Kuypers, M. M., Laroche, J., 2013. Giant Hydrogen
Sulfide Plume in the Oxygen Minimum Zone off Peru Supports Chemolithoautotrophy. *PLoS One* 8 (8), e68661.

Sousa, T., Domingos, T., Poggiale, J.-C., Kooijman, S. A. L. M., 2010. Dynamic energy budget theory restores coherence in
biology. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 365 (1557), 3413–3428.

Stotz, W., Gonzalez, S., 1997. Abundance, growth, and production of the sea scallop *Argopecten purpuratus* (Lamarck 1819): bases
for sustainable exploitation of natural scallop beds in north-central Chile. *Fish. Res.* 32, 173–183.

Thomas, Y., Flye-Sainte-Marie, J., Chabot, D., Aguirre-Velarde, A., L., P., 2018. Effects of hypoxia on metabolic functions in
marine organisms: observed patterns and modelling assumptions within the context of Dynamic Energy Budget (DEB) theory.
J. Sea Res., this issue.

Thomas, Y., Mazurié, J., Alunno-Bruscia, M., Bacher, C., Bouget, J. F., Gohin, F., Pouvreau, S., Struski, C., 2011. Modelling
spatio-temporal variability of *Mytilus edulis* (L.) growth by forcing a dynamic energy budget model with satellite-derived
environmental data. *J. Sea Res.* 66 (4), 308–317.

Wang, W., Widdows, J., 1993. Calorimetric studies on the energy metabolism of an infaunal bivalve, *Abra tenuis*, under normoxia,
hypoxia and anoxia. *Marine Biology* 116 (1), 73–79.

Willows, R. I., 1992. Optimal digestive investment: A model for filter feeders experiencing variable diets. *Limnol. Oceanogr.*
37 (4), 829–847.

Wolff, M., 1987. Population dynamics of the Peruvian scallop *Argopecten purpuratus* during the El Niño phenomenon of 1983.
Can. J. Fish. Aquat. Sci. 44, 1684–1691.

Wolff, M., 1988. Spawning and recruitment in the Peruvian scallop *Argopecten purpuratus*. *Mar. Ecol. Prog. Ser.* 42, 213–217.

Wu, R. S. S., 2002. Hypoxia: from molecular responses to ecosystem responses. *Mar. Pollut. Bull.* 45, 35–45.




Figure 5: Simulations (lines) of shell height (a), total wet weight (WW) (b), somatic wet weight (c) and gonad wet weight (d) for the two size groups (initial shell height of 3.65 and 6.68 cm for group 2 and 1, respectively) of *A. purpuratus* scallops cultivated on the bottom and in suspended cages in Paracas Bay between August 2012 and March 2013. Points represent observed means and bars the 95% confidence intervals. Dotted vertical lines correspond to the different milky water events observed in Paracas Bay. Black lines correspond to simulations performed without the effect of hypoxia and milky waters. Continuous black line: bottom, dashed black line: suspended.