



**HAL**  
open science

## Origin and fate of long-chain polyunsaturated fatty acids in the Kerguelen Islands region (Southern Ocean) in late summer

Marine Remize, Frédéric Planchon, Ai Ning Loh, Fabienne Le Grand, Antoine Bideau, Eleonora Puccinelli, Aswani Volety, Philippe Soudant

### ► To cite this version:

Marine Remize, Frédéric Planchon, Ai Ning Loh, Fabienne Le Grand, Antoine Bideau, et al.. Origin and fate of long-chain polyunsaturated fatty acids in the Kerguelen Islands region (Southern Ocean) in late summer. *Journal of Marine Systems*, 2022, 228, pp.103693. 10.1016/j.jmarsys.2021.103693 . hal-03566357

**HAL Id: hal-03566357**

**<https://hal.univ-brest.fr/hal-03566357>**

Submitted on 12 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 Origin and fate of long-chain polyunsaturated fatty acids in the Kerguelen 2 Islands region (Southern Ocean) in late summer

3 **Marine REMIZE<sup>1,2</sup>, Frédéric PLANCHON<sup>1,\*</sup>, Ai Ning LOH<sup>2</sup>, Fabienne LE GRAND<sup>1</sup>, Antoine**  
4 **BIDEAU<sup>1</sup>, Eleonora PUCCINELLI<sup>1</sup>, Aswani VOLETY<sup>3</sup>, Philippe SOUDANT<sup>1\*</sup>**

5 <sup>1</sup> Univ Brest, CNRS, IRD, Ifremer, UMR 6539 LEMAR F-29280 Plouzané, France

6 <sup>2</sup> University of North Carolina Wilmington, Department of Earth and Ocean Sciences, Center for Marine Science, 5600  
7 Marvin K. Moss Ln, WILMINGTON NC 28403 USA

8 <sup>3</sup> Elon University, 50 Campus Drive, ELON NC 27244 USA

9  
10 \* Correspondence: frederic.planchon@univ-brest.fr; philippe.soudant@univ-brest.fr

## 11 **Highlights:**

- 12 - FA profiles of SPOM from the Kerguelen region during the post-bloom period was  
13 dominantly composed of PUFA with high proportions of the two LC-PUFA 20:5n-3 and  
14 22:6n-3.
- 15 - Abundance of LC-PUFA in the mixed layer derived from the phytoplankton community  
16 composed of small species (prymnesiophytes and prasinophytes) and diatoms.
- 17 - In the upper mesopelagic, LC-PUFA are maintained at high proportions according to  
18 distinct pathways, export of diatoms for 20:5n-3 and zooplankton fecal material for 22:6n-  
19 3.
- 20 - SPOM revealed high nutritional quality in the upper water column (0-300m) both in the  
21 iron-fertilized area on the Plateau and outside in HNLC waters.

## 22 **Abstract:**

23 Long-chain polyunsaturated fatty acids (LC-PUFA) are molecules produced at the basis of  
24 marine food webs and essential for ecosystem functioning. This study reports detailed fatty acid  
25 (FA) composition including the two LC-PUFA 20:5n-3 and 22:6n-3, in suspended organic matter  
26 (SPOM) from the upper 300 m collected in the Kerguelen Island region in the Southern Ocean  
27 during the post-bloom period (February–March 2018; project MOBYDICK). FA profiles were  
28 largely dominated by PUFA (53-69 % of Total Fatty Acid, TFA) regardless of stations and among  
29 PUFA, proportions of LC-PUFA were especially high, making up 27-44 % of TFA both in the ML

30 and upper mesopelagic. 20:5n-3 and 22:6n-3 co-occurred in the ML as a result of the post-bloom  
31 phytoplankton community showing a mixed composition dominated by small size  
32 phytoplankton (prymnesiophytes and prasinophytes) supplying 22:6n-3, and with diatoms in  
33 lower proportions supplying 20:5n-3. Elevated levels of LC-PUFA were observed both inside the  
34 iron-fertilized area on the Kerguelen Plateau and downstream, and outside in High Nutrient Low  
35 Chlorophyll waters located upstream of the Plateau, and appeared unrelated to site. In the upper  
36 mesopelagic, both LC-PUFA were maintained at high relative proportions suggesting an efficient  
37 and possibly fast vertical transfer from the surface. Transfer with depth seems to proceed via  
38 distinct pathways according to LC-PUFA. 20:5n-3 may be exported along with diatoms,  
39 presumably in the form of large intact cells, aggregates as well as resting spores. For 22:6n-3,  
40 transfer may involve a channeling through the heterotrophic food web resulting in its association  
41 with fecal material at depth. Channeling of 22:6n-3 could involve heterotrophic protists such as  
42 dinoflagellates and ciliates grazing on small phytoplankton, as well as larger zooplankton such as  
43 copepods and salps, possibly feeding on microzooplankton and producing fecal pellets rich in  
44 22:6n-3. According to LC-PUFA content, SPOM present throughout the upper water column (0-  
45 300 m) appeared of high nutritional quality both on- and off-plateau, and represented a valuable  
46 source of food for secondary consumers and suspension feeders.

47 **Keywords:** Essential fatty acids, vertical distribution, fatty acid export, phytoplankton  
48 diversity, diatoms, heterotrophic interactions, nutritional quality.

49

## 50 A. INTRODUCTION

51 The Southern Ocean (SO) is a vast and contrasted environment where the characteristics of  
52 pelagic ecosystems are particularly diverse. The Antarctic Circumpolar Current (ACC) composed  
53 of successive hydrographic fronts (Orsi et al., 1995) acts as a strong physical and biogeochemical  
54 boundary for biological activity, and to the south and until the Antarctic sea ice, lies the largest  
55 High Nutrient Low Chlorophyll (HNLC) zone of the world ocean. The productivity is low  
56 throughout the year in HNLC open waters essentially due to a lack of dissolved iron (Blain et al.,  
57 2007; de Baar et al., 1995; Martin, 1990), and high concentrations of unused macronutrients  
58 (phosphate, nitrate, silicic acid) persist in surface waters. In contrast, high productivity regimes  
59 are only found close or downstream of Subantarctic islands (South-Georgia, Crozet, Kerguelen,  
60 Heard) and continental shelves, which receive natural and persistent iron inputs (Blain et al., 2008;  
61 Perissinotto and Duncombe Rae, 1990; Pollard et al., 2007; van der Merwe et al., 2015; Venables  
62 and Moore, 2010). Iron fertilization allows phytoplankton, mainly initially composed of diatoms  
63 and also *Phaeocystis*, to thrive during intense and long-lasting blooms in austral spring/summer  
64 (Cavagna et al., 2015; Korb and Whitehouse, 2004; Mongin et al., 2008; Schallenberg et al., 2018;  
65 Schlosser et al., 2018; Seeyave et al., 2007). These blooms provide abundant food and energy to  
66 heterotrophic organisms and higher trophic levels, and help to maintain highly productive food  
67 webs extended from microbes to zooplankton, micronekton, up to emblematic top-predators of  
68 the SO (marine mammals, sea-birds, penguins, whales) (El-Sayed, 1988; Evans and Brussaard,  
69 2012; Pakhomov and McQuaid, 1996).

70 In the current context of climate change, major modifications are predicted for the SO such as  
71 sea surface warming, increasing stratification and reduction in nutrient supplies, southward  
72 migration of oceanographic fronts, shrinking of sea ice, as well as ocean acidification, and it  
73 already concerns some areas like the Antarctic peninsula (Constable et al., 2014; Moline et al.,  
74 2004). These climate-induced changes are likely to affect phytoplankton distribution and  
75 composition (Deppeler and Davidson, 2017), and hence to have positive or negative cascade  
76 effects on the whole food web. It is therefore important to determine what parameters control the  
77 present-day stability, health and resilience of these pristine ecosystems.

78 Among the parameters that maintains ecosystem stability, the nutritional value of  
79 phytoplankton is crucial, as it determines the amount of energy that can flow through the entire  
80 food web and controls its overall functioning. Part of nutritional quality for marine organisms is  
81 attributable to lipids, in particular to n-3 long-chain polyunsaturated fatty acids (n-3 LC-PUFA)  
82 such as 20:5n-3 (eicosapentaenoic acid, EPA) and 22:6n-3 (docosahexaenoic acid, DHA). n-3 LC-  
83 PUFA are molecules essential to all organisms and are involved in a variety of physiological  
84 processes such as growth, immunity, cell membrane function regulation and energy storage  
85 (Guschina and Harwood, 2006; Parrish, 2013). They play important roles in trophic interactions,  
86 starting from secondary consumers such as copepods and other crustacean zooplankton, for  
87 which they determine growth potential, reproduction success and general fitness (Brett et al.,  
88 2009; Brett and Müller-Navarra, 1997; Müller-Navarra et al., 2000; Pond et al., 2005). LC-PUFA are  
89 synthesized at the basis of the food web, essentially by phytoplankton and to some extent by small  
90 heterotrophs, and the supplies to consumers occur necessarily through diet (Bec et al., 2006; Guo  
91 et al., 2017; Klein-Breteler et al., 1999). The production of LC-PUFA by phytoplankton, and fatty  
92 acids (FA) in general, is highly species-specific both at the phylum and class levels (Cañavate,  
93 2019; Jónasdóttir, 2019; Parrish, 2013) and enables their use as valuable biomarkers (Dalsgaard et  
94 al., 2003). Another consequence is that marine algae do not all have the same nutritional value,  
95 the most valuable algae in terms of LC-PUFA proportions includes Prymnesiophyceae,  
96 Pavlovaphyceae, Bacillariophyceae, Dinophyceae, Cryptophyceae, Eustigmatophyceae followed  
97 by, Prasinophyceae and Mamiellophyceae (Jónasdóttir, 2019). The lowest contents are found in  
98 Chlorophyceae, Trebouxiophyceae and especially in Cyanobacteria, which are devoid of LC-  
99 PUFA (Jónasdóttir, 2019).

100 In the SO, LC-PUFA are presumably supplied by two distinct groups of phytoplankton,  
101 diatoms and flagellates, characterized by specific FA profiles. Diatoms produce preferentially  
102 20:5n-3 along with C16 monounsaturated fatty acids (MUFA) 16:1n-7, and C<sub>16</sub> PUFA 16:3n-4 and  
103 16:4n-1 (Dalsgaard et al., 2003; Parrish, 2013; Volkman et al., 1998), with only small amounts of  
104 22:6n-3 found in centric diatoms (Dunstan et al., 1993). The other essential n-3 LC-PUFA 22:6n-3  
105 together with C18 PUFA 18:5n-3 are produced preferentially by dinoflagellates as well as some  
106 Prymnesiophyceae like the coccolithophore *Emiliana huxleyi* (Jónasdóttir, 2019; Okuyama et al.,

107 1992; Thomson et al., 2004) but this latter group is scarce in the SO, and thus it is not assumed to  
108 be a potential producer. Another important Prymnesiophyceae species, *Phaeocystis sp.*, produces  
109 low levels of LC-PUFA (Nichols et al., 1991; Skerratt et al., 1995), but it can represent an important  
110 part of the autotrophic biomass especially in the early and late phase of the diatom bloom. This  
111 general picture of LC-PUFA sources is generally adopted in diet studies and relies on the  
112 identification of diagnostic FA in consumers (Atkinson et al., 2006; Ericson et al., 2018; Hellessey  
113 et al., 2020). This generalized view is probably oversimplified as it mostly relies on FA signatures  
114 obtained from culture and feeding experiments without accounting for field observations.  
115 Moreover, it only takes into account a single autotrophic production pathway for LC-PUFA,  
116 whereas alternative routes via heterotrophy and upgrading may exist. As an example, interactions  
117 among grazers such as copepods and heterotrophic protozoans (dinoflagellates and ciliates) were  
118 shown to substantially increase the nutritional quality of *Phaeocystis globosa*, most probably via the  
119 production of LC-PUFA by heterotrophic upgrading (Tang et al., 2001).

120 For the SO, available data on the natural occurrence of LC-PUFA in phytoplankton, as well as  
121 in protists in general, remain scarce, and it received a disproportionately low attention in  
122 comparison to larger size organisms (Hellessey et al., 2020; Pond et al., 2005). The few available  
123 data concern only surface samples obtained from highly productive regions of the marginal ice  
124 zone and coastal Antarctic zone (Fahl and Kattner, 1993; Gillan et al., 1981; Hernando et al., 2018;  
125 Nichols et al., 1993; Thomson et al., 2004) as well as of the frontal Subantarctic zone of the Indian  
126 sector of the SO (Mayzaud et al., 2007, 2002). This clearly limits our understanding of LC-PUFA  
127 cycling in both the surface euphotic zone where production takes place, and in the water column  
128 where organic matter (OM) and associated LC-PUFA are considered as labile fraction, and could  
129 be degraded, transformed, and even transferred to benthic communities (Budge and Parrish, 1998;  
130 Rembauville et al., 2018; Wilson et al., 2010). Recent results obtained from five long-term sediment  
131 traps deployed in naturally iron-fertilized settings and HNLC waters of the Polar Front (PF)  
132 region (South Georgia, Crozet, Kerguelen) indicate large regional and seasonal fluctuations in FA  
133 and PUFA composition of sinking OM (Rembauville et al., 2018). Variations are tightly linked to  
134 the ecological vectors responsible for the export of OM (e.g., diatom resting spores, phytoplankton  
135 aggregates, zooplankton fecal pellets) (Rembauville et al., 2016, 2015a, 2015b). On the Kerguelen

136 Plateau, LC-PUFA such as 20:5n-3, which is usually associated to diatoms resting spores (mainly  
137 *Chaetoceros*), could be transferred at depth towards the sediments (Rembauville et al., 2018).

138 Among naturally Fe-fertilized regions, the Kerguelen Island area has received considerable  
139 interest over recent decades as it is well suited for studying the contrasted properties of pelagic  
140 ecosystems found in the SO. The Kerguelen Plateau, which extends to the southeast of the islands,  
141 is a large-scale topographic feature with shallow bathymetry (< 700 m) that benefits from  
142 enhanced Fe inputs via diapycnal mixing (Blain et al., 2008; Park et al., 2008) and is subjected to a  
143 large-scale (45,000 km<sup>2</sup>) and long-lasting (October to February) diatom bloom (Mongin et al.,  
144 2008). Bloom phenology consists of two successive phases (Pellichero et al., 2020) with a first phase  
145 occurring in spring and dominated by small size diatoms forming long-chains (*Chaetoceros*,  
146 *Pseudo-nitzschia*) (Lasbleiz et al., 2016) and a second phase occurring in summer and dominated  
147 by larger size diatoms (*Eucampia*, *Corethron*) (Armand et al., 2008; Blain et al., 2021; Liu et al., 2020).  
148 Productivity is also stimulated downstream of the Plateau (according to the West to East ACC  
149 circulation) due to lateral advection of Fe-rich waters (Qu  rou   et al., 2015; Trull et al., 2015; van  
150 der Merwe et al., 2015), which promotes a diatom bloom from October to January. This latter  
151 bloom is shorter in duration and lower in intensity, and composed of different phytoplankton  
152 community composition in comparison to the central Plateau (Armand et al., 2008; Lasbleiz et al.,  
153 2016). Upstream of the Plateau, south-west of the Kerguelen Islands, open waters are away from  
154 the influence of the Fe fertilized area, productivity is drastically reduced and considered  
155 representative of HNLC conditions found in the SO (Rembauville et al., 2017). Phytoplankton  
156 biomass stays low throughout the year (climatological Chlorophyll a (Chl-a) < 0.3 µg.L<sup>-1</sup>) except  
157 during a short time period (~1 month) in December/January (Fiala et al., 1998; Kopczy  nska et al.,  
158 1998). Under HNLC regime, pico- and nano- communities composed mostly by *Phaeocystis sp.*  
159 predominate most of the time, except during summer when microphytoplankton, such as small  
160 diatoms (*Fragilariopsis sp.*) and autotrophic dinoflagellates, are more abundant (Armand et al.,  
161 2008; Fiala et al., 1998; Kopczy  nska et al., 1998; Lasbleiz et al., 2016; Rembauville et al., 2017).

162 The MOBYDICK project (Marine Ecosystem Biodiversity and Dynamics of Carbon around  
163 Kerguelen: an integrated view) was designed to complement the available description of the

164 Kerguelen region ecosystems and the main objectives were 1) to track carbon from its initial  
165 fixation at the surface to its channeling through the food web and its transfer at depth via export,  
166 and 2) to perform a detailed description of the diversity at each trophic level. The oceanographic  
167 survey was carried out in late summer/early autumn (February-March 2018) corresponding to the  
168 demise of the diatom bloom, a period that was not previously investigated by other surveys  
169 carried out during the onset (KEOPS2, October-November 2011) and decline (KEOPS1, January-  
170 February 2005) of the bloom. In this study, we investigated the distribution of FA, including the  
171 essential LC-PUFA 20:5n-3 and 22:6n-3, in suspended organic matter (SPOM) collected in the  
172 upper water column. Our aims were to explore the origin and fate of essential LC-PUFA from the  
173 surface to the upper mesopelagic zone at sites with contrasted seasonal productivity regimes  
174 found on and off the Kerguelen Plateau. Statistical analyses were used to identify the main drivers  
175 of FA profile variability. A selection of FA specific to phytoplankton classes and zooplankton  
176 activity was used to identify the main producers of LC-PUFA and to explore the impact of  
177 heterotrophic interactions in the upper water column. Finally, PUFA contents were used to  
178 discuss the nutritional quality of suspended OM and a comparison between iron-fertilized and  
179 HNLC conditions was addressed.

## 180 B. MATERIAL & METHODS


### 181 **Studied area**

182 The MOBYDICK survey was conducted in the Kerguelen Islands area in the Indian sector of  
183 the SO on board of *R/V Marion Dufresne II* during late-austral summer 2018 from February 18<sup>th</sup> to  
184 March 28<sup>th</sup> coinciding with the post bloom phase. Four stations (M1, M2, M3, and M4, Figure 1),  
185 were selected to represent the contrasted production regimes encountered in the area and defined  
186 on a seasonal basis. Station M2 (bottom depth 520 m) is located in the Fe-enriched area on the  
187 central Kerguelen Plateau. This station corresponds to the station A3 investigated during KEOPS1  
188 and KEOPS2 programs. Station M1 (bottom depth 2723 m), located downstream of the Plateau,  
189 benefits also from enhanced Fe supplies and exhibit a seasonal moderate production regime.  
190 Station M3 (bottom depth 1730 m) and M4 (bottom depth 4731 m) are both located upstream of  
191 the Plateau and are representative of HNLC conditions. Station M3, previously investigated as



192 part of the KERFIX time series program (1990-1995) (Jeandel et al., 1998), was at the time of  
193 sampling north of the PF in the Polar Frontal Zone (PFZ) according to Pauthenet et al. (2018),  
194 while the three other stations were permanently south of the PF. During the survey, repeated  
195 visits were performed at most stations at around 10-day interval except at station M1, which was  
196 visited only once. As shown in Table 1, three visits were performed at station M2 (M2-1, M2-2,  
197 and M2-3) and two at stations M3 and M4 (M3-1, M3-3, M4-1, M4-2).

### 198 Hydrological and biogeochemical data



**Figure 1** Map of MOBYDICK station locations and monthly mean (March 2018) surface Chlorophyll-a concentrations ( $\mu\text{g.L}^{-1}$ ) obtained from 4 km resolution Global Ocean Satellite Observations (Copernicus-Globcolour, Copernicus Marine Service, <http://marine.copernicus.eu/>). The black line refers to 1000 m bathymetry and the blue refers to the approximate position of the PF for the Feb.-Mar. period and drawn according to Pauthenet et al. (2018).

199 Vertical profiles of temperature, dissolved oxygen, and salinity were obtained at all stations  
200 using a SeaBird 911-plus CTD (Conductivity, Temperature, and Density) unit mounted on the  
201 rosette. Chl-a concentrations and PAR (photosynthetically available radiation) levels were  
202 measured using a fluorometer and LI-COR sensor, respectively. The mean depth of the mixed  
203 layer (MLD) was estimated based on a difference in potential density of 0.03 to the surface value  
204 (10 m) and using all CTD casts performed during the occupation of stations (Lafond et al., 2020).

205 The depth of the euphotic zone ( $Z_e$ ) corresponded to the depth where light intensity was at least  
 206 1% of incident light at the surface (Table 1). Averaged Chl-a, silicic acid ( $\text{Si}(\text{OH})_4$ ) and ammonium  
 207 ( $\text{NH}_4^+$ ) concentrations for the mixed layer (ML) are also shown in Table 1, details on the analytical  
 208 methods can be found elsewhere (Irion et al., 2020; Lafond et al., 2020).

Table 1. MOBYDICK station details, averaged depth of the mixed layer (MLD), depth of the euphotic zone ( $Z_e$ ), and averaged concentrations of Chl-a, silicic acid, ammonium in the mixed layer.

Station	Lat	Lon	Bottom depth m	date	Visit	MLD <sup>a</sup> m	$Z_e^b$ m	Chl-a		$\text{Si}(\text{OH})_4$		$\text{NH}_4$	
	°S	°E						$\mu\text{g.L}^{-1}$	S.D	$\mu\text{mol.L}^{-1}$	S.D	$\mu\text{mol.L}^{-1}$	S.D
M1	49.9	74.9	2723	10/03/2018	M1	63	89	0.34	0.04	6.7	0.3	0.57	0.21
M2	50.6	72.0	520	27/02/2018	M2-1	79	64	0.27	0.02	1.4	0.4	0.75	0.08
				08/03/2018	M2-2	73	61	0.31	0.05	1.7	0.8	1.12	0.03
				17/03/2018	M2-3	80	58	0.59	0.03	2.8	0.3	0.95	0.06
M3	50.7	68.1	1730	05/03/2018	M3-1	74	93	0.21	0.03	2.9	1.0	0.63	0.22
				19/03/2018	M3-3	96	105	0.14	0.00	2.3	0.2	0.73	0.01
M4	52.6	67.2	4731	03/03/2018	M4-1	69	95	0.19	0.01	4.4	0.4	0.37	0.03
				14/03/2018	M4-2	96	101	0.21	0.01	5.3	1.0	0.54	0.11

<sup>a</sup> MLD: Average mixed layer depth estimated from multiple CTD casts at each station and according to potential density fluctuations  $< 0.03 \text{ kg.m}^{-3}$  relative to 10 m depth.

<sup>b</sup>  $Z_e$ : Depth of the euphotic zone defined as 1% of surface photosynthetic active radiation.

209

210 **Sampling of suspended particles**

211 Suspended particulate organic matter (SPOM) was sampled for FA determination, at all  
 212 visited stations (8 stations in total) using five *in situ* pumping systems (ISP) allowing large volume  
 213 of seawater to be filtered (200-1500 L) and sufficient amount of material to be collected. The five  
 214 ISP were deployed from the surface euphotic zone to the upper mesopelagic, down to 200 m (M1,  
 215 M2-2, M2-3, M4-2), 300 m (M2-1, M3-1, M3-3), and 600 m depth (M4-1). For three stations (M1,  
 216 M2-1, M4-1), only four depths could be sampled due to ISP failures. ISP were equipped for  
 217 sequential filtration, and we considered here only the 1-50  $\mu\text{m}$  size fraction, which corresponds to  
 218 the largely dominant fraction in terms of Particulate Organic Carbon (POC, averaged weight  
 219 proportion 93% of total POC,  $n=37$ ). For the first six stations (M1, M2-1, M2-2, M4-1, M4-2, M3-1),

220 the 1-50  $\mu\text{m}$  size fraction was obtained from a single filter made of high-purity quartz microfiber  
221 (QMA, Sartorius, France) of 1  $\mu\text{m}$  nominal pore size, and placed after two successive nylon  
222 (NITEX) screen pre-filters of 300 and 50  $\mu\text{m}$  mesh size. For the last two stations (M2-3 and M3-3),  
223 due to a change in the mesh size of screens (50 and 20  $\mu\text{m}$ ), the 1-50  $\mu\text{m}$  fraction was obtained  
224 from the QMA filter (1-20  $\mu\text{m}$ ) and from the intermediate screen of 20  $\mu\text{m}$  mesh size (20-50  $\mu\text{m}$ ).

### 225 **Samples processing on board**

226 After ISP recovery, the QMA filters were processed on-board as follows: subsampling was  
227 conducted directly on the filter using a 25 mm plexiglass punch previously cleaned with ethanol.  
228 For FA analyses, four punches were taken and then extracted in 6 mL of 2:1 (v:v)  
229 chloroform:methanol solvent and preserved at  $-20^{\circ}\text{C}$ . For POC measurements, four other punches  
230 were taken, dried at  $55^{\circ}\text{C}$  for 24 hours, and stored at room temperature. For the NITEX screens,  
231 the 142 mm filters were cut into quarters using a scalpel (ethanol cleaned), and one quarter was  
232 dedicated to POC analyses and another to FA analyses. The two remaining quarters were kept as  
233 spare at  $-20^{\circ}\text{C}$ . Particles from the NITEX were resuspended using filtered seawater (0.4  $\mu\text{m}$ ) and  
234 recollected on a pre-combusted 0.7  $\mu\text{m}$  nominal pore-size glass fiber filters (Whatman GF/F,  
235 Maidstone, UK). The GF/F filters were processed as described for QMA filters for FA (dived in 6  
236 mL of chloroform:methanol solvent and stored at  $-20^{\circ}\text{C}$ ) and POC analyses. Lipid extracts  
237 dedicated to FA analyses were stored at  $-20^{\circ}\text{C}$  during the cruise, shipped to home laboratory  
238 (France) with dry ice and stored at  $-20^{\circ}\text{C}$  until FA analyses.

### 239 **Particulate organic carbon analyses**

240 POC samples were fumed with hydrochloric acid (10M) for 4 hours to remove particulate  
241 inorganic carbon, dried at ambient temperature for 2 hours, subsampled with a 13 mm punch and  
242 encapsulated into tin caps following protocols by Brook et al. (2003) and Trull et al. (2015). POC  
243 content was determined using a Flash Elemental Analyzer 2000 coupled to a Thermo Fisher Delta  
244 V Plus stable light isotope ratio mass spectrometer system (IRMS, Bremen, Germany). Data were  
245 corrected for filter blank contribution, and acetanilide ( $\text{C}_8\text{H}_9\text{NO}$ ) was used as a standard for POC  
246 content. Final concentrations of POC in seawater are reported in  $\mu\text{mol.L}^{-1}$ .

247 **Fatty acids analysis**

248 *a. Lipid extraction and purification*

249 Samples for FA analyses were re-extracted to remove any residual seawater. The initial lipid  
250 extract was transferred in a new 22 mL vial, and then re-extracted with 3 mL of chloroform. The  
251 sample was vigorously shaken and then centrifuged to insure a good phase separation. For each  
252 sample, three re-extractions were performed. Lipid extracts were then evaporated with N<sub>2</sub> gas,  
253 resuspended in 6 mL of chloroform:methanol (2:1 v:v) and stored at -20°C until analysis. Lipid  
254 extracts were separated into neutral and polar lipids following the method of Remize et al.(2020).  
255 Briefly, 3 mL of total lipid extract was evaporated with nitrogen, recovered with three washes  
256 using chloroform:methanol (final volume 1.5 mL, 98:2 v:v) and spotted at the top of a silica gel  
257 column (40 mm ×4 mm, silica gel 60A 63–200 µm rehydrated with 6% H<sub>2</sub>O, 70–230 mesh, Sigma-  
258 Aldrich, Darmstadt, Germany). The neutral lipid fraction (NL) was eluted using  
259 chloroform:methanol (10 mL 98:2 v:v) and the polar lipid fraction (PL) with methanol (20 mL).  
260 Both fractions were then collected in glass vials, and an internal standard (C23:0, 2.3 µg) was  
261 added.

262 *b. Transesterification of FAME*

263 Fatty acids methyl esters (FAME) transesterification was conducted according to the protocol  
264 described by Mathieu-Resuge et al. (2019). In brief, after evaporation to dryness of the NL and LP  
265 fractions, transesterification was performed by adding 0.8 mL of H<sub>2</sub>SO<sub>4</sub>/methanol mixture (3.4%  
266 v:v) to the lipid extract and heated at 100 °C for 10 min. Hexane (0.8 mL) and distilled water  
267 saturated with hexane (1.5 mL) were added. The lower MeOH–water phase was discarded after  
268 homogenization and centrifugation. Hexane fraction containing FAME was washed two more  
269 times with another 1.5 mL of distilled water.

270 *c. Fatty acid analysis by gas chromatography*

271 Analyses of FAME were performed on a Varian CP8400 gas chromatograph (Agilent, Santa Clara  
272 CA, USA) using simultaneously two separations on a polar column (ZBWAX: 30 mm × 0.25 mm  
273 ID × 0.2 µm, Phenomenex, Torrance, CA, USA) and an apolar column (ZB5HT: 30 m × 0.25 mm

274 ID × 0.2 μm, Phenomenex, Torrance, CA, USA). The temperature program used by the gas  
 275 chromatograph was the following: first, initial heating to 0 from 150 °C at 50 °C.min<sup>-1</sup>, then to 170  
 276 °C at 3.5 °C.min<sup>-1</sup>, to 185 °C at 1.5 °C.min<sup>-1</sup>, to 225 at 2.4 °C.min<sup>-1</sup> and finally to 250 °C at 5.5 °C.min<sup>-1</sup>  
 277 and maintained for 15 min. The FAME were identified by comparison of their retention time with  
 278 commercial and in-house standards mixtures as described in Remize et al. (2020).

Table 2. Complete list of FA determined in this study and grouped by compound family with saturated fatty acids (SAFA), monounsaturated fatty acids (MUFA), polyunsaturated fatty acids (PUFA), and branched fatty acids (Branched). Major FA (>1% of TFA of the mean of all samples) are highlighted in bold.

Group	Fatty acids
SAFA	<b>14:0</b> , 15:0, <b>16:0</b> , 17:0, <b>18:0</b> , 20:0, 21:0, 22:0, 24:0
MUFA	14:1n-5, 15:1n-5, 16:1n-9, <b>16:1n-7</b> , 16:1n-5, 17:1n-7, <b>18:1n-9</b> , <b>18:1n-7</b> , 18:1n-5, 20:1n-9, 20:1n-7, 22:1n-9, 22:1n-7, 24:1n-9
	C <sub>16</sub> : 16:2n-7, 16:2n-6, 16:2n-4, 16:3n-4, 16:3n-6, 16:3n-3, 16:4n-3, <b>16:4n-1</b>
	C <sub>18</sub> : <b>18:2n-6</b> , 18:2n-4, 18:3n-6, 18:3n-4, <b>18:3n-3</b> , 18:4n-1, <b>18:4n-3</b> , <b>18:5n-3</b>
PUFA	C <sub>20</sub> : 20:2n-6, 20:3n-6, 20:3n-3, 20:4n-6, 20:4n-3, <b>20:5n-3</b>
	C <sub>21</sub> : 21:5n-3
	C <sub>22</sub> : 22:2n-6, 22:4n-6, 22:5n-6, 22:5n-3, <b>22:6n-3</b>
Branched	iso15:0, ante15:0, iso16:0, iso17:0

279 FA concentrations reported here correspond to the total lipid fraction (sum of PL and NL  
 280 fractions) measured in 1-50 μm SPOM. Concentrations are reported in μg.L<sup>-1</sup> (of seawater) as well  
 281 as in percent mass proportion relative to total fatty acids (%TFA). The complete list of FA  
 282 considered and grouped by compound family is shown in Table 2. FA are reported using a  
 283 shorthand notation of A:Bn-x, where A indicates the number of carbon atoms, B is the number of  
 284 double bonds and x indicates the position of the first double bond relative to the terminal methyl  
 285 group (Budge et al., 2006).

## 286 Fatty acid profiles in phytoplankton classes

287 In Table S1, we summarized available information on FA composition of phytoplankton  
288 classes provided by the meta-analysis of Jónasdóttir (2019) and Cañavate (2019) as well as the  
289 screening study of Mitani et al. (2017). This allows to highlight the relationships between some  
290 specific FA abundances and phytoplankton classes.

291 As seen in Table S1, the percentage of 14:0 is usually above 10% in Mamiellophyceae (MAM),  
292 Pavlovophyceae (PAV), Pelagophyceae (PEL), Coscinodiscophyceae (COS) and Mediophyceae (MED)  
293 and above 15% in Coccolithophyceae (COC), Prymnesiophyceae (PRY), Pinguicophyceae (PIN) and  
294 Fragilariophyceae (FRA). The percentage of 16:1n-7 is above 15% in PAV, Bacillariophyceae (BAC),  
295 COS, FRA, and MED. The sum of 16:2n-7, 16:2n-4, 16:3n-4 and 16:4n-1 account for 12% in BAC,  
296 21% in COS, 12% in FRA and 19% in MED while they are only present in trace amounts in other  
297 phytoplankton classes. The percentage of 16:4n-3 is above 10% in Chlorophyceae (CHL),  
298 Chlorodendrophyceae (CHD), Pyramimonadophyceae (PYR), Prasinophyceae (PRA) and MAM.  
299 Percentage of 18:2n-6 is present at 11% in Cyanophyceae (CYA) and 14.5% in Trebouxiophyceae  
300 (TRE) while 18:3n-6 is only above 5% in CYA. 18:3n-3 and 18:4n-3 are present in most  
301 phytoplankton classes; 18:3n-3 is above 15% in CYA, CHL, TRE, CHD, and Cryptophyceae (CRY)  
302 while 18:4n-3 is above 12% in PYR, PRA, MAM, CRY, PRY, Raphidophyceae (RAP), and PEL. The  
303 percentage of 18:5n-3 is especially high in Dinophyceae (DIN) (16%) and present above 5% in PYR,  
304 PRA, COC, and PRY. The percentage of 16:4n-3 is above 12% in CHL, CHD, PYR, PRA and MAM,  
305 all belonging to the chlorophyta phylum. 20:5n-3 (EPA) ranged between 15% and 20% in PAV,  
306 RAP, BAC, COS, FRA, and MED and above 20% in Porphyridophyceae and PIN. 22:6n-3 (DHA)  
307 is above 10% in COC, and PRY and above 15% in DIN.

### 308 **Fatty acids associated to heterotrophic organisms**

309 Branched FA (iso15:0, anteiso15:0, iso16:0 and iso17:0) are associated to bacteria (Parrish, 2013;  
310 Volkman et al., 1998; Wilson et al., 2010). The MUFA 20:1n-9, 22:1n-9, 22:1n-7 and 22:1n-11 are  
311 specifically produced by zooplankton such as Calanoids (Brett et al., 2009; Dalsgaard et al., 2003;  
312 Kattner and Hagen, 1995; Mayzaud et al., 2007; Parrish, 2013; Wilson et al., 2010) and are not found  
313 in phytoplankton.

### 314 **FA-based nutritional quality index**

315 FA data were used to estimate the nutritional quality of the 1-50  $\mu\text{m}$  SPOM following the FA-  
316 based nutritional quality index (NQI) developed by Cañavate (2019). The NQI was calculated as  
317 follow:

$$318 \quad \text{NQI} = [(15 * \text{DHA} + 10 * \text{EPA} + 2 * \text{ARA}) * 0.8 + (1.8 * \sum \text{C18PUFA})] * \log\left(\frac{n-3}{n-6}\right)$$

319 DHA, EPA and 20:4n-6 (arachidonic acid, ARA) are expressed as % of TFA.  $\sum \text{C18PUFA}$  is the  
320 sum of 18:2n-6, 18:3n-3, 18:4n-3 and 18:5n-3 percentages and n-3/n-6 is the ratio of total n-3 PUFA  
321 to total n-6 PUFA.

### 322 **Statistical analysis**

323 Principal Component Analysis (PCA) was used to characterize relationships among FA and  
324 to discriminate FA profiles according to depth and station. The PCA was performed with 36 FA  
325 out of 55 analyzed whose average abundance was >0.1% of TFA. Resulting components 1, 2 and  
326 3 were analyzed using one-way analysis of variance (ANOVA) to identify differences of FA  
327 profiles according to depth intervals (ML, MLD-150 m, 150-300 m) and stations. Categories of  
328 branched FA, SAFA, MUFA, short-chain PUFA (SC-PUFA), and LC-PUFA were analyzed by non-  
329 parametric one-way ANOVA (Kruskal-Wallis) with post-hoc analysis to determine significant  
330 differences between depth zones (ML, MLD-150 m, 150-300 m). Furthermore, spearman tests were  
331 conducted to explore the relationship between some FA and depth. In the surface mixed layer  
332 (ML), FA profiles of the 1-50  $\mu\text{m}$  fraction were analyzed by non-parametric one-way ANOVA  
333 (Kruskal-Wallis) with post-hoc analysis to determine significant differences among stations.  
334 Spearman tests were also conducted to explore the relationship between FA profiles and physico-  
335 chemical parameters (Temperature, Salinity, Oxygen,  $\text{NH}_4^+$ ,  $\text{Si}(\text{OH})_4$ ,  $\text{NO}_3^-$ ) in the ML. Differences  
336 were considered statistically significant if  $p < 0.05$ . All statistical analyses were performed using  
337 Statgraphics Plus statistical software (Manugistics, Rockville, MD, USA).

## 338 **C. RESULTS**

### 339 **Hydrological and biogeochemical context**

340 During the first half of the MOBYDICK survey, from late February to early March, the MLD  
341 was relatively shallow at all stations (Table 1), varying from 63 m at station M1 to 79 m at station  
342 M2-1, and it was consistent with late summer conditions (stable upper water column and low  
343 wind stress). After a storm event that occurred on the 10<sup>th</sup> of March, the MLD deepened  
344 consistently at stations M4-2 and M3-3 reaching 96 m, and to a lesser extent on the Plateau at  
345 station M2-3 (80 m). The depth of Ze was shallow on the plateau (58-64 m) and above the MLD.  
346 Ze was deeper at the other stations (89 m at M1 down to 105 m at M3-3) and it was below the  
347 MLD indicating no light limitation in the ML.

348 In the ML, Chl-a exhibited low values at all stations, with 0.27-0.59  $\mu\text{g.L}^{-1}$  and 0.14-0.34  $\mu\text{g.L}^{-1}$   
349 on- and off-plateau, respectively. At station M2, Chl-a almost doubled between the second (M2-  
350 2) and the third visit (M2-3). Silicic acid had variable concentrations in the ML (Table 1). These  
351 were the lowest on the plateau at M2-1 and M2-2 (1.4-1.7  $\mu\text{M}$ ), intermediate at M3 (2.3-2.9  $\mu\text{M}$ )  
352 and the highest at M4 (4.8  $\mu\text{M}$ ) and M1 (6.7  $\mu\text{M}$ ). As a comparison, ammonium concentrations, an  
353 indicator of heterotrophic excretion, exhibited a reverse trend with highest values on the plateau  
354 (0.7-1.1  $\mu\text{M}$ ) decreasing to 0.6-0.7  $\mu\text{M}$  at M3 and M1, and were the lowest at M4 (0.3-0.5  $\mu\text{M}$ ).

#### 355 **Vertical distribution of TFA concentrations**

356 Depth weighted average (DWA) of TFA concentrations obtained for three depth intervals  
357 (ML, MLD-150 m, 150-300 m) are shown in Figure 2. Considering all stations and visits, TFA  
358 concentrations were significantly ( $p < 0.001$ ) higher in the ML (range: 1.4-3.6  $\mu\text{g.L}^{-1}$ ) as compared  
359 to the deeper zones (MLD-150 m and 150-300 m). Although TFA concentrations tended to be lower  
360 in the 150-300 m depth interval (range: 0.2-0.7  $\mu\text{g.L}^{-1}$ ) compared to the MLD-150 m zone (range:  
361 0.6-1.6  $\mu\text{g.L}^{-1}$ ), the difference was not significant. TFA concentrations were significantly related to  
362 POC ( $R^2 = 0.91$  and  $p < 0.001$ ,  $n = 24$ ) and therefore to biomass variability. Based on the slope of the  
363 regression, the average concentration of TFA in biomass was estimated to be  $79 \pm 5 \mu\text{g.mgC}^{-1}$ .



364 In the ML, TFA concentrations were not significantly ( $p>0.05$ ) different between on- and off-  
 365 Plateau stations despite different temporal trends. On the Plateau, TFA concentrations increased  
 366 from 1.7 to 2.9  $\mu\text{g.L}^{-1}$  between stations M2-1 and M2-3. Off the Plateau, TFA concentrations  
 367 decreased during the course of the survey, from 2.3 to 1.5  $\mu\text{g.L}^{-1}$  at station M3 and from 2.4 to 1.4  
 368  $\mu\text{g.L}^{-1}$  at station M4. The highest TFA concentration in the ML (3.6  $\mu\text{g.L}^{-1}$ ) was found at station M1  
 369 downstream of the Plateau. Below the ML, no significant differences ( $p>0.05$ ) were observed  
 370 according to station locations. On the Plateau, TFA concentration was higher during the first visit  
 371 of M2-1 (1.6  $\mu\text{g.L}^{-1}$ ) decreasing to 0.6  $\mu\text{g.L}^{-1}$  at M2-2, ten days later.




Figure 2: Depth weighted average TFA concentrations ( $\mu\text{g.L}^{-1}$ ) obtained in three depth intervals (ML, MLD-150m, 150-300m) according the station-visit sampled during the cruise. Also shown, the MLD (m) which was estimated for all stations.

### 372 FA partitioning according to unsaturation level

373 A preliminary partitioning of FA was performed according to the unsaturation level as  
 374 described in Table 2 and considering branched FA, SAFA, MUFA and PUFA. Within PUFA, we  
 375 further distinguished SC-PUFA with 16 and 18 atoms of carbon from long-chain PUFA (LC-  
 376 PUFA) containing 20 and 22 atoms of carbon, which included the essential 20:5n-3 and 22:6n-3.

377 Depth weighted average percentages (relative to TFA) of these five groups are shown in Figure 3.  
 378 Considering all stations and depths, PUFA were the predominant category (62% on average),




Figure 3: Depth-weighted averaged percentages (relative to TFA) of branched FA (BRCH), saturated FA (SAFA), monounsaturated FA (MUFA), short-chain (C16-C18) and long-chain (C20-C22) polyunsaturated FA (SC-PUFA and LC-PUFA) in three depth intervals (ML, MLD-150, 150-300m).

379 followed by SAFA (21% on average), MUFA (15% on average), and Branched FA (2% on average).  
 380 Within PUFA, LC-PUFA were dominant (range: 34-44%) and showed increasing and significantly  
 381 different ( $p < 0.001$ ) proportions according to depth interval, 34% in the ML, 37% in the MLD-150  
 382 m, and 45% in the 150-300 m. On the opposite, SC-PUFA decreased substantially with depth, from  
 383 30% in the ML down to 12% in the 150-300 m depth interval, and differences between all depth  
 384 intervals were significant ( $p < 0.001$ ). For SAFA and MUFA, proportions increased with depth.  
 385 Differences were significant ( $p < 0.01$ ) only in the 150-300 m compared to the ML and MLD-150 m  
 386 for SAFA, and in the ML compared to the MLD-150 m and 150-300 m for MUFA.

387 In the ML, no significant differences were observed for SAFA, MUFA, and Branched between  
 388 on- and off-plateau. LC-PUFA tended to be higher off-plateau compared to on-plateau while SC-  
 389 PUFA tended to be higher on the plateau vs off-plateau. Below the ML, no significant differences

390 were observed among stations. However, the decrease of SC-PUFA with depth was faster on the  
391 plateau (from 34 to 20%) as compared to off-plateau (from 28 to 24%).

392 LC-PUFA were dominated by 20:5n-3 and 22:6n-3, both increasing significantly with depth  
393 from 13% to 19% for 20:5n-3 and from 18% to 20% for 22:6n-3. SC-PUFA decreased significantly  
394 with depth due to the decreases of 18:3n-3, 18:4n-3 and 18:5n-3. In contrast, C16 PUFA associated  
395 to diatoms were significantly higher in deep zones (MLD-150m and 150m-300m divisions) than  
396 in the ML.

### 397 **Overall pattern in individual FA abundance**

398 Principal component analysis was applied to 36 FA relative abundances (expressed as % of  
399 TFA) by examining 37 cases including FA compositions from 8 stations at various depths and  
400 visits. As shown in Figure 4, axis 1 and 2 explained 34.1% and 17.1% of the variability, respectively,  
401 and axis 3 12.3% (not shown in Fig. 4). These three axis accounted for >63% of total variability.  
402 The positive side of axis 1 was driven by 15:0, 16:0, 18:1n-7, 20:1n-9, 22:1n-9, and 22:1n-7 while  
403 17:1n-7, 18:2n-6, 18:3n-6, 18:3n-3, 18:4n-3 and 18:5n-3 were correlated with the negative side of  
404 axis 1. The second axis of the PCA was positively associated to diatom markers including 16:2n-  
405 7, 16:2n-4, 16:3n-4, 16:4n-1 and 20:5n-3 and negatively associated to 18:1n-9. The third axis was  
406 positively correlated to 16:4n-3. All correlations previously listed (positive or negative) between  
407 FA variables and principal components 1, 2, and 3 were highly significant and had correlation  
408 coefficients greater than 0.7. As PC1, PC2, and PC3 revealed good correlations (positive and  
409 negative) between FA variables, non-parametric analyses of variance with PC1, PC2, and PC3 as  
410 the dependent variables and depth intervals and stations as the independent variables were  
411 performed. The analysis showed that differences in FA profiles attributable to depth were highly  
412 significant (Figure 5) for both PC1 and 2 with p values <0.001 in both cases. While PC1 value  
413 increased steadily with depth zones, PC2 value of the MLD-150 m was higher than in the ML and  
414 150-300m zones. While PC1 and PC2 were not significantly different according to stations, PC3  
415 allowed significant differentiation between on-plateau station M2 and off-plateau stations M3 and  
416 M4 (p<0.01, Figure 5).

417




Figure 4: Correlation circle of 36 fatty acid variables according to the two principal components 1 and 2 obtained from the Principal Component Analysis, which used 37 cases including FA compositions of 1-50  $\mu\text{m}$  suspended particulate organic matter from 8 stations at various depths. The outer circle represents the unit circle in terms of correlation coefficient (positive or negative along each axis). The closer the FA variables are to the outer circle, the higher the correlation coefficient.




Figure 5: (Left) Non-parametric analysis of variance with PC1 and PC2 as the dependent variables (all stations pooled) and depth intervals as the independent variable; (Right) Non-parametric analysis of variance with PC2 and PC3 as the dependent variables (all depth intervals pooled) and stations as the independent variable. Letters a, b and c indicate statistical differences among conditions.

421 **FA variability according to depth and stations**




Figure 6: Percentages (relative to TFA) of C18 PUFA (sum of 18:3n-3, 18:4n-3 and 18:5n-3), C16 PUFA (sum of 16:2n-7, 16:2n-4, 16:3n-4 and 16:4n-1), 22:6n-3 and 20:5n-3 according to depth intervals. Letters a, b and c indicate statistical differences among conditions.

422 Furthermore, we explored the relationships between proportions of individual FA and depth  
 423 (considering all stations together). Depth explained significantly the variability of 17:1n-7  
 424 ( $R^2=61\%$ ,  $p<0.0001$ ), 18:2n-6 ( $R^2=30\%$ ,  $p<0.001$ ), 18:3n-6 ( $R^2=37\%$ ,  $p<0.001$ ), 18:3n-3 ( $R^2=44\%$ ,  
 425  $p<0.0001$ ), 18:4n-3 ( $R^2=49\%$ ,  $p<0.0001$ ), 18:5n-3 ( $R^2=44.7\%$ ,  $p<0.0001$ ) through negative linear  
 426 regressions.  $R^2$  values were improved using Log regression for 18:2n-6 ( $R^2=50\%$ ,  $p<0.0001$ ), 18:3n-  
 427 3 ( $R^2=61\%$ ,  $p<0.0001$ ), 18:4n-3 ( $R^2=68\%$ ,  $p<0.0001$ ), 18:5n-3 ( $R^2=68\%$ ,  $p<0.0001$ ). At the opposite,  
 428 depth explained significantly the variability of 16:0 ( $R^2=68\%$ ,  $p<0.0001$ ), 18:1n-9 ( $R^2=56\%$ ,  $p<0.001$ ),  
 429 20:1n-9 ( $R^2=78\%$ ,  $p<0.0001$ ), 22:1n-7 ( $R^2=41\%$ ,  $p<0.0001$ ), 22:1n-9 ( $R^2=36\%$ ,  $p<0.001$ ), and 22:5n-3  
 430 ( $R^2=79\%$ ,  $p<0.0001$ ), through positive linear regressions. The best fitting regressions for 18:4n-3,  
 431 18:5n-3, 20:1n-9 and 22:5n-3 as a function of depth are presented in Figure S1. Variations according  
 432 to depth concerned also LC-PUFA 20:5n-3 and 22:6n-3, and diatom C16 PUFA (the sum of 16:2n-

433 7, 16:2n-4, 16:3n-4 and 16:4n-1) as illustrated in Figure 6 for the three depth intervals considered.  
 434 Averaged proportions of 20:5n-3 and diatom C16 PUFA were significantly higher in the MLD-  
 435 150m and 150-300m depth intervals in comparison to the ML. For 22:6n-3, relative proportions  
 436 remained high throughout the upper water column (0-300m) ranging from 18 to 20% of TFA, and  
 437 were significantly higher in the deepest depth zone (150-300m).




Figure 7: Percentages (relative to TFA) of C18 PUFA (sum of 18:3n-3, 18:4n-3 and 18:5n-3), C16 PUFA (sum of 16:2n-7, 16:2n-4, 16:3n-4 and 16:4n-1), 22:6n-3 and 20:5n-3 according to stations in the ML. Letters a and b indicate statistical differences among conditions.

438 Considering the significant differences between depth intervals, we compared FA profiles among  
 439 stations within depth zones (Figure 7 for the ML). In the ML, the sum of 18:3n-3, 18:4n-3 and 18:5n-  
 440 3 proportions was significantly higher at M1 and M2 stations compared to M3 and M4 stations off  
 441 the plateau. The sum of diatoms C16 PUFA (16:2n-7, 16:2n-4, 16:3n-4 and 16:4n-1) and the

442 proportion of 20:5n-3 were significantly higher at station M4 than at stations M2 and M3. No  
443 difference among stations were found within both MLD-150m and 150m-300m zones.

444 To attempt explaining difference of FA composition among stations, relationships between  
445 physico-chemical parameters and FA composition were explored. We only considered regressions  
446 with  $R^2 > 40\%$  with  $p$  value  $< 0.01$ . Diatoms C16 PUFA and 20:5n-3 proportions were statistically  
447 negatively correlated to temperature ( $R^2=47\%$  and  $44\%$ , respectively, with  $p < 0.01$ , Figure S2). The  
448 sum of 18:3n-3, 18:4n-3 and 18:5n-3 was positively correlated to  $\text{NH}_4^+$  ( $R^2=41\%$ ,  $p < 0.01$ , Figure S2).

#### 449 Diatom C16 PUFA and 20:5n-3




Figure 8: Correlation between diatoms C16 PUFA and 20:5n-3 proportions (% of total fatty acid-TFA) for the mixed layer (ML) (a) and for ML depth (MLD)-300m depth interval (b).

450 As diatoms C16 PUFA and 20:5n-3 were gathered in the PCA, we explored further this  
451 relationship in Figure 8. For the ML (Fig. 8a), C16 PUFA and 20:5n-3 appeared highly related  
452 regardless of stations ( $R^2=0.945$ ,  $p < 0.001$ ). The slope of the regression indicated an average ratio of  
453 20:5n-3 to C16 PUFA of  $3.6 \pm 0.2$ , substantially higher than the ratio obtained for four diatom classes  
454 (Bacillariophyceae, Coscinodiscophyceae, Fragilariophyceae, and Mediophyceae) gathered from  
455 literature (see Table S1), which ranged from 0.9 to 1.6. Considering the MLD-300m depth interval  
456 (Fig. 8b), the correlation between C16PUFA and 20:5n-3 was lower but still significant ( $R^2=0.487$ ,  
457  $p < 0.001$ ) and the average 20:5n-3 to C16 PUFA ratio decreased to  $1.6 \pm 0.2$ .






Figure 9: Linear regressions between 18:3n-3 and 22:6n-3 (% total fatty acid-TFA) in the mixed layer for off-plateau stations (M1, M3 and M4) and on-plateau station (M2).

459 From the PCA, 22:6n-3 was poorly correlated to axis 1 ( $R^2=12.5\%$ ,  $p<0.05$ ) and not correlated  
 460 to axes 2 and 3 when considering all stations and depths together. To further explore the  
 461 relationship with non-diatom markers, we plotted in Figure 9 18:3n-3 against 22:6n-3 proportions  
 462 in the ML. Although the numbers of data points reduced its statistical strength, separating the  
 463 samples according to station locations (off-Plateau versus on-Plateau) allowed to obtain a  
 464 significant ( $R^2=0.907$ ,  $p<0.001$ ) positive relationship between 18:3n-3 and 22:6n-3 for off-plateau  
 465 stations (M1, M3, and M4). For the on-plateau station M2, the relationship was not significant  
 466 ( $p=0.17$ ) although 18:3n-3 and 22:6n-3 were negatively correlated.

467 22:6n-3 to n-3 C18 PUFA ratio increased linearly with depth between the surface and 300 m  
 468 ( $R^2=0.726$ ,  $p<0.0001$ ; Figure 10). Significant relationships were observed between 22:6n-3/n-3 C18

469 PUFA ratio and 22:5n-3 and 20:1n-9 ( $R^2=0.874$  with  $p<0.0001$  and  $R^2=0.551$  with  $p<0.001$ ,  
470 respectively) are shown in Figure S3.




Figure 10: Linear regression between 22:6n-3 to n-3 C18 PUFA ratio according to depth (m).

471

472 **FA-based nutritional quality of SPOM**

473 The nutritional quality index (NQI) deduced from FA is presented in Figure 11. NQI ranged  
474 from 354 to 477 (average  $394 \pm 37$ ) and did not vary significantly according to depth zones. It has  
475 to be noted that NQI was more variable with increasing depth. When considering together the  
476 three depth zones, NQI was slightly higher on-plateau than off-plateau.




Figure 11: FA-based nutritional quality index of SPOM (1-50  $\mu\text{m}$ ) according to stations and depth intervals.

477 **D. Discussion**

478 This study investigated the upper water column distribution of 36 FA in SPOM (1-50  $\mu\text{m}$ )  
479 collected in the Indian sector of the Southern Ocean in the region of Kerguelen Islands. Principal  
480 component analysis (PCA) revealed contrasted FA profiles attributable to depth for both PC1 and  
481 2. The PC3 also revealed significant differences between HNLC stations M3 and M4 and the  
482 naturally-iron fertilized station M2. In the following, we discuss first how the post-bloom  
483 phytoplanktonic community and depth impact the FA composition of SPOM in terms of SC-PUFA  
484 and MUFA. We then consider the potential sources of essential LC-PUFA, 20:5n-3 and 22:6n-3, in

485 the ML and their fate in the upper mesopelagic. Finally, the FA-based nutritional quality of SPOM  
486 is discussed according to depth and stations location.

### 487 **Post-bloom phytoplankton community and FA profiles**

488 The MOBYDICK survey took place in late summer/early autumn coinciding with post-bloom  
489 conditions on and off the Kerguelen Plateau. Chl-a was low in the ML ( $<0.6 \mu\text{g.L}^{-1}$ ), and POC and  
490 TFA concentrations were  $<45 \text{ mg.L}^{-1}$  and  $<4.5 \mu\text{g.L}^{-1}$ , respectively. The phytoplanktonic  
491 community, as described using 18S rDNA amplicon sequencing, pigments, flow cytometry, and  
492 microscopic enumeration (Irion et al., 2021, 2020; Lafond et al., 2020), revealed a mixed  
493 composition with three main groups, prymnesiophytes, diatoms and prasinophytes showing  
494 variable proportions on- and off the Plateau. Nano-phytoplankton (prymnesiophytes) were the  
495 most abundant group (37–53% and 59–70% of total Chl-a on- and off-plateau, respectively)  
496 followed by diatoms (27-40% and 18-33% of total Chl-a on- and off-plateau, respectively), and  
497 finally by pico-phytoplankton (prasinophytes) (Irion et al., 2021, 2020). This latter group was  
498 found in low proportions at all stations except on the plateau where it reached up to 16% of total  
499 Chl-a at the end of the survey (M2-3) (Irion et al., 2020). The mixed composition of the  
500 phytoplankton community was well reflected in the FA profile of SPOM, especially when  
501 considering chloroplastic C16 and n-3 C18 PUFA (Alonso et al., 1998; Guschina and Harwood,  
502 2006), which derive mainly from autotrophic biomass. n-3 C18 PUFA (sum of 18:3n-3, 18:4n-3, and  
503 18:5n-3) are characteristic of non-diatom phytoplankton including prymnesiophytes (24% of n-3  
504 C18 PUFA) and prasinophytes (42% n-3 C18 PUFA), and were particularly abundant in the ML  
505 representing between 17 to 32% of TFA. n-3 C18 PUFA in the ML were significantly ( $p<0.05$ )  
506 higher on the Plateau ( $30\pm 4\%$  on average at station M2) and downstream (26% at M1) in  
507 comparison to HNLC off-plateau stations ( $19\pm 4\%$  on average at stations M3 and M4). On the  
508 Plateau, n-3 C18 PUFA increased from 22% (M2-1) to 32% (M2-2 and M2-3) during the survey  
509 probably illustrating the increasing proportion of Prasinophytes.

510 For diatoms, their abundance could be identified using C16 PUFA (sum of 16:2n-7, 16:2n-4,  
511 16:3n-4 and 16:4n-1), which are highly specific to this phylum with an averaged proportion of  
512  $16\pm 4\%$  (see Table S1 for details). Diatom C16 PUFA were found at all stations in the ML. C16 PUFA

513 were the lowest on the Plateau at station M2 ( $1.2 \pm 0.2\%$  on average) and upstream at station M3  
514 ( $1.3 \pm 0.2\%$  on average) and were significantly ( $p < 0.05$ ) higher upstream at station M4 ( $2.3 \pm 0.6\%$  on  
515 average) compared to the other stations. Variability of C16 PUFA according to stations in the ML  
516 could be compared to estimates of diatom abundance deduced from pigments (Irion et al., 2020),  
517 diatom biomass contribution to POC as well as BSi (Lafond et al., 2020). For off-plateau stations  
518 (M1, M3, and M4), C16 PUFA matched the trends in BSi concentrations ( $2.2 \mu\text{mol.L}^{-1}$  at M4  $> 1.0$   
519  $\mu\text{mol.L}^{-1}$  at M1  $> 0.4 \mu\text{mol.L}^{-1}$  at M3) as well as diatom contribution to POC ( $24 \pm 6\%$  at M4  $> 16 \pm 1\%$   
520 at M1  $> 6 \pm 2\%$  at M3). On the Plateau, differences arose and C16 PUFA proportions were the lowest  
521 while diatoms were found to be more abundant, 27-40% of total Chl-a (Irion et al., 2020) and 17 to  
522 43% of POC (Lafond et al., 2020). This was especially the case at the last visit on the Plateau (station  
523 M2-3), where a bloom of *Corethron inerme* was observed (Lafond et al., 2020) which contributed to  
524 an increase of TFA and POC concentrations ( $2.1$  to  $2.9 \mu\text{g.L}^{-1}$  and  $27$  to  $40 \mu\text{g.L}^{-1}$ , respectively). In  
525 contrast, this effect was not reflected in C16 PUFA, whose proportions stayed relatively stable  
526 between station M2-2 ( $1.0\%$ ) and M2-3 ( $1.2\%$ ). This pattern could indicate that this late summer  
527 diatom bloom was probably poor in chloroplastic C16 PUFA.

### 528 **Changes in FA profiles with depth**

529 Below the ML, TFA and POC decreased rapidly and substantial changes were evidenced at all  
530 stations with a consistent decrease in n-3 C18 PUFA (from  $24 \pm 6\%$  in the ML to  $16 \pm 6\%$  in the  
531 MLD-150m down to  $5 \pm 1\%$  in the 150-300 m depth zones) and an increase in C16 PUFA (from  $1.6$   
532  $\pm 0.6\%$  in the ML to  $3.5 \pm 1.1\%$  in the MLD-150m and to  $3.1 \pm 1.4\%$  in the 150-300 m intervals). The  
533 observed pattern in phytoplankton-derived FA was in agreement with the vertical distribution of  
534 pigments showing a rapid decrease in prymnesiophytes pigments below the ML and an  
535 increasing contribution of diatom pigments, the latter representing up to 77-96% of total Chl-a at  
536 250 m depth (Irion et al., 2021). In addition, we noted variable vertical trends of n-3 C18 PUFA  
537 and C16 PUFA as observed with pigments (Irion et al., 2021). On the Plateau and downstream  
538 (Station M2 and M1) n-3 C18 PUFA decreased rapidly below the MLD, while at station M3 C18  
539 PUFA remained relatively high in the MLD-150m depth interval ( $22\%$  on average) and decreased  
540 only below 150 m ( $6\%$  on average). For C16 PUFA and diatoms, the increase with depth was

541 particularly rapid on the Plateau at the first two visits (1.5 to 5.5% at M2-1 and 1.0 to 4.3% at M2-  
542 2 between the ML and the MLD-150m depth zone) indicating an important accumulation of  
543 diatoms just below the MLD. Considering that biomass and associated FA increased below the  
544 ML as a result of vertical transfer from the surface, the observed trends in n-3 C18 PUFA and C16  
545 PUFA proportions may be consistent with a more efficient export of diatoms in comparison to  
546 nano- and pico-phytoplankton below the ML. This feature is commonly observed in the SO (Salter  
547 et al., 2012, 2007) and was documented in details on the Plateau (Blain et al., 2021; Rembauville et  
548 al., 2015a, 2015b) and also in the surrounding HNLC areas (Rembauville et al., 2017).

549 Unexpectedly and along with n-3 C18 PUFA, the MUFA 17:1n-7 decreased linearly with  
550 depth. As its distribution was concomitant with the one of chloroplastic n-3 C18 PUFA, it  
551 suggested its production was mostly autotrophic and occurred in the ML. 17:1n-7 is not listed in  
552 the meta-analysis of Jónasdóttir (2019) and Cañavate (2019) and is only rarely cited in papers  
553 analyzing FA in phytoplankton species. The cyanobacteria *Synechococcus elongates*, *Microcystis*  
554 *aeruginosa* and *Anabaena variabilis* were reported to contain respectively 0.6%, 0.5% and 1.3 of  
555 17:1n-7 (Bec et al., 2006; Martin-Creuzburg et al., 2008). Interestingly, the heterotrophic  
556 nanoflagellates *Paraphysomonas sp.*, concentrated this MUFA at 3.4% and 1.5%, when fed *S.*  
557 *elongates*, *M. aeruginosa*, respectively (Bec et al., 2006). Indeed, these planktonic groups were  
558 observed during the MOBYDICK survey. Heterotrophic nanoflagellates, pico and nano-plankton  
559 varied between 3.3 and 4.7 10<sup>6</sup> cells.L<sup>-1</sup>, 0.8 and 2.0 10<sup>6</sup> cells.L<sup>-1</sup>, and 73 and 111 10<sup>3</sup> cells.L<sup>-1</sup>  
560 respectively, in the ML (Christaki et al., 2020). However, while heterotrophic nanoflagellates were  
561 more abundant at M2 than at off-plateau stations (M3 and M4) (Christaki et al., 2020), no  
562 significant difference was observed according to stations for both concentration (in µg.L<sup>-1</sup>) and  
563 relative proportion of 17:1n-7.

564 The long chain MUFA 20:1n-9, 22:1n-7 and 22:1n-9, which are mainly produced by  
565 zooplankton such as calanoid copepods and other crustaceans (Brett et al., 2009; Dalsgaard et al.,  
566 2003; Kattner and Hagen, 1995; Mayzaud et al., 2007; Parrish, 2013; Wilson et al., 2010) can be used  
567 to detect potential interactions with zooplankton via the production of fecal pellets and their  
568 presence into SPOM (Mayzaud et al., 2007; Sheridan et al., 2002; Wilson et al., 2010). Fecal pellets

569 are known to play an important role in the vertical transfer of OM in the Kerguelen region  
570 (Ebersbach and Trull, 2008; Laurenceau-Cornec et al., 2015), particularly on the Plateau during the  
571 post-bloom period (Rembauville et al., 2015a). In our samples, long chain MUFA 20:1n-9, 22:1n-7  
572 and 22:1n-9 were detected at all stations, their proportions were low in the ML (sum of 20:1n-9,  
573 22:1n-7, and 22:1n-9 ranging between 0.11 and 0.25%) and increased linearly with depth ( $R^2=0.683$ ,  
574  $p<0.0001$ ) reaching the highest values in the 150-300m depth zone (range: 0.21-0.61%). Although  
575 particularly low, these proportions still appeared comparable to those measured in fecal pellets  
576 of copepods and euphausiids (range: 1.3-2.2%) sampled in the Subantarctic zone of the SO  
577 (Mayzaud et al., 2007). This could indicate that a fraction of the SPOM sampled during the post-  
578 bloom period was composed of fecal material excreted by zooplankton, which increased with  
579 depth. Moreover, and as outlined in the PCA, the PUFA 22:5n-3 was clearly associated with long  
580 chain MUFA. Indeed, a similar linear increase of 22:5n-3 with depth ( $R^2=0.79$ ,  $p<0.001$ ) was  
581 reported suggesting a common origin. This PUFA can be formed from the 20:5n-3 through  
582 elongation and could be a synthesis intermediate of 22:6n-3 in crustaceans including copepods  
583 (Kabeya et al., 2021; Monroig and Kabeya, 2018; Nielsen et al., 2019). This PUFA was found in  
584 three zooplankton species *Themisto libellula*, *Calanus marshallae/glacialis* (*Calanus spp.*), and  
585 *Thysanoessa raschii* collected in the Bering Sea and ranged between 0.5-0.6% of TFA (Wang et al.,  
586 2015). As part of MOBYDICK, 22:5n-3 accounted on average for 0.77% of total phospholipids  
587 extracted from 70 zooplankton samples (Puccinelli E. in prep) and its abundance in SPOM reached  
588  $1.3\pm 0.1\%$  on average in the 150-300m depth interval.

### 589 **Origin and fate of 20:5n-3**

590 Synthesis of the LC-PUFA 20:5n-3 is assumed to occur essentially via autotrophy and can be  
591 performed by several phytoplankton phyla such as Diatoms, ochrophytes, haptophytes, and  
592 rhodophytes (see Table S1 for details). In the ocean and in particular the SO, the main suppliers  
593 of 20:5n-3 are generally considered to be diatoms (Ericson et al., 2018; Hellesey et al., 2020). In  
594 the case of MOBYDICK and for the Kerguelen area, this assumption seems to be true. 20:5n-3 was  
595 clearly associated with diatom C16 PUFA in the PCA and furthermore in the ML, proportions of  
596 20:5n-3 appeared tightly related to C16 PUFA through a positive linear correlation ( $R^2=0.945$ ,

597  $p < 0.05$ ) regardless of stations. Abundance of 20:5n-3 accounted for a significant contribution to  
598 TFA in the ML ( $13 \pm 2\%$  on average), highest proportions were found at station M4 ( $15 \pm 2\%$  on  
599 average), significantly ( $p < 0.001$ ) different in comparison to stations M2 and M3 ( $12 \pm 1\%$  on average  
600 at both stations) and to a lesser extent to station M1 ( $14\%$  on average). Measured proportions in  
601 1-50  $\mu\text{m}$  SPOM were comparable to literature data for diatoms ranging from 15 to 20% of 20:5n-3  
602 (see details in Table S1). However, the literature range corresponds to diatoms monoculture and  
603 differs from the MOBYDICK situation. Because diatoms were a minor group in the ML (18–40%  
604 of total Chl-a) (Irion et al., 2020), proportions of 20:5n-3 in SPOM are expected to be lower  
605 compared to literature data if only provided by diatoms. Based on the highly significant linear  
606 relationship ( $p < 0.001$ ) between 20:5n-3 and diatoms C16 PUFA in the ML, we calculated a 20:5n-  
607 3/C16 PUFA ratio of  $3.6 \pm 0.2$ , which appeared to be  $\sim 2$  fold higher compared to literature data  
608 (average of 1.5 calculated from Table S1). This difference suggests that the diatom community of  
609 the Kerguelen region, both on and off-Plateau, had higher proportions of 20:5n-3 than cultured  
610 diatoms. Such a case was previously reported by Vaezi et al. (2013), who found that the cold-water  
611 *Fragilariopsis cylindrus* was able to produce up to 31.4% of 20:5n-3, approximately twice the  
612 average value usually found in the literature (Table S1). Additionally, we reported a significant  
613 negative regression between 20:5n-3 and temperature in the ML ( $R^2 = 44\%$ ,  $p < 0.01$ ). Indeed, 20:5n-  
614 3 proportion was the highest at station M4 where temperature was the lowest. Based on literature  
615 data, Hixson and Arts (2016) established a negative linear regression between temperature and  
616 20:5n-3 in diatoms. Thus, it can be speculated that the high 20:5n-3 content in diatoms of the  
617 Kerguelen region partially reflected homeoviscous adaptation to low temperatures.

618 In the upper mesopelagic, relative proportions of 20:5n-3 increased, reaching  $17 \pm 2\%$  in the  
619 MLD-150m zone and  $19 \pm 3\%$  in the 150-300m zone, and were significantly ( $p < 0.001$ ) different  
620 compared to the ML. Linear relationship between C16 PUFA and 20:5n-3 was still significant  
621 below the MLD ( $R^2 = 0.487$ ,  $p < 0.001$ ) with an average ratio 20:5n-3/C16 PUFA of 1.3. This ratio was  
622 lower than in the ML and closer to the literature value (1.5, calculated from Table S1). This  
623 suggests that the vertical distribution of 20:5n-3 was still mostly determined by the fate of diatoms  
624 in the mesopelagic with no significant difference among stations. Accordingly, trends in 20:5n-3  
625 proportions were likely a result of the increasing proportion of diatoms observed below the MLD.



626 However, according to the low 20:5n-3/C16 PUFA ratio, the FA composition of diatoms has  
627 probably been modified with depth, and this could be related to several factors. For instance,  
628 diatoms assemblages exhibited consistent variations according to stations and depth (Lafond et  
629 al., 2020). This was particularly the case at the plateau station M2, where in the ML diatom biomass  
630 was dominated by the large centric *Corethron inerme* (59 to 83% of diatom biomass), while below  
631 the ML, *Eucampia antarctica* at the first visit (M2-1), and *Chaetoceros* resting spores and *Chaetoceros*  
632 *spp.* at the following visits (M2-2 and M2-3) were the dominant contributors (Lafond et al., 2020).  
633 At other stations, the community structure was more diverse and included *Actinocyclus spp.*,  
634 *Thalassiosira spp.*, *Chaetoceros atlanticus* and *Chaetoceros* resting spores (Lafond et al., 2020).  
635 Considering that the amount of 20:5n-3 produced varies at a species level, diatom diversity were  
636 thus likely to influence the FA content available. Other factors could also involve the physiological  
637 state of diatom cells, which greatly evolved in the upper mesopelagic zone as a consequence of  
638 grazing, parasitic infection and mortality (Lafond et al., 2020; Sassenhagen et al., 2020). Indeed,  
639 abundance of detrital cells in the form of empty, broken or crushed frustules rapidly increased  
640 below the ML and this was particularly pronounced at off-plateau stations M1, M3, and M4 in  
641 comparison to on-plateau station M2, where up to 48% of cells were still intact between 175-200  
642 m depth (Lafond et al., 2020). Finally, the relative proportions of 20:5n-3 as well as 20:5n-3/C16  
643 PUFA ratio could have been modified as a result of the increasing proportion of fecal material  
644 observed in the mesopelagic zone. The FA composition of fecal pellets reflects the composition of  
645 the zooplankton organism (Hamm et al., 2001; Mayzaud et al., 2007) and can contain high and  
646 variable proportions of 20:5n-3 (up to 25-30%) as observed in krill (Hellessey et al., 2020), with  
647 substantial differences in comparison to the phytoplankton food source.

#### 648 **Origin and fate of 22:6n-3**

649 The LC-PUFA 22:6n-3 is also produced at the basis of the marine food web by unicellular  
650 eukaryotic organisms (Dalsgaard et al., 2003; Parrish, 2013) and it is produced by several  
651 phytoplankton taxa (as detailed in Table S1). The taxa with the highest 22:6n-3 proportions, grown  
652 in autotrophic conditions, include Dinophyceae (17.5% in average), Coccolithophyceae (11.6% in  
653 average) and Prymnesiophyceae (10.7% in average) and exclude diatoms, which produce only

654 small amounts of 22:6n-3 (<2.5%). In this study and considering first the ML, where autotrophic  
655 production was likely dominant, the 1-50  $\mu\text{m}$  SPOM was found to be rich in 22:6n-3 at all stations  
656 representing  $18\pm 1\%$  on average. The highest proportions were found off the Plateau at stations  
657 M1 and M3, with 19.1% and  $19.5\pm 0.5\%$ , respectively, which were slightly higher to the upstream  
658 station M4 ( $17.0\pm 1.0\%$  in average) and the on-plateau station M2 ( $16.4\pm 0.8\%$  in average).  
659 According to the composition of the post-bloom phytoplankton community (Irion et al., 2021,  
660 2020), it can be hypothesized that Prymnesiophyceae, as the dominant group, were the most  
661 important suppliers of 22:6n-3 in the ML both on and off-Plateau. Autotrophic dinoflagellates  
662 (Dinophyceae), which are generally considered as important producers in the SO (Dalsgaard et  
663 al., 2003; Ericson et al., 2018; Falk-Petersen et al., 2000) were likely marginally involved as their  
664 contribution to the phytoplankton community was very modest (Irion et al., 2020). In addition to  
665 Prymnesiophyceae, contribution of pico-phytoplankton (Prasinophyceae) to 22:6n-3 proportions  
666 was also probable, especially on the Plateau where they were more abundant. Prasinophyceae are  
667 able to produce 22:6n-3 but in much lower proportions (7.0%) compared to Prymnesiophyceae. In  
668 order to test the assumption that Prasinophyceae and Prymnesiophyceae contributed to 22:6n-3  
669 production, we considered the relationship between 22:6n-3 and 18:3n-3 in the ML. 22:6n-3  
670 appeared strongly related to 18:3n-3 only at off plateau stations (M1, M3, and M4) through a  
671 positive correlation ( $R^2=0.907$ ,  $p<0.05$ ). The slope of the regression corresponding to the averaged  
672 22:6n-3/18:3n-3 ratio found in off Plateau SPOM was 7.2 and was substantially higher than the  
673 ratio deduced from literature data for Prymnesiophyceae (2.1) and Prasinophyceae (0.6). The fact  
674 that the SPOM 22:6n-3/18:3n-3 ratio was closer to that of Prymnesiophyceae may indicate that  
675 22:6n-3 was at least partially produced by Prymnesiophyceae rather than Prasinophyceae at these  
676 off-Plateau stations. The situation was markedly different on the Plateau at station M2, where  
677 22:6n-3 and 18:3n-3 were negatively related but not significantly, suggesting a more complex  
678 pattern. The observed trend in SPOM could be consistent with a multiple origin of 22:6n-3 with  
679 some from Prymnesiophyceae having a high 22:6n-3/18:3n-3 ratio and some from Prasinophyceae  
680 having a low 22:6n-3/18:3n-3 ratio. Different contributions of Prymnesiophyceae and  
681 Prasinophyceae on and off Plateau seemed to be confirmed by the decrease of 22:6n-3/18:3n-3 ratio  
682 in SPOM observed at station M2 according to visits (7.9 at M2-1, 5.7 at M2-2, and 5.1 at M2-3)

683 which fits with the increasing proportion of Prasinophyceae deduced from pigments and reaching  
684 up to 16% of total Chl-a at M2-3 (Irion et al., 2020).

685 Considering Prymnesiophyceae, it is worth noting that this group was mostly composed by a  
686 single species, *Phaeocystis antarctica* (Irion et al., 2020), whose FA profile remains to be  
687 documented. As a comparison, data available for other species of *Phaeocystis* (*pouchetti* and *globosa*)  
688 exhibit contrasted FA profiles, with some studies reporting only trace or minor amounts of 22:6n-  
689 3 (maximum 6.2% TFA) (Claustre et al., 1990; Nichols et al., 1991; Skerratt et al., 1995), while others  
690 reported much higher proportions, up to 15% (Hamm et al., 2001; Sargent et al., 1985; Virtue et  
691 al., 1993). In our case, *Phaeocystis antarctica* was necessarily high in 22:6n-3 to match the high  
692 proportions observed in the ML (18±1% on average) and further work is needed to confirm this  
693 hypothesis.

694 In addition to autotrophic origin, trophic upgrading by heterotrophic protists such as  
695 dinoflagellates and ciliates, which composed the microzooplankton, may also have contributed to  
696 the observed 22:6n-3 content in the ML (Broglia et al., 2003; Lund et al., 2008). Heterotrophic  
697 dinoflagellates and ciliates were documented at both on and off-plateau stations and ranged  
698 between 0.29 and 2.3 10<sup>3</sup> cells L<sup>-1</sup> (Christaki et al., 2021). Dinoflagellates, dominated by  
699 *Gymnodinium* sp., were more abundant than ciliates at all stations and grazed actively on  
700 phytoplankton with grazing rates exceeding phytoplankton growth rates (Christaki et al., 2021).  
701 Heterotrophic dinoflagellates can contain especially high proportions of 22:6n-3, >40% as reported  
702 by Lim et al. (2020) for *Gymnodinium smaydae*, and up to >50% for *Crypthecodinium cohnii* (Jiang  
703 and Chen, 2000). Interestingly for this latter species, 22:6n-3 proportions were dependent on  
704 cultivation temperature, increasing from 17% to 53% with decreasing temperature from 30°C to  
705 15°C (Jiang and Chen, 2000). Considering the low temperature environment of the Kerguelen  
706 region, this may have favored high 22:6n-3 proportions in heterotrophic protists. Another  
707 characteristic of heterotrophic dinoflagellates is their low content in chloroplastic n-3 C18 PUFA  
708 and as a consequence, their high 22:6n-3/n-3 C18 PUFA ratio (Lim et al., 2020; Lund et al., 2008).  
709 As an example, *Gymnodinium smaydae* had a 22:6n-3/18:3n-3 ratio >80, much higher than the ratio  
710 of its prey *Heterocapsa rotundata* of 5.4 (Lim et al., 2020). This may be one explanation for the higher

711 22:6n-3/18:3n-3 ratio found in the ML (7.2) in comparison to autotrophic producers (0.6-2.1) and  
712 could be consistent with a small contribution (<10%) of heterotrophic protists to 22:6n-3 content.

713 Below the ML in the upper mesopelagic, 22:6n-3 concentrations (in  $\mu\text{g.L}^{-1}$ ) decreased  
714 substantially and similarly to TFA concentrations ( $R^2=0.975$ ,  $p<0.05$ ) it was maintained at high  
715 relative proportions with  $17\pm 2\%$  in the MLD-150m depth interval comparable to the ML, and up  
716 to  $20\pm 2\%$  in the 150-300m depth interval, significantly ( $p<0.001$ ) higher in comparison to the upper  
717 layers. This vertical trend was in a marked contrast compared to n-3 C18 PUFA, which decreased  
718 substantially both in concentration and proportion below the ML. This contrasted pattern was  
719 initially outlined by the PCA, where 22:6n-3 was not correlated to any of the three axes and  
720 furthermore, not associated to n-3 C18 PUFA when considering the whole dataset. These different  
721 trends clearly indicate that the fate of 22:6n-3 in the mesopelagic could not be governed by the  
722 vertical transfer of non-diatom phytoplankton which was initially involved in its production in  
723 the ML, and most probably relied on another pathway. This pathway is further supported by the  
724 linear increase of 22:6n-3 to n-3 C18 PUFA ratio according to depth ( $R^2=0.726$ ,  $p<0.0001$ ) indicating  
725 a progressive enrichment of 22:6n-3 compared to n-3 C18 PUFA in SPOM between the surface and  
726 300 m depth. Moreover, this increase correlates fairly well with the heterotrophic FA 22:5n-3  
727 ( $R^2=0.874$ ,  $p<0.0001$ ) and also but to a lesser extent with the MUFA 20:1n-9 ( $R^2=0.551$ ,  $p<0.001$ ).  
728 Variations of 22:5n-3 and 20:1n-9 were attributed earlier in the discussion to an increasing  
729 abundance of fecal material at depth. Accordingly, it can be postulated that 22:6n-3 in SPOM may  
730 be at least partially associated with this fecal material in the upper mesopelagic and thus its fate  
731 may be controlled by interactions with the heterotrophic food web. It is worth noting that  
732 correlations were made regardless of stations, indicating that this potential transfer pathway  
733 seemed to affect both iron-fertilized and HNLC regions. Such an association of 22:6n-3 with fecal  
734 material has already been documented in some regions of the world ocean, including the high  
735 latitude North Pacific where fecal OM and high proportions of 22:6n-3 (range 8-17% of TFA) were  
736 evidenced throughout the water column (Sheridan et al., 2002; Wilson et al., 2010) and also in the  
737 Arctic, where it was demonstrated that 22:6n-3 rich OM (~20% of TFA) derived from *Phaeocystis*  
738 *pouchetii*, was exported below the euphotic zone via krill fecal strings (Hamm et al., 2001).

739 In the present case, the processing of 22:6n-3 from phytoplankton to suspended fecal material  
740 is likely to be complex and could involve a number of heterotrophic intermediates. Among  
741 grazers, mesozooplankton such as copepods and other large size crustacean zooplankton require  
742 to be considered as C20-C22 MUFA are representative of fecal material excreted by these  
743 organisms (Brett et al., 2009; Dalsgaard et al., 2003; Kattner and Hagen, 1995; Mayzaud et al., 2007;  
744 Parrish, 2013; Wilson et al., 2010). Mesozooplankton were documented at all stations and  
745 exhibited significant variations according to sites and visits from 207 ind.m<sup>-3</sup> at M2-1 to 1636  
746 ind.m<sup>-3</sup> at M4-1 (Christaki et al., 2021). Moreover, grazing experiments showed that copepods  
747 were not grazing directly on phytoplankton but were feeding primarily on microzooplankton  
748 controlling their abundance during the post-bloom period (Christaki et al., 2021). This feeding  
749 strategy on heterotrophic protists as well as the selective storage of 22:6n-3 as an essential LC-  
750 PUFA for mesozooplankton (Arendt et al., 2005) may have favoured its accumulation relative to  
751 n-3 C18 PUFA and as a result, impacted the composition of fecal material produced by these  
752 organisms. In addition to mesozooplankton, other grazers such as salps may also be involved in  
753 the channelling of 22:6n-3. Salps are important grazers of small phytoplankton (Moline et al., 2004)  
754 and flagellates (Pakhomov and Hunt, 2017; von Harbou et al., 2011), and produce easily  
755 fragmented tabular fecal pellets (Iversen et al., 2017). Salps population (*Salpa thompsoni*) were  
756 particularly abundant on the Plateau at station M2 and downstream at station M1 representing  
757 up to 40% of total micronekton biomass (Henschke et al., 2021).

#### 758 **Nutritional food value according to production regimes**

759 The nutritional quality of OM as food for higher trophic levels is partly determined by the  
760 abundance of the essential omega-3 LC-PUFA 20:5n-3 and 22:6n-3 and the essential omega-6  
761 20:4n-6. In the present study, LC-PUFA essentially composed of omega-3 LC-PUFA were found  
762 in high proportions, making up 27-44% of TFA, indicating a high nutritional value of SPOM both  
763 in the ML and upper mesopelagic. Following the approach developed by Cañavate et al. (2019),  
764 the NQI of SPOM was also high and ranged from 354 to 403 in the ML and from 333 to 477 in the  
765 MLD-300m depth interval. As a comparison, much lower NQI were obtained in the estuary of  
766 Guadalquivir River, with values always <200 (Cañavate et al., 2021), in the northern part of the

767 Indian sector of the ACC, with NQI of 193 and 242 in surface SPOM of the subtropical and  
768 Subantarctic zone, respectively (Mayzaud et al., 2007), in the Northern Pacific, NQI was <300 in  
769 >51  $\mu\text{m}$  SPOM of the Subarctic area (Wilson et al., 2010) as well as in the coastal Antarctic zone,  
770 NQI was <260 for a diatom sea-ice community of McMurdo Sound (Nichols et al., 1993). This  
771 comparison confirms that quality of SPOM in the Kerguelen area was especially high in the post-  
772 bloom period. High NQI values were observed over the whole area and seemed unrelated to the  
773 contrasted productivity regimes that took place earlier in the season during the bloom. In the ML,  
774 the high nutritional value of SPOM was mainly attributable to the mixed phytoplanktonic  
775 community relying both on diatoms producing 20:5n-3 and small phytoplankton producing  
776 22:6n-3. This mixed production of both LC-PUFA substantially promoted the nutritional quality  
777 of SPOM and offered a valuable source of food for secondary consumers that could be of  
778 importance for the ecosystem functioning.

779 In the upper mesopelagic, LC-PUFA in SPOM were maintained at similar, and even higher,  
780 proportions in comparison to the ML ( $34\pm 2\%$  and  $40\pm 3\%$  on average in the MLD-150m and 150-  
781 300m depth intervals, respectively). Similarly, NQI remained high until 300 m depth (333-477)  
782 even though TFA concentrations in seawater (in  $\mu\text{g}\cdot\text{L}^{-1}$ ) decreased substantially with depth. This  
783 preservation of food quality in the upper mesopelagic was rather unexpected because LC-PUFA  
784 are generally considered as a labile fraction of OM that could be easily degraded/consumed by  
785 heterotrophic organisms during their transfer from the surface (Budge et al., 2006; Conte et al.,  
786 2003, 1995; Wakeham, 1995). This is illustrated by the low NQI values that can be deduced from  
787 sediment trap data, with values between 90 to 180 in trap samples collected between 50-100 m  
788 depth in the North Atlantic (Budge and Parrish, 1998) and NQI <30 in samples collected at 300 m  
789 depth in Breid Bay, Antarctica (Hayakawa et al., 1996). For the Kerguelen region, sediment trap  
790 data obtained on the Plateau at station M2 (Rembauville et al., 2018) do not allow for NQI  
791 estimates, however considering LC-PUFA concentrations per unit mass of OC, ranging from 0.03  
792 to  $3.4 \mu\text{g}\cdot\text{mgOC}^{-1}$  in the trap samples and from 14 to  $44 \mu\text{g}\cdot\text{mgOC}^{-1}$  in our SPOM samples, it  
793 appears that sinking material from traps was of much lower quality than suspended particles  
794 collected on and off the Plateau. In the present case, the high quality of SPOM in the upper  
795 mesopelagic could represent a valuable nutritional option for heterotrophic organisms residing

796 in the part of the water column and feeding on suspended particles such as zooplankton,  
797 coprophages, detritivores, and bacteria.

## 798 E. CONCLUSION

799 In this study, FA distribution including essential LC-PUFA 20:5n-3 and 22:6n-3 was  
800 documented in the upper water column of the Kerguelen Islands Region several weeks after the  
801 bloom period in late summer/early autumn. Our results illustrate the important role of the mixed  
802 phytoplankton community, composed of nano- and pico-size phytoplankton and diatoms, in  
803 providing high proportions of both LC-PUFA to surface waters. Diatoms were identified as the  
804 likely suppliers of 20:5n-3 and small phytoplankton (prymnesiophytes and prasinophytes)  
805 supplying 22:6n-3. Elevated proportions are unrelated to past productivity regimes and were  
806 observed both inside the iron-fertilized area on the Kerguelen Plateau (station M2) and  
807 downstream (station M1), as well as outside in HNLC waters located upstream of the Plateau  
808 (stations M3 and M4). As a consequence, and despite reduced productivity and biomass  
809 abundance during the post-bloom period, phytoplankton-derived OM reveal an especially high  
810 nutritional value that could be relevant for secondary consumers such as micro- and  
811 mesozooplankton.

812 In the upper mesopelagic, both LC-PUFA were maintained at high relative proportions in  
813 SPOM suggesting an efficient and probably fast vertical transfer from the surface. This vertical  
814 transfer seems to proceed via distinct pathways according to LC-PUFA. For 20:5n-3, its fate in the  
815 mesopelagic seems to mainly follow the one of diatoms and could be associated with the export  
816 of large intact cells, diatoms aggregates as well as resting spores. For 22:6n-3, its fate with depth  
817 appears not to be simply related to small phytoplankton but is most likely due to a complex  
818 channeling through the heterotrophic food web resulting in its association with fecal material at  
819 depth. Processing of 22:6n-3 could involve different intermediates such as heterotrophic protists  
820 with dinoflagellates and ciliates (microzooplankton), which probably grazed on small  
821 phytoplankton as well as larger zooplankton organisms such as copepods and salps, which  
822 presumably fed on microzooplankton and produced fecal pellets rich in 22:6n-3. The complex  
823 pathway of 22:6n-3 cannot be easily resolved using only SPOM data and additional evidences are

824 needed to better understand this pattern. Despite these contrasted pathways, FA-based  
825 nutritional quality of SPOM remained especially high in the upper mesopelagic and could  
826 represent an interesting food source for suspension feeders residing there. Finally, these results  
827 provide a detailed framework of FA abundances that can be used in trophic and diet studies  
828 dedicated to the Subantarctic islands.

## 829 F. Acknowledgments

830 We thank B. Quéguiner, the PI of the MOBYDICK project, for providing us the opportunity to  
831 participate to this cruise, the chief scientist I. Obernosterer and the captain and crew of the R/V  
832 Marion Dufresne for their enthusiasm and support aboard during the MOBYDICK–THEMISTO  
833 cruise (<https://doi.org/10.17600/18000403>). This work was supported by the French oceanographic  
834 fleet (“Flotte océanographique française”), the French ANR (“Agence Nationale de la Recherche”,  
835 AAPG 2017 program, MOBYDICK Project number: ANR-17-CE01-0013), the French Research  
836 program of INSU-CNRS LEFE/CYBER (“Les enveloppes fluides et l’environnement” – “Cycles  
837 biogéochimiques, environnement et ressources”). Marine Remize PhD fellowship and studies was  
838 supported by the University of Brest (UBO, France), the Center of Marine Sciences (CMS) of the  
839 University of North Carolina Wilmington (UNCW, USA), the “Laboratoire d’Excellence”  
840 LabexMer (ANR-10-LABX-19), and the Walter-Zellidja grant of the Académie Française. E.  
841 Puccinelli was supported by ISblue project, Interdisciplinary graduate school for the blue planet  
842 (ANR-17-EURE-0015) and co-funded by a grant from the French government under the program  
843 “Investissements d’Avenir”. We would like to thank our colleagues of the Technical Division of  
844 the INSU, Emmanuel de Saint-Leger, Lionel Scouarnec, Lionel Fischen, and Olivier Desprez De  
845 Gesincourt for their logistical and technical help with the in-situ pumps, the colleagues of the UBO  
846 Open Factory for their help in designing and preparing the NITEX membranes, and also to  
847 Stephane Blain and Bernard Quéguiner for their helpful comments and suggestions on the  
848 manuscript.

## 849 G. References



850 Alonso, D.L., Belarbi, E.-H., Rodríguez-Ruiz, J., Segura, C.I., Giménez, A., 1998. Acyl lipids of  
851 three microalgae. *Phytochemistry* 47, 1473–1481. [https://doi.org/10.1016/S0031-9422\(97\)01080-7](https://doi.org/10.1016/S0031-9422(97)01080-7)

852 Arendt, K.E., Jónasdóttir, S.H., Hansen, P.J., Gärtner, S., 2005. Effects of dietary fatty acids on  
853 the reproductive success of the calanoid copepod *Temora longicornis*. *Marine Biology* 146, 513–  
854 530. <https://doi.org/10.1007/s00227-004-1457-9>

855 Armand, L.K., Cornet-Barthaux, V., Mosseri, J., Quéguiner, B., 2008. Late summer diatom  
856 biomass and community structure on and around the naturally iron-fertilised Kerguelen Plateau  
857 in the Southern Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography* 55, 653–676.  
858 <https://doi.org/10.1016/j.dsr2.2007.12.031>

859 Atkinson, A., Shreeve, R.S., Hirst, A.G., Rothery, P., Tarling, G.A., Pond, D.W., Korb, R.E.,  
860 Murphy, E.J., Watkins, J.L., 2006. Natural growth rates in Antarctic krill (*Euphausia superba*): II.  
861 Predictive models based on food, temperature, body length, sex, and maturity stage. *Limnology*  
862 *and Oceanography* 51, 973–987. <https://doi.org/10.4319/lo.2006.51.2.0973>

863 Bec, A., Martin-Creuzburg, D., von Elert, E., 2006. Trophic upgrading of autotrophic  
864 picoplankton by the heterotrophic nanoflagellate *Paraphysomonas* sp. *Limnology and*  
865 *Oceanography* 51, 1699–1707. <https://doi.org/10.4319/lo.2006.51.4.1699>

866 Blain, S., Queguiner, B., Armand, L., Belviso, S., Bombled, B., Bopp, L., Bowie, A., Brunet, C.,  
867 Brussaard, C., Carlotti, F., Christaki, U., Corbiere, A., Durand, I., Ebersbach, F., Fuda, J.-L., Garcia,  
868 N., Gerringa, L., Griffiths, B., Guigue, C., Guillerm, C., Jacquet, S., Jeandel, C., Laan, P., Lefevre,  
869 D., Lo Monaco, C., Malits, A., Mosseri, J., Obernosterer, I., Park, Y.-H., Picheral, M., Pondaven, P.,  
870 Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scouarnec, L., Souhaut, M., Thuiller, D.,  
871 Timmermans, K., Trull, T., Uitz, J., van Beek, P., Veldhuis, M., Vincent, D., Viollier, E., Vong, L.,  
872 Wagener, T., 2007. Effect of natural iron fertilization on carbon sequestration in the Southern  
873 Ocean. *Nature* 446, 1070–1074. <https://doi.org/10.1038/nature05700>

874 Blain, S., Rembauville, M., Crispi, O., Obernosterer, I., 2021. Synchronized autonomous  
875 sampling reveals coupled pulses of biomass and export of morphologically different diatoms in  
876 the Southern Ocean. *Limnology and Oceanography* 66, 753–764. <https://doi.org/10.1002/lno.11638>

877 Blain, S., Sarthou, G., Laan, P., 2008. Distribution of dissolved iron during the natural iron-  
878 fertilization experiment KEOPS (Kerguelen Plateau, Southern Ocean). *Deep Sea Research Part II:  
879 Topical Studies in Oceanography* 55, 594–605. <https://doi.org/10.1016/j.dsr2.2007.12.028>

880 Brett, M.T., Müller-Navarra, D., 1997. The role of highly unsaturated fatty acids in aquatic food  
881 web processes. *Freshwater Biology* 38, 483–499. <https://doi.org/10.1046/j.1365-2427.1997.00220.x>

882 Brett, M.T., Müller-Navarra, D.C., Persson, J., 2009. Crustacean zooplankton fatty acid  
883 composition, in: Kainz, M., Brett, M., Arts, M. (Eds.), *Lipids in Aquatic Ecosystems*. Springer, New  
884 York, NY.

885 Broglio, E., Jónasdóttir, S.H., Calbet, A., Jakobsen, H.H., Saiz, E., 2003. Effect of heterotrophic  
886 versus autotrophic food on feeding and reproduction of the calanoid copepod *Acartia tonsa*:  
887 relationship with prey fatty acid composition. *Aquat Microb Ecol* 31, 267–278.

888 Brooks, P.D., Geilmann, H., Werner, R.A., Brand, W., 2003. Improved precision of coupled <sup>13</sup>C  
889 and <sup>15</sup>N measurements from single samples using an elemental analyser. *Rapid Communications  
890 in Mass Spectrometry* 12, 1924–1926. <https://doi.org/10.1002/rcm.1134>

891 Budge, S.M., Iverson, S.J., Koopman, H.N., 2006. Studying trophic ecology in marine  
892 ecosystems using fatty acids: a primer on analysis and interpretation. *Marine Mammal Science* 22,  
893 759–801.

894 Budge, S.M., Parrish, C.C., 1998. Lipid biogeochemistry of plankton, settling matter and  
895 sediments in Trinity Bay, Newfoundland. II. Fatty acids. *Organic Geochemistry* 29, 1547–1559.  
896 [https://doi.org/10.1016/S0146-6380\(98\)00177-6](https://doi.org/10.1016/S0146-6380(98)00177-6)

897 Cañavate, J.P., 2019. Advancing assessment of marine phytoplankton community structure  
898 and nutritional value from fatty acid profiles of cultured microalgae. *Reviews in Aquaculture* 11,  
899 527–549. <https://doi.org/10.1111/raq.12244>

900 Cañavate, J.-P., van Bergeijk, S., González-Ortegón, E., Vílas, C., 2021. Contrasting fatty acids  
901 with other indicators to assess nutritional status of suspended particulate organic matter in a

902 turbid estuary. *Estuarine, Coastal and Shelf Science* 254, 107329.  
903 <https://doi.org/10.1016/j.ecss.2021.107329>

904 Cavagna, A.J., Fripiat, F., Elskens, M., Mangion, P., Chirurgien, L., Closset, I., Lasbleiz, M.,  
905 Florez-Leiva, L., Cardinal, D., Leblanc, K., Fernandez, C., Lefèvre, D., Oriol, L., Blain, S.,  
906 Quéguiner, B., Dehairs, F., 2015. Production regime and associated N cycling in the vicinity of  
907 Kerguelen Island, Southern Ocean. *Biogeosciences* 12, 6515–6528. [https://doi.org/10.5194/bg-12-](https://doi.org/10.5194/bg-12-6515-2015)  
908 6515-2015

909 Christaki, U., Gueneugues, A., Liu, Y., Blain, S., Catala, P., Colombet, J., Debeljak, P., Jardillier,  
910 L., Irion, S., Planchon, F., Sassenhagen, I., Sime-Ngando, T., Obernosterer, I., 2020. Seasonal  
911 microbial food web dynamics in contrasting Southern Ocean productivity regimes. *Limnology*  
912 and *Oceanography* n/a. <https://doi.org/10.1002/lno.11591>

913 Christaki, U., Skouroliakou, I.-D., Delegrange, A., Irion, S., Courcot, L., Jardillier, L.,  
914 Sassenhagen, I., 2021. Microzooplankton diversity and potential role in carbon cycling of  
915 contrasting Southern Ocean productivity regimes. *Journal of Marine Systems* 219, 103531.  
916 <https://doi.org/10.1016/j.jmarsys.2021.103531>

917 Claustre, H., Poulet, S.A., Williams, R., Marty, J.C., Coombs, S., Mlih, F.B., Hapette, A.M.,  
918 Jezequel-Martin, V., 1990. A biochemical investigation of a *Phaeocystis* sp. bloom in the Irish Sea.  
919 *Journal of the Marine Biological Association of the United Kingdom* 70, 197–207.  
920 <https://doi.org/10.1017/S0025315400034317>

921 Constable, A.J., Melbourne-Thomas, J., Corney, S.P., Arrigo, K.R., Barbraud, C., Barnes,  
922 D.K.A., Bindoff, N.L., Boyd, P.W., Brandt, A., Costa, D.P., Davidson, A.T., Ducklow, H.W.,  
923 Emmerson, L., Fukuchi, M., Gutt, J., Hindell, M.A., Hofmann, E.E., Hosie, G.W., Iida, T., Jacob, S.,  
924 Johnston, N.M., Kawaguchi, S., Kokubun, N., Koubbi, P., Lea, M.-A., Makhado, A., Massom, R.A.,  
925 Meiners, K., Meredith, M.P., Murphy, E.J., Nicol, S., Reid, K., Richerson, K., Riddle, M.J., Rintoul,  
926 S.R., Smith Jr, W.O., Southwell, C., Stark, J.S., Sumner, M., Swadling, K.M., Takahashi, K.T.,  
927 Trathan, P.N., Welsford, D.C., Weimerskirch, H., Westwood, K.J., Wienecke, B.C., Wolf-Gladrow,  
928 D., Wright, S.W., Xavier, J.C., Ziegler, P., 2014. Climate change and Southern Ocean ecosystems I:

929 how changes in physical habitats directly affect marine biota. *Global Change Biology* 20, 3004–  
930 3025. <https://doi.org/10.1111/gcb.12623>

931 Conte, M.H., Dickey, T.D., Weber, J.C., Johnson, R.J., Knap, A.H., 2003. Transient physical  
932 forcing of pulsed export of bioreactive material to the deep Sargasso Sea. *Deep Sea Research Part*  
933 *I: Oceanographic Research Papers* 50, 1157–1187. [https://doi.org/10.1016/S0967-0637\(03\)00141-9](https://doi.org/10.1016/S0967-0637(03)00141-9)

934 Conte, M.H., Eglinton, G., Madeira, L.A.S., Rabouille, C., Labeyrie, L., Mudge, S., Eglinton,  
935 G., Elderfield, H., Whitfield, M., Williams, P.J.Le.B., 1995. Origin and fate of organic biomarker  
936 compounds in the water column and sediments of the eastern North Atlantic. *Philosophical*  
937 *Transactions of the Royal Society of London. Series B: Biological Sciences* 348, 169–178.  
938 <https://doi.org/10.1098/rstb.1995.0059>

939 Dalsgaard, J., St. John, M., Kattner, G., Müller-Navarra, D., Hagen, W., 2003. Fatty acid trophic  
940 markers in the pelagic marine environment, in: *Advances in Marine Biology*. Academic Press, pp.  
941 225–340. [https://doi.org/10.1016/S0065-2881\(03\)46005-7](https://doi.org/10.1016/S0065-2881(03)46005-7)

942 de Baar, H.J.W., de Jong, J.T.M., Bakker, D.C.E., Loscher, B.M., Veth, C., Bathmann, U.,  
943 Smetacek, V., 1995. Importance of iron for plankton blooms and carbon dioxide drawdown in the  
944 Southern Ocean. *Nature* 373, 412–415. <https://doi.org/10.1038/373412a0>

945 Deppeler, S.L., Davidson, A.T., 2017. Southern Ocean Phytoplankton in a Changing Climate.  
946 *Frontiers in Marine Science* 4. <https://doi.org/10.3389/fmars.2017.00040>

947 Dunstan, G.A., Volkman, J.K., Barrett, S.M., Leroi, J.-M., Jeffrey, S.W., 1993. Essential  
948 polyunsaturated fatty acids from 14 species of diatom (Bacillariophyceae). *Phytochemistry* 35,  
949 155–161. [https://doi.org/10.1016/S0031-9422\(00\)90525-9](https://doi.org/10.1016/S0031-9422(00)90525-9)

950 Ebersbach, F., Trull, T.W., 2008. Sinking particle properties from polyacrylamide gels during  
951 the Kerguelen Ocean and Plateau compared Study (KEOPS): Zooplankton control of carbon  
952 export in an area of persistent natural iron inputs in the Southern Ocean. *Limnol. Oceanogr.* 53,  
953 212–224. <https://doi.org/10.4319/lo.2008.53.1.0212>

954 El-Sayed, S.Z., 1988. Productivity of the southern ocean: a closer look. *Comparative*  
955 *Biochemistry and Physiology Part B: Comparative Biochemistry* 90, 489–498.  
956 [https://doi.org/10.1016/0305-0491\(88\)90287-8](https://doi.org/10.1016/0305-0491(88)90287-8)

957 Ericson, J.A., Hellessey, N., Nichols, P.D., Kawaguchi, S., Nicol, S., Hoem, N., Virtue, P., 2018.  
958 Seasonal and interannual variations in the fatty acid composition of adult *Euphausia superba*  
959 Dana, 1850 (Euphausiacea) samples derived from the Scotia Sea krill fishery. *Journal of*  
960 *Crustacean Biology* 38, 662–672. <https://doi.org/10.1093/jcbiol/ruy032>

961 Evans, C., Brussaard, C.P.D., 2012. Viral lysis and microzooplankton grazing of  
962 phytoplankton throughout the Southern Ocean. *Limnology and Oceanography* 57, 1826–1837.  
963 <https://doi.org/10.4319/lo.2012.57.6.1826>

964 Fahl, K., Kattner, G., 1993. Lipid content and fatty acid composition of algal communities in  
965 sea-ice and water from the Weddell Sea (Antarctica). *Polar Biology* 13, 405–409.  
966 <https://doi.org/10.1007/BF01681982>

967 Falk-Petersen, S., Hagen, W., Kattner, G., Clarke, A., Sargent, J., 2000. Lipids, trophic  
968 relationships, and biodiversity in Arctic and Antarctic krill. *Canadian Journal of Fisheries and*  
969 *Aquatic Sciences* 57, 178–191. <https://doi.org/10.1139/f00-194>

970 Fiala, M., Kopczynska, E.E., Jeandel, C., Oriol, L., Vétion, G., 1998. Seasonal and interannual  
971 variability of size-fractionated phytoplankton biomass and community structure at station Kerfix,  
972 off the Kerguelen Islands, Antarctica. *Journal of Plankton Research* 20, 1341–1356.  
973 <https://doi.org/10.1093/plankt/20.7.1341>

974 Gillan, F.T., McFadden, G.I., Wetherbee, R., Johns, R.B., 1981. Sterols and fatty acids of an  
975 Antarctic sea ice diatoms, *Stauroneis amphioxys*. *Phytochemistry* 20, 1935–1937.

976 Guo, F., Bunn, S.E., Brett, M.T., Kainz, M., 2017. Polyunsaturated fatty acids in stream food  
977 webs - high dissimilarity among producers and consumers. *Freshwater Biology* 62, 1325–1334.  
978 <https://doi.org/10.1111/fwb.12956>

979 Guschina, I.A., Harwood, J.L., 2006. Lipids and lipid metabolism in eukaryotic algae. Progress  
980 in Lipid Research 45, 160–186. <https://doi.org/10.1016/j.plipres.2006.01.001>

981 Hamm, C., Reigstad, M., Riser, C.W., Mühlebach, Anneke, Wassmann, P., 2001. On the trophic  
982 fate of *Phaeocystis pouchetii*. VII. Sterols and fatty acids reveal sedimentation of *P. pouchetii*-  
983 derived organic matter via krill fecal strings. Mar Ecol Prog Ser 209, 55–69.

984 Hayakawa, K., Handa, N., Ikuta, N., Fukuchi, M., 1996. Downward fluxes of fatty acids and  
985 hydrocarbons during a phytoplankton bloom in the austral summer in Breid Bay, Antarctica.  
986 Organic Geochemistry 24, 511–521. [https://doi.org/10.1016/0146-6380\(96\)00047-2](https://doi.org/10.1016/0146-6380(96)00047-2)

987 Hellessey, N., Johnson, R., Ericson, J.A., Nichols, P.D., Kawaguchi, S., Nicol, S., Hoem, N.,  
988 Virtue, P., 2020. Antarctic Krill Lipid and Fatty acid Content Variability is Associated to Satellite  
989 Derived Chlorophyll a and Sea Surface Temperatures. Scientific Reports 10, 6060.  
990 <https://doi.org/10.1038/s41598-020-62800-7>

991 Hernando, M., Schloss, I.R., Almandoz, G.O., Malanga, G., Varela, D.E., De Troch, M., 2018.  
992 Combined effects of temperature and salinity on fatty acid content and lipid damage in Antarctic  
993 phytoplankton. Journal of Experimental Marine Biology and Ecology 503, 120–128.  
994 <https://doi.org/10.1016/j.jembe.2018.03.004>

995 Hixson, S.M., Arts, M.T., 2016. Climate warming is predicted to reduce omega-3, long-chain,  
996 polyunsaturated fatty acid production in phytoplankton. Global Change Biology 22, 2744–2755.  
997 <https://doi.org/10.1111/gcb.13295>

998 Irion, S., Christaki, U., Berthelot, H., L’Helguen, S., Jardillier, L., 2021. Small phytoplankton  
999 contribute greatly to CO<sub>2</sub>-fixation after the diatom bloom in the Southern Ocean. The ISME  
1000 Journal. <https://doi.org/10.1038/s41396-021-00915-z>

1001 Irion, S., Jardillier, L., Sassenhagen, I., Christaki, U., 2020. Marked spatiotemporal variations  
1002 in small phytoplankton structure in contrasted waters of the Southern Ocean (Kerguelen area).  
1003 Limnology and Oceanography. <https://doi.org/10.1002/lno.11555>

1004 Iversen, M.H., Pakhomov, E.A., Hunt, B.P.V., van der Jagt, H., Wolf-Gladrow, D., Klaas, C.,  
1005 2017. Sinkers or floaters? Contribution from salp pellets to the export flux during a large bloom  
1006 event in the Southern Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography* 138,  
1007 116–125. <https://doi.org/10.1016/j.dsr2.2016.12.004>

1008 Jiang, Y., Chen, F., 2000. Effects of medium glucose concentration and pH on docosahexaenoic  
1009 acid content of heterotrophic *Cryptocodinium cohnii*. *Process Biochemistry* 35, 1205–1209.  
1010 [https://doi.org/10.1016/S0032-9592\(00\)00163-1](https://doi.org/10.1016/S0032-9592(00)00163-1)

1011 Jónasdóttir, S.H., 2019. Fatty Acid Profiles and Production in Marine Phytoplankton. *Marine*  
1012 *Drugs* 17, 151.

1013 Kabeya, N., Ogino, M., Ushio, H., Haga, Y., Satoh, S., Navarro, J.C., Monroig, Ó., 2021. A  
1014 complete enzymatic capacity for biosynthesis of docosahexaenoic acid (DHA, 22 : 6n-3) exists in  
1015 the marine Harpacticoida copepod *Tigriopus californicus*. *Open Biology* 11, 200402.  
1016 <https://doi.org/10.1098/rsob.200402>

1017 Kattner, G., Hagen, W., 1995. Polar herbivorous copepods – different pathways in lipid  
1018 biosynthesis. *ICES Journal of Marine Science* 52, 329–335. [https://doi.org/10.1016/1054-](https://doi.org/10.1016/1054-3139(95)80048-4)  
1019 [3139\(95\)80048-4](https://doi.org/10.1016/1054-3139(95)80048-4)

1020 Klein-Breteler, W.C.M., Schogt, N., Baas, M., Schouten, S., Kraay, G.W., 1999. Trophic  
1021 upgrading of food quality by protozoans enhancing copepod growth: role of essential lipids.  
1022 *Marine Biology* 135, 191–198. <https://doi.org/10.1007/s002270050616>

1023 Koczyńska, E.E., Fiala, M., Jeandel, C., 1998. Annual and interannual variability in  
1024 phytoplankton at a permanent station off Kerguelen Islands, Southern Ocean. *Polar Biology* 20,  
1025 342–351. <https://doi.org/10.1007/s003000050312>

1026 Korb, R.E., Whitehouse, M., 2004. Contrasting primary production regimes around South  
1027 Georgia, Southern Ocean: large blooms versus high nutrient, low chlorophyll waters. *Deep Sea*  
1028 *Research Part I: Oceanographic Research Papers* 51, 721–738.  
1029 <https://doi.org/10.1016/j.dsr.2004.02.006>

1030 Lafond, A., Leblanc, K., Legras, J., Cornet, V., Quéguiner, B., 2020. The structure of diatom  
1031 communities constrains biogeochemical properties in surface waters of the Southern Ocean  
1032 (Kerguelen Plateau). *Journal of Marine Systems* 212, 103458.  
1033 <https://doi.org/10.1016/j.jmarsys.2020.103458>

1034 Lasbleiz, M., Leblanc, K., Armand, L.K., Christaki, U., Georges, C., Obernosterer, I.,  
1035 Quéguiner, B., 2016. Composition of diatom communities and their contribution to plankton  
1036 biomass in the naturally iron-fertilized region of Kerguelen in the Southern Ocean. *FEMS*  
1037 *Microbiology Ecology* 92, fiw171.

1038 Laurenceau-Cornec, E.C., Trull, T.W., Davies, D.M., Bray, S.G., Doran, J., Planchon, F., Carlotti,  
1039 F., Jouandet, M.P., Cavagna, A.J., Waite, A.M., Blain, S., 2015. The relative importance of  
1040 phytoplankton aggregates and zooplankton fecal pellets to carbon export: insights from free-  
1041 drifting sediment trap deployments in naturally iron-fertilised waters near the Kerguelen Plateau.  
1042 *Biogeosciences* 12, 1007–1027. <https://doi.org/10.5194/bg-12-1007-2015>

1043 Lim, A.S., Jeong, H.J., You, J.H., Park, S.A., 2020. Semi-continuous cultivation of the  
1044 mixotrophic dinoflagellate *Gymnodinium smaydae*, a new promising microalga for omega-3  
1045 production. *Algae* 35, 277–292. <https://doi.org/10.4490/algae.2020.35.9.2>

1046 Liu, Y., Blain, S., Crispi, O., Rembauville, M., Obernosterer, I., 2020. Seasonal dynamics of  
1047 prokaryotes and their associations with diatoms in the Southern Ocean as revealed by an  
1048 autonomous sampler. *Environmental Microbiology* 22, 3968–3984. [https://doi.org/10.1111/1462-](https://doi.org/10.1111/1462-2920.15184)  
1049 [2920.15184](https://doi.org/10.1111/1462-2920.15184)

1050 Lund, E.D., Chu, F.-L.E., Harvey, E., Adlof, R., 2008. Mechanism(s) of long chain n-3 essential  
1051 fatty acid production in two species of heterotrophic protists: *Oxyrrhis marina* and *Gyrodinium*  
1052 *dominans*. *Marine Biology* 155, 23–36. <https://doi.org/10.1007/s00227-008-1003-2>

1053 Martin, J.H., 1990. Glacial-interglacial CO<sub>2</sub> change: The Iron Hypothesis. *Paleoceanography* 5,  
1054 1–13. <https://doi.org/10.1029/PA005i001p00001>



1055 Martin-Creuzburg, D., von Elert, E., Hoffmann, K.H., 2008. Nutritional constraints at the  
1056 cyanobacteria—*Daphnia magna* interface: The role of sterols. *Limnology and Oceanography* 53,  
1057 456–468. <https://doi.org/10.4319/lo.2008.53.2.0456>

1058 Mathieu-Resuge, M., Schaal, G., Kraffe, E., Corvaisier, R., Lebeau, O., Lluch-Costa, S.E.,  
1059 Salgado García, R.L., Kainz, M.J., Le Grand, F., 2019. Different particle sources in a bivalve species  
1060 of a coastal lagoon: evidence from stable isotopes, fatty acids, and compound-specific stable  
1061 isotopes. *Marine Biology* 166, 89–101.

1062 Mayzaud, P., Laureillard, J., Merien, D., Brinis, A., Godard, C., Razouls, S., Labat, J. –P, 2007.  
1063 Zooplankton nutrition, storage and fecal lipid composition in different water masses associated  
1064 with the Agulhas and Subtropical Fronts. *Marine Chemistry* 107, 202–213.  
1065 <https://doi.org/10.1016/j.marchem.2007.07.001>

1066 Mayzaud, P., Tirelli, V., Errhif, A., Labat, J.P., Razouls, S., Perissinotto, R., 2002. Carbon intake  
1067 by zooplankton. Importance and role of zooplankton grazing in the Indian sector of the Southern  
1068 Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography* 49, 3169–3187.  
1069 [https://doi.org/10.1016/S0967-0645\(02\)00077-2](https://doi.org/10.1016/S0967-0645(02)00077-2)

1070 Mitani, E., Nakayama, F., Matsuwaki, I., Ichi, I., Kawabata, A., Kawachi, M., Kato, M., 2017.  
1071 Fatty acid composition profiles of 235 strains of three microalgal divisions within the NIES  
1072 Microbial Culture Collection. *Microb. Resour. Syst* 33, 1929.

1073 Moline, M.A., Claustre, H., Frazer, T.K., Schofield, O., Vernet, M., 2004. Alteration of the food  
1074 web along the Antarctic Peninsula in response to a regional warming trend. *Global Change*  
1075 *Biology* 10, 1973–1980. <https://doi.org/10.1111/j.1365-2486.2004.00825.x>

1076 Mongin, M., Molina, E., Trull, T.W., 2008. Seasonality and scale of the Kerguelen plateau  
1077 phytoplankton bloom: A remote sensing and modeling analysis of the influence of natural iron  
1078 fertilization in the Southern Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography*  
1079 55, 880–892. <https://doi.org/10.1016/j.dsr2.2007.12.039>

1080 Monroig, Ó., Kabeya, N., 2018. Desaturases and elongases involved in polyunsaturated fatty  
1081 acid biosynthesis in aquatic invertebrates: a comprehensive review. *Fisheries Science* 84, 911–928.  
1082 <https://doi.org/10.1007/s12562-018-1254-x>

1083 Müller-Navarra, D.C., Brett, M.T., Liston, A.M., Goldman, C.R., 2000. A highly unsaturated  
1084 fatty acid predicts carbon transfer between primary producers and consumers. *Nature* 403, 74–77.  
1085 <https://doi.org/10.1038/47469>

1086 Nichols, D.S., Nichols, P.D., Sullivan, C.W., 1993. Fatty acid, sterol and hydrocarbon  
1087 composition of Antarctic sea ice diatom communities during the spring bloom in McMurdo  
1088 Sound. *Antarctic Science* 5, 271–278.

1089 Nichols, P.D., Skerratt, J.H., Davidson, A., Burton, H., McMeekin, T.A., 1991. Lipids of  
1090 cultured *Phaeocystis pouchetii*: Signatures for food-web, biogeochemical and environmental  
1091 studies in Antarctica and the Southern ocean. *Phytochemistry* 30, 3209–3214.  
1092 [https://doi.org/10.1016/0031-9422\(91\)83177-M](https://doi.org/10.1016/0031-9422(91)83177-M)

1093 Nielsen, B.L.H., Gøtterup, L., Jørgensen, T.S., Hansen, B.W., Hansen, L.H., Mortensen, J.,  
1094 Jepsen, P.M., 2019. n-3 PUFA biosynthesis by the copepod *Apocyclops royi* documented using  
1095 fatty acid profile analysis and gene expression analysis. *Biology Open* 8.  
1096 <https://doi.org/10.1242/bio.038331>

1097 Okuyama, H., Morita, N., Kogame, K., 1992. Occurrence of octadecapentaenoic acid in lipids  
1098 of a cold stenothermic alga, prymnesiophyte strain B. *Journal of Phycology* 28, 465–472.

1099 Orsi, A.H., Whitworth, T., Nowlin, W.D., 1995. On the meridional extent and fronts of the  
1100 Antarctic Circumpolar Current. *Deep Sea Research Part I: Oceanographic Research Papers* 42,  
1101 641–673. [https://doi.org/10.1016/0967-0637\(95\)00021-W](https://doi.org/10.1016/0967-0637(95)00021-W)

1102 Pakhomov, E.A., McQuaid, C.D., 1996. Distribution of surface zooplankton and seabirds  
1103 across the Southern Ocean. *Polar Biology* 16, 271–286. <https://doi.org/10.1007/s0030000050054>

1104 Park, Y.-H., Fuda, J.-L., Durand, I., Naveira Garabato, A.C., 2008. Internal tides and vertical  
1105 mixing over the Kerguelen Plateau. *Deep Sea Research Part II: Topical Studies in Oceanography*  
1106 55, 582–593. <https://doi.org/10.1016/j.dsr2.2007.12.027>

1107 Parrish, C.C., 2013. Lipids in marine ecosystems. *ISRN Oceanography* 2013, 604045.  
1108 <https://doi.org/10.5402/2013/604045>

1109 Pauthenet, E., Roquet, F., Madec, G., Guinet, C., Hindell, M., McMahon, C.R., Harcourt, R.,  
1110 Nerini, D., 2018. Seasonal Meandering of the Polar Front Upstream of the Kerguelen Plateau.  
1111 *Geophysical Research Letters* 45, 9774–9781. <https://doi.org/10.1029/2018gl079614>

1112 Pellichero, V., Boutin, J., Claustre, H., Merlivat, L., Sallée, J.-B., Blain, S., 2020. Relaxation of  
1113 Wind Stress Drives the Abrupt Onset of Biological Carbon Uptake in the Kerguelen Bloom: A  
1114 Multisensor Approach. *Geophysical Research Letters* 47, e2019GL085992.  
1115 <https://doi.org/10.1029/2019gl085992>

1116 Perissinotto, R., Duncombe Rae, C.M., 1990. Occurrence of anticyclonic eddies on the Prince  
1117 Edward Plateau (Southern Ocean): effects on phytoplankton biomass and production. *Deep Sea*  
1118 *Research Part A. Oceanographic Research Papers* 37, 777–793. [https://doi.org/10.1016/0198-](https://doi.org/10.1016/0198-0149(90)90006-H)  
1119 [0149\(90\)90006-H](https://doi.org/10.1016/0198-0149(90)90006-H)

1120 Pollard, R., Sanders, R., Lucas, M., Statham, P., 2007. The Crozet Natural Iron Bloom and  
1121 Export Experiment (CROZEX). *Deep Sea Research Part II: Topical Studies in Oceanography* 54,  
1122 1905–1914. <https://doi.org/10.1016/j.dsr2.2007.07.023>

1123 Pond, D.W., Atkinson, A., Shreeve, R.S., Tarling, G., Ward, P., 2005. Diatom fatty acid  
1124 biomarkers indicate recent growth rates in Antarctic krill. *Limnology and Oceanography* 50, 732–  
1125 736. <https://doi.org/10.4319/lo.2005.50.2.0732>

1126 Quéroué, F., Sarthou, G., Planquette, H.F., Bucciarelli, E., Chever, F., van der Merwe, P.,  
1127 Lannuzel, D., Townsend, A.T., Cheize, M., Blain, S., d’Ovidio, F., Bowie, A.R., 2015. High  
1128 variability of dissolved iron concentrations in the vicinity of Kerguelen Island (Southern Ocean).  
1129 *Biogeosciences Discuss.* 12, 231–270. <https://doi.org/10.5194/bgd-12-231-2015>

1130 Rembauville, M., Blain, S., Armand, L., Quéguiner, B., Salter, I., 2015a. Export fluxes in a  
1131 naturally iron-fertilized area of the Southern Ocean – Part 2: Importance of diatom resting spores  
1132 and faecal pellets for export. *Biogeosciences* 12, 3171–3195. [https://doi.org/10.5194/bg-12-3171-](https://doi.org/10.5194/bg-12-3171-2015)  
1133 2015

1134 Rembauville, M., Blain, S., Manno, C., Tarling, G., Thompson, A., Wolff, G., Salter, I., 2018.  
1135 The role of diatom resting spores in pelagic–benthic coupling in the Southern Ocean.  
1136 *Biogeosciences* 15, 3071–3084. <https://doi.org/10.5194/bg-15-3071-2018>

1137 Rembauville, M., Salter, I., Dehairs, F., Miquel, J.-C., Blain, S., 2017. Annual particulate matter  
1138 and diatom export in a high nutrient, low chlorophyll area of the Southern Ocean. *Polar Biology*.  
1139 <https://doi.org/10.1007/s00300-017-2167-3>

1140 Rembauville, M., Salter, I., Leblond, N., Gueneugues, A., Blain, S., 2015b. Export fluxes in a  
1141 naturally iron-fertilized area of the Southern Ocean – Part 1: Seasonal dynamics of particulate  
1142 organic carbon export from a moored sediment trap. *Biogeosciences* 12, 3153–3170.  
1143 <https://doi.org/10.5194/bg-12-3153-2015>

1144 Remize, M., Planchon, F., Loh, A.N., Le Grand, F., Bideau, A., Le Goic, N., Fleury, E., Miner,  
1145 P., Corvaisier, R., Volety, A., Soudant, P., 2020. Study of Synthesis Pathways of the Essential  
1146 Polyunsaturated Fatty Acid 20:5n-3 in the Diatom *Chaetoceros Muelleri* Using <sup>13</sup>C-Isotope  
1147 Labeling. *Biomolecules* 10. <https://doi.org/10.3390/biom10050797>

1148 Salter, I., Kemp, A.E.S., Moore, C.M., Lampitt, R.S., Wolff, G.A., Holtvoeth, J., 2012. Diatom  
1149 resting spore ecology drives enhanced carbon export from a naturally iron-fertilized bloom in the  
1150 Southern Ocean. *Global Biogeochemical Cycles* 26, GB1014. <https://doi.org/10.1029/2010gb003977>

1151 Salter, I., Lampitt, R.S., Sanders, R., Poulton, A., Kemp, A.E.S., Boorman, B., Saw, K., Pearce,  
1152 R., 2007. Estimating carbon, silica and diatom export from a naturally fertilised phytoplankton  
1153 bloom in the Southern Ocean using PELAGRA: A novel drifting sediment trap. *Deep Sea Research*  
1154 *Part II: Topical Studies in Oceanography* 54, 2233–2259. <https://doi.org/10.1016/j.dsr2.2007.06.008>

1155 Sargent, J.R., Eilertsen, H.C., Falk-Petersen, S., Taasen, J.P., 1985. Carbon assimilation and lipid  
1156 production in phytoplankton in northern Norwegian fjords. *Marine Biology* 85, 109–116.  
1157 <https://doi.org/10.1007/BF00397428>

1158 Sassenhagen, I., Irion, S., Jardillier, L., Moreira, D., Christaki, U., 2020. Protist Interactions and  
1159 Community Structure During Early Autumn in the Kerguelen Region (Southern Ocean). *Protist*  
1160 171, 125709. <https://doi.org/10.1016/j.protis.2019.125709>

1161 Schallenberg, C., Bestley, S., Klocker, A., Trull, T.W., Davies, D.M., Gault-Ringold, M., Eriksen,  
1162 R., Roden, N.P., Sander, S.G., Sumner, M., Townsend, A.T., van der Merwe, P., Westwood, K.,  
1163 Wuttig, K., Bowie, A., 2018. Sustained Upwelling of Subsurface Iron Supplies Seasonally  
1164 Persistent Phytoplankton Blooms Around the Southern Kerguelen Plateau, Southern Ocean.  
1165 *Journal of Geophysical Research: Oceans* 123, 5986–6003. <https://doi.org/10.1029/2018jc013932>

1166 Schlosser, C., Schmidt, K., Aquilina, A., Homoky, W.B., Castrillejo, M., Mills, R.A., Patey,  
1167 M.D., Fielding, S., Atkinson, A., Achterberg, E.P., 2018. Mechanisms of dissolved and labile  
1168 particulate iron supply to shelf waters and phytoplankton blooms off South Georgia, Southern  
1169 Ocean. *Biogeosciences* 15, 4973–4993. <https://doi.org/10.5194/bg-15-4973-2018>

1170 Seeyave, S., Lucas, M.I., Moore, C.M., Poulton, A.J., 2007. Phytoplankton productivity and  
1171 community structure in the vicinity of the Crozet Plateau during austral summer 2004/2005. *Deep*  
1172 *Sea Research Part II: Topical Studies in Oceanography* 54, 2020–2044.  
1173 <https://doi.org/10.1016/j.dsr2.2007.06.010>

1174 Sheridan, C.C., Lee, C., Wakeham, S.G., Bishop, J.K.B., 2002. Suspended particle organic  
1175 composition and cycling in surface and midwaters of the equatorial Pacific Ocean. *Deep Sea*  
1176 *Research Part I: Oceanographic Research Papers* 49, 1983–2008. [https://doi.org/10.1016/S0967-](https://doi.org/10.1016/S0967-0637(02)00118-8)  
1177 [0637\(02\)00118-8](https://doi.org/10.1016/S0967-0637(02)00118-8)

1178 Skerratt, J.H., Nichols, P.D., McMeekin, T.A., Burton, H., 1995. Seasonal and inter-annual  
1179 changes in planktonic biomass and community structure in eastern Antarctica using signature  
1180 lipids. *Marine Chemistry* 51, 93–113. [https://doi.org/10.1016/0304-4203\(95\)00047-U](https://doi.org/10.1016/0304-4203(95)00047-U)

1181 Tang, K.W., Jakobsen, H.H., Visser, A.W., 2001. *Phaeocystis globosa* (Prymnesiophyceae) and  
1182 the planktonic food web: Feeding, growth, and trophic interactions among grazers. *Limnology*  
1183 and *Oceanography* 46, 1860–1870. <https://doi.org/10.4319/lo.2001.46.8.1860>

1184 Thomson, P.G., Wright, S.W., Bolch, C.J.S., Nichols, P.D., Skerratt, J.H., McMinn, A., 2004.  
1185 Antarctic distribution, pigment and lipid composition, and molecular identification of the brine  
1186 dinoflagellate *Polarella glacialis* (Dinophyceae). *Journal of Phycology* 40, 867–873.  
1187 <https://doi.org/10.1111/j.1529-8817.2004.03169.x>

1188 Trull, T.W., Davies, D.M., Dehairs, F., Cavagna, A.J., Lasbleiz, M., Laurenceau-Cornec, E.C.,  
1189 d’Ovidio, F., Planchon, F., Leblanc, K., Quéguiner, B., Blain, S., 2015. Chemometric perspectives  
1190 on plankton community responses to natural iron fertilisation over and downstream of the  
1191 Kerguelen Plateau in the Southern Ocean. *Biogeosciences* 12, 1029–1056.  
1192 <https://doi.org/10.5194/bg-12-1029-2015>

1193 Vaezi, R., Napier, J.A., Sayanova, O., 2013. Identification and functional characterization of  
1194 genes encoding omega-3 polyunsaturated fatty acid biosynthetic activities from unicellular  
1195 microalgae. *Marine drugs* 11, 5116–5129.

1196 van der Merwe, P., Bowie, A.R., Quéroúé, F., Armand, L., Blain, S., Chever, F., Davies, D.,  
1197 Dehairs, F., Planchon, F., Sarthou, G., Townsend, A.T., Trull, T.W., 2015. Sourcing the iron in the  
1198 naturally fertilised bloom around the Kerguelen Plateau: particulate trace metal dynamics.  
1199 *Biogeosciences* 12, 739–755. <https://doi.org/10.5194/bg-12-739-2015>

1200 Venables, H., Moore, C.M., 2010. Phytoplankton and light limitation in the Southern Ocean:  
1201 Learning from high-nutrient, high-chlorophyll areas. *Journal of Geophysical Research: Oceans*  
1202 115. <https://doi.org/10.1029/2009jc005361>

1203 Virtue, P., Nichols, P.D., Nicol, S., McMinn, A., Sikes, E.L., 1993. The lipid composition of  
1204 *Euphausia superba* Dana in relation to the nutritional value of *Phaeocystis pouchetii* (Hariot)  
1205 Lagerheim. *Antarctic Science* 5, 169–177. <https://doi.org/10.1017/S0954102093000239>

1206 Volkman, J.K., Barrett, S.M., Blackburn, S.I., Mansour, M.P., Sikes, E.L., Gelin, F., 1998.  
1207 Microalgal biomarkers: A review of recent research developments. *Organic Chemistry* 29, 1163–  
1208 1179. [https://doi.org/10.1016/S0146-6380\(98\)00062-X](https://doi.org/10.1016/S0146-6380(98)00062-X)

1209 Wakeham, S.G., 1995. Lipid biomarkers for heterotrophic alteration of suspended particulate  
1210 organic matter in oxygenated and anoxic water columns of the ocean. *Deep Sea Research Part I:*  
1211 *Oceanographic Research Papers* 42, 1749–1771. [https://doi.org/10.1016/0967-0637\(95\)00074-G](https://doi.org/10.1016/0967-0637(95)00074-G)

1212 Wang, S.W., Budge, S.M., Iken, K., Gradinger, R.R., Springer, A.M., Wooller, M.J., 2015.  
1213 Importance of sympagic production to Bering Sea zooplankton as revealed from fatty acid-carbon  
1214 stable isotope analyses. *Mar Ecol Prog Ser* 518, 31–50.

1215 Wilson, S.E., Steinberg, D.K., Chu, F.L.E., Bishop, J.K.B., 2010. Feeding ecology of mesopelagic  
1216 zooplankton of the subtropical and subarctic North Pacific Ocean determined with fatty acid  
1217 biomarkers. *Deep Sea Research Part I: Oceanographic Research Papers* 57, 1278–1294.  
1218 <https://doi.org/10.1016/j.dsr.2010.07.005>

1219