



HAL
open science

Diversity and molecular epidemiology of Ostreid herpesvirus 1 in farmed *Crassostrea gigas* in Australia: Geographic clusters and implications for "microvariants" in global mortality events

Suzanne Trancart, Alison Tweedie, Olivia Liu, Ika Paul-Pont, Paul Hick,
Maryline Houssin, Richard J Whittington

► To cite this version:

Suzanne Trancart, Alison Tweedie, Olivia Liu, Ika Paul-Pont, Paul Hick, et al.. Diversity and molecular epidemiology of Ostreid herpesvirus 1 in farmed *Crassostrea gigas* in Australia: Geographic clusters and implications for "microvariants" in global mortality events. *Virus Research*, 2022, 323, 10.1016/j.virusres.2022.198994 . hal-04032451

HAL Id: hal-04032451

<https://hal.univ-brest.fr/hal-04032451>

Submitted on 29 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Diversity and molecular epidemiology of *Ostreid herpesvirus 1* in farmed *Crassostrea***
2 ***gigas* in Australia: geographic clusters and implications for “microvariants” in global**
3 **mortality events**

4
5
6
7 Suzanne Trancart ^a, Alison Tweedie ^{bc}, Olivia Liu^{bd}, Ika Paul-Pont ^{be}, Paul Hick^{bc}, Maryline
8 Houssin ^{af} and Richard J Whittington ^{b*}

9
10 ^a LABÉO Research department, 1 Route de Rosel, 14053 Caen, Cedex 4, France

11
12 ^b The University of Sydney, Sydney School of Veterinary Science, Faculty of Science,
13 Camden, NSW 2570, Australia

14
15 ^c Present address : Elizabeth Macarthur Agricultural Institute, Department of Primary
16 Industries, Menangle, NSW 2568, Australia

17
18 ^d Present address : Department of Agriculture, Water and the Environment, Canberra, ACT
19 2601, Australia

20
21 ^e Present address : LEMAR, Rue Dumont d'Urville, 29280 Plouzané, FRANCE

22
23 ^f UMR BOREA Université de Caen Normandie, MNHN, CNRS 8067, SU, IRD 207, UCN,
24 UA, Esplanade de la Paix, 14032 Caen Cedex 4, France

25
26 *Corresponding author: RJ Whittington, The University of Sydney, Sydney School of
27 Veterinary Science, Faculty of Science, The University of Sydney, 425 Werombi Rd, Camden,
28 NSW 2570, Australia

29 E-mail: richard.whittington@sydney.edu.au;

30 Tel: +61 2 9351 1619 | +61 2 9351 1618 (fax)| +61 (0) 438 983 195 (mobile)

31

32 **Highlights**

- 33 • A relatively low diversity of OsHV-1 genotypes was found in Australia
- 34 • Distinct *OsHV-1* genotypes were geographically clustered in estuaries
- 35 • Australian *OsHV-1* genotypes grouped in a globally distinct cluster
- 36 • Australian POMS outbreaks were not due to OsHV-1 microvariants

37

38 **Abstract**

39 Since 2010, mass mortality events known as Pacific oyster mortality syndrome (POMS) have
40 occurred in *Crassostrea gigas* in Australia associated with *Ostreid herpesvirus 1*. The virus was
41 thought to be an OsHV-1 μ Var or “microvariant”, i.e. one of the dominant variants associated
42 with POMS in Europe, but there are few data to characterize the genotype in Australia.
43 Consequently, the genetic identity and diversity of the virus was determined to understand the
44 epidemiology of the disease in Australia. Samples were analysed from diseased *C. gigas* over
45 five summer seasons between 2011 and 2016 in POMS-affected estuaries: Georges River in
46 New South Wales (NSW), Hawkesbury River (NSW) and Pitt Water in Tasmania. Sequencing
47 was attempted for six genomic regions. Numerous variants were identified among these regions
48 ($n = 100$ isolates) while twelve variants were identified from concatenated nucleotide sequences
49 ($n = 61$ isolates). Nucleotide diversity of the seven genotypes of C region among Australian
50 isolates ($P_i 0.99 \times 10^{-3}$) was the lowest globally. All Australian isolates grouped in a cluster
51 distinct from other OsHV-1 isolates worldwide. This is the first report that Australian outbreaks
52 of POMS were associated with OsHV-1 distinct from OsHV-1 reference genotype, μ Var and
53 other microvariants from other countries. The findings illustrate that microvariants are not the
54 only variants of OsHV-1 associated with mass mortality events in *C. gigas*. In addition, there
55 was mutually exclusive spatial clustering of viral genomic and amino acid sequence variants
56 within estuaries, and a possible association between genotype/amino acid sequence and the

57 prevalence and severity of POMS, as this differed between these estuaries. The sequencing
58 findings supported prior epidemiological evidence for environmental reservoirs of OsHV-1 for
59 POMS outbreaks in Australia.

60

61 **Keywords: Ostreid herpesvirus, variant, diversity, polymorphism, phylogeny, *Crassostrea***
62 ***gigas*, Australia**

63

64 **1. Introduction**

65 Since late last century, the Pacific oyster, *Crassostrea gigas*, has been subject to
66 recurrent disease events (Nicolas et al., 1992; Renault et al., 2000; Renault et al., 1994; Samain
67 et al., 2007). However, in 2008 a marked increase in mass mortality outbreaks occurred in
68 France (Cochennec-Laureau et al., 2010; Martenot et al., 2011; Segarra et al., 2010) and later
69 in other areas in the world (Abbadi et al., 2018; Batista et al., 2015; Gittenberger et al., 2016;
70 Hwang et al., 2013; Keeling et al., 2014; Mortensen et al., 2016; Peeler et al., 2012; Roque et
71 al., 2012; Segarra et al., 2010). The disease was called Pacific oyster mortality syndrome
72 (POMS) when it first appeared in Australia in 2010 in the Georges and Parramatta rivers, which
73 are both major estuaries in Sydney (Jenkins et al., 2013; Paul-Pont et al., 2013a). POMS causes
74 considerable economic losses in Australia, New Zealand and Europe (Fuhrmann et al., 2019).
75 It is a multifactorial disease influenced by environmental conditions such as water temperature
76 and host characteristics such as age and size (Hick et al., 2018; Petton et al., 2013) and although
77 it has a proposed polymicrobial etiology involving bacteria (de Lorgeril et al., 2018; Friedman
78 et al., 2005; Petton et al., 2021), outbreaks are induced by infection with *Ostreid herpesvirus 1*
79 microvariants (Batista et al., 2015; Gittenberger et al., 2016; Hwang et al., 2013; Jenkins et al.,
80 2013; Keeling et al., 2014; Mortensen et al., 2016; Peeler et al., 2012; Renault et al., 2012;
81 Roque et al., 2012). The disease typically occurs in summer and mortality can be reproduced

82 experimentally by controlled exposure of oysters to the virus alone in a dose-responsive and
83 temperature-dependent manner (de Kantzow et al., 2016; Paul-Pont et al., 2015).

84 *Ostreid herpesvirus 1*, OsHV-1, which is the only species included in the genus
85 *Ostreavirus*, is a double-stranded DNA virus (Davison et al., 2009). The presence of the gene
86 coding for the ATPase subunit of a terminase involved in the packaging of viral DNA inside
87 the capsid, in addition to intranuclear replication, places OsHV-1 in the *Malacoherpesviridae*
88 under the order *Herpesvirales* (Davison et al., 2009; Davison et al., 2005). Members of this
89 taxon are widespread, both in terms of the range of species infected and also geographically.
90 They have been detected in many hosts besides *C. gigas* including *Crassostrea virginica* (Farley
91 et al., 1972), *Ostrea angasi* (Hine and Thorne, 1997), *Ostrea edulis* (Comps and Cochenec,
92 1993), *Tiostrea chilensis* (Hine et al., 1998), *Pecten maximus* (Arzul et al., 2001a), *Ruditapes*
93 *philippinarum* (Renault et al., 2001), *Ruditapes decussatus* (Renault and Arzul, 2001),
94 *Crassostrea angulata* (Batista et al., 2015), *Scapharca broughtonii* (Bai et al., 2016; Xia et al.,
95 2015), *Carcinus maenas* (Bookelaar et al., 2018), *Chlamys farreri* (Bai et al., 2019), *Octopus*
96 *vulgaris* (Prado-Alvarez et al., 2021) and in numerous sites around the world including Europe,
97 Asia, Oceania and the Americas (Abadi et al., 2018 ; Batista et al., 2015; Burge et al., 2021;
98 Gittenberger et al., 2016; Hwang et al., 2013; Jenkins et al., 2013; Keeling et al., 2014;
99 Mortensen et al., 2016; Peeler et al., 2012; Roque et al., 2012; Segarra et al., 2010; Shimahara
100 et al., 2012).

101 A specific microvariant, OsHV-1 μ Var, was described in association with the large
102 increase in mortality in *C. gigas* on the French coast in 2008 (Segarra et al., 2010), but the
103 detection of this virus in a specimen collected before 2008 (Martenot et al., 2012) raises the
104 question of its actual or exclusive involvement in the increasing global phenomenon of oyster
105 mortality. OsHV-1 μ Var was strictly characterised by mutations upstream of ORF 4 consisting
106 mainly of 12 deletions in the microsatellite zone and a few mutations upstream and downstream

107 which impact the amino acid sequence. In addition, at ORF 42-43 OsHV-1 μ Var had a
108 substitution of a thymine instead of a cytosine downstream of the sequence without showing any
109 impact on the amino acid sequence (Segarra et al., 2010). Following the publication of the
110 OsHV-1 μ Var sequence from region C, partly located at ORF 4 (Segarra et al., 2010), several
111 isolates were analysed and this revealed a range of « microvariants » close to the OsHV-1 μ Var
112 sequence (Burge et al., 2021; Burioli et al., 2018; Martenot et al., 2012; Morga et al., 2021).
113 Data for numerous OsHV-1 genotypes characterised by sequencing of various coding and non-
114 coding regions or whole genomes are available: in Europe, the reference genome isolated from
115 *C. gigas* larvae (Davison et al., 2005), the variant OsHV-1 var sampled from *P. maximus* (Arzul
116 et al., 2001a) and the microvariant OsHV-1 μ Var from *C. gigas* spat and juveniles (Abbadì et
117 al., 2018; Burioli et al., 2017; Segarra et al., 2010); in China, Acute Viral Necrosis Virus (Ren
118 et al., 2013), OsHV-1-SB isolated from *S. broughtonii* (Bai et al., 2016; Xia et al., 2015) and
119 the variant isolated from *C. farreri* (Bai et al., 2019); in Sweden OsHV-1 SW6 found in *O.*
120 *edulis* (Morga et al., 2021); and in many parts of the world, so called « microvariants » or
121 μ Vars, which are similar to OsHV-1 μ Var (Burge et al., 2021; Morga et al., 2021). Double-
122 stranded DNA viruses are known to be genetically stable; DNA viruses in general have
123 mutation rates of 10^{-8} to 10^{-6} substitutions per nucleotide site per cell infection, a lower rate than
124 RNA viruses, due the viral DNA polymerase that exhibits proofreading activity (Peck and
125 Lauring, 2018). However, Morga et al. (2021) recently reported a faster evolutionary rate than
126 usual for OsHV-1. The emergence of variations may be due to selection pressure and the ability
127 of the virus to adapt to its environment and host as reported in other members of the
128 *Herpesvirales* (Renzette et al., 2013). The available data suggest that pathogenic subtypes of
129 OsHV-1 known as microvariants possess genetic diversity beyond that defined for a single
130 genotype by Segarra et al. (2010), but the extent of diversity and how this relates to their global
131 emergence is unclear.

132 The first detection of OsHV-1 associated with mortality of *C. gigas* in Australia occurred
133 in 2010 but despite recurrent outbreaks there has been little characterisation of the genetic
134 diversity of the viruses affecting *C. gigas* in Australia (Jenkins et al., 2013). Available data
135 suggest that the genotype responsible for the index case in 2010 was close to OsHV-1 μ Var but
136 had some differences (Jenkins et al., 2013). Given that OsHV-1 genetic diversity could affect
137 the gravity of the disease (Delmotte et al., 2022; Martenot et al., 2011; Morga et al., 2021), it is
138 important to study the genetic composition of the Australian isolates to better understand
139 epidemiological patterns in Australia. Detailed, long term, epidemiological studies in Australia
140 revealed that POMS outbreaks recurred at the same sites over successive years (Whittington et
141 al., 2019). The source of OsHV-1 for these disease events between 2011 and 2016 was
142 environmental rather than spread through oyster farming activities (Whittington et al., 2018).

143 The aim of this report is to (i) study the identity and genetic diversity of OsHV-1 in
144 archival samples collected during mortality events in Australia using six regions sequenced by
145 Sanger technology, (ii) compare the Australian viruses with international isolates previously
146 published, and iii) make inferences based on diversity and phylogeny about the source and
147 transmission of the virus to inform management decisions for disease prevention and control.

148

149 **2. Materials and methods**

150

151 **2.1. Oyster sample selection**

152

153 Oyster samples were selected over the five summer seasons between 2011 and 2016
154 from the entire geographical distribution of POMS disease outbreaks in commercial aquaculture
155 in Australia. Comprehensive data for each sample including location, age, size, mortality rate
156 and viral load are provided in Supplementary data 1, linked by references to scientific

157 publications on these populations. In summary, all samples were collected on oyster farms by
158 researchers from the University of Sydney and all oysters in the final dataset were initially
159 recruited as spat from an OsHV-1 free hatchery in Tasmania (Table 1). All samples were tested
160 using OsHV-1 real-time PCR (Martenot et al., 2010) and positive samples containing the
161 highest concentration of OsHV-1 DNA from among the samples collected that day were
162 selected for sequencing. A total of 118 samples were selected and sequence was subsequently
163 obtained from at least one region of the genome from 107 samples to which a unique “isolate
164 ID” was then assigned. For seven of these (all wild *C. gigas* or *Saccostrea commercialis* from
165 the Georges River in the 2013-2014 season) sequence was obtained only for Region 3 which
166 had no variation (see below); these isolates were excluded from further analysis, leaving 100
167 isolates available for analysis, all being samples from farmed *C. gigas* (Supplementary data 1).
168 Of the 100 samples of *C. gigas*, 38 were collected during periods of OsHV-1 infection in the
169 Hawkesbury River estuary north of Sydney NSW (from Coba Bay n=2; Kimmerikong Bay
170 n=5; Mooney Mooney n=15; Mullet Creek n=10; Porto Bay n=6), 60 were collected in the
171 Georges River estuary south of Sydney NSW (from Limekiln Bar n=1; Neverfail Bay n=2; Site
172 A n=13; Site B n=10; Site C n=25; Sylvania Waters n=1; The Shed n=8), while two were
173 collected in Tasmania (from Pitt Water n=2) (Figure 1). Exact locations are illustrated in maps
174 in Whittington et al. (2019) and in de Kantzow et al. (2017). Sampled oysters had an age range
175 of 5.5 months to 17 months and a size range of 5mm to 110mm. They were all the subject of
176 published studies in which details of the oyster populations are reported (de Kantzow et al.,
177 2017; Hick et al., 2018; Paul-Pont et al., 2013a; Paul-Pont et al., 2013b; Paul-Pont et al., 2014;
178 Whittington et al., 2015a; Whittington et al., 2015b; Whittington et al., 2019) (Supplementary
179 data 1).

180

181 2.2. Description of the selected regions

182

183 Six regions of the OsHV-1 genome were targeted for Sanger sequencing. Firstly, four
184 regions were selected based on their discriminatory value for distinguishing microvariant
185 genotypes from among other OsHV-1 variants. Region 1 and Region 2 targeted ORF 4 and
186 ORF 43, respectively. They were selected in order to distinguish OsHV-1 μ Var (Segarra et al.,
187 2010), OsHV-1 μ Var Δ 9, OsHV-1 μ Var Δ 15 described in Europe (Martenot et al., 2012;
188 Martenot et al., 2011) and other microvariants (Bai et al., 2016). Region 3 covered ORFs 35,
189 36, 37 and 38 and bounded a large deletion in OsHV-1 μ Var that induced the suppression of
190 ORF 36 and 37 and disrupted ORF 38 (Burioli et al., 2017; Martenot et al., 2015; Martenot et
191 al., 2013; Morga et al., 2021; Renault et al., 2012). Region 4 was selected for the high level of
192 polymorphism suitable for distinguishing genotypes of OsHV-1 (Batista et al., 2015). It is
193 located between ORF 49 and ORF 50 and is a non-coding site. Region 5 located on ORF 11,
194 showed a large deletion of 1386bp in microvariants (Morga et al., 2021 ; Burioli et al., 2017 ;
195 Martenot et al., 2013). Finally, Region 6, which could be involved in virulence mechanisms is
196 a polymorphic region of interest; it targets ORF 88 coding for a transmembrane glycoprotein
197 (Martenot et al., 2013). PCR and sequencing reaction primers used for each region are provided
198 in Table 2 .

199

200 2.3. Sample processing

201 2.3.1. Extraction

202 Each sample was derived from whole animal or dissected mantle and gill tissues of one
203 or more oysters, depending on their size (Supplementary data 1). The DNA extraction was
204 performed according to previous studies (Evans et al., 2014; Whittington et al., 2019). Briefly,
205 oysters were collected in the field and transported directly to the laboratory where they were
206 frozen at -80°C until processed. A 400 ± 100 mg sample was homogenised by bead beating

207 using sterile stainless steel beads (Aussie Sapphires) and 1 mL distilled water and placed into a
208 TissueLyser II machine (Qiagen) for 2 min at frequency 30, repeated once. All samples were
209 clarified by centrifugation at 1340 g for 2 min in a microcentrifuge and supernatants were
210 removed and stored in sterile tubes at -80°C . Nucleic acids were purified using a 5X
211 MagMAXTM-96Viral RNA Isolation Kit (Ambion, Life TechnologiesTM,
212 Mulgrave,Australia) and then MagMAXTM Express 96 magnetic particle processor (Applied
213 BiosystemsTM, Life TechnologiesTM, Mulgrave, Australia) according to manufacturers'
214 instructions for a final volume of 50 μL using the AM1836 deep-well standard programme
215 (Ambion, Life TechnologiesTM, Mulgrave, Australia). Purified nucleic acids were stored at
216 -20°C .

217

218 2.3.2. PCR

219 Various PCR were performed in order to target the 6 regions described above in part 2.2.
220 The primers are reported in Table 2. For Region 1 and Region 2, each reaction contained 5 μl
221 DNA extract, 5 μl 10X PCR buffer (66.6mM Tris-HCl,16.6mM $(\text{NH}_4)_2\text{SO}_4$, 2.5mM MgCl_2 ,
222 1.65mg/ml bovine serum albumin,10mM beta-mercaptoethanol), 10 μl dNTP mix (1mM),
223 250nM each forward and reverse primer, 5U DNA polymerase (taq:pfu mix) and nuclease-free
224 water to a final volume of 50 μl . Thermocycling was performed (Corbett Research CGI960)
225 according to the following conditions: 1 cycle at 94°C for 2min; 35 cycles consisted in 94°C
226 for 1min, 50°C for 1 min and 72°C for 1min, and a final extension at 72°C for 5 min. For the
227 remaining regions, each reaction contained 10 μl 5X HiFi reaction buffer (Bioline) (containing
228 10mM Mg^{2+}), 0.5 μl dNTP mix (100mM), 1 μl each forward and reverse primer (400nM), 1 μl
229 (2U) Velocity DNA polymerase (Bioline), 1.5 μl DMSO (BIO-21098 -Bioline Aust Pty. Ltd),
230 5 μl template DNA and nuclease-free water to a final volume of 50 μl ; thermocycling was

231 performed according to the following conditions: 1 cycle at 98°C for 2min; 35 cycles consisted
232 in 98°C for 30s, 62°C for 30s and 72°C for 30s, and a final extension at 72°C for 10 min.

233 The amplicons were visualised using 5µl of each amplification product on a 2% agarose
234 gel, assessed against a molecular weight marker and visualised using RedSafe (iNtRON
235 Biotechnology) on a GelDoc transilluminator (Biorad).

236

237 2.3.3. Sequencing

238 Amplified PCR product was purified by incubation with ExoSAP-IT/Cleansweep
239 (ThermoFisher Scientific) then submitted for Sanger sequencing. For several samples where a
240 single amplicon was not obtained, bands of the expected size were excised from the agarose gel
241 and DNA was purified using the QIAquick Gel Extraction Kit (Qiagen). The amplicons were
242 sent to the Australian Genome Research Facility or to the Monash Health Translation Precinct
243 Medical Genomics Facility Australia for sequencing. Forward and reverse sequencing were
244 performed in reactions with relevant PCR primers and internal primers (Table 2).
245 Chromatograms were reviewed, analyzed and primer sequence was removed, using FinchTV
246 (Geospiza).

247

248 2.4. DNA data processing

249

250 For each region targeted, nucleotide sequences were aligned with the reference genome
251 NC_005881.2 and the two microvariant genomes KY242788 and KY271630, using MEGA10
252 (Kumar et al., 2018) with the MUSCLE algorithm. The sequences were trimmed and the
253 chromatograms were again specifically checked for all mutations on forward and reverse
254 strands. Base positions described in results are mapped to the reference genome NC_005881.2.

255 Phylogenetic trees were constructed using MEGA 10 with the Neighbour Joining
256 method and 1000 replications of bootstraps and the method based on Maximum Composite
257 Likelihood was applied. Two trees were obtained: the first was based on the concatenated
258 sequence of Regions 1, 2, 3, 4 and 6 (Region 5 was not included due to too few sequences - see
259 results); the second tree was based on Region 1. The trees were unrooted because the ancestry
260 of the reference genotype is uncertain. In addition, the use of another *malacoherpesviridae*,
261 Haliotid herpesvirus, was impracticable because the ORFs had no sequence in common making
262 alignment impossible. The unrooted tree allows the analysis of diversity but does not orient the
263 genotypes temporally.

264 Nucleotide diversity (Pi) for Region 1 (C region) including the microsatellite region was
265 calculated using DnaSP v6: DNA Sequence Polymorphism Analysis of Large Datasets (Rozas
266 et al., 2017) and compared with the results of Mineur et al. (2015), after adding six sequences
267 (JN800089, JN800075, JN800082, JN800083, JN800088 and JN800072) to be consistent with
268 these authors. Briefly, the calculation was performed on the alignment; the indel polymorphism
269 module with the “Multiallelic”gap option was used to consider all InDel events. Nucleotide
270 diversity was listed according to geographic region. Pi values reported by Mineur et al. (2015)
271 were corrected for frequency equal to one because multiple occurrences of an identical
272 genotype may not be reported. To make the data comparable, Pi in the current study was not
273 corrected because only one sequence per genotype was included in the calculation.

274

275

276 2.5. Amino acid sequence prediction

277

278 In order to estimate whether some putative mutations could induce a modification in the
279 phenotype of the variant, a prediction of amino acid sequence was performed on coding sites in

280 Regions 1, 2, 3, 5 and 6. The translations of the ORFs were performed with ORF Finder (<https://www.ncbi.nlm.nih.gov/orffinder/>) and the Bioedit sequence alignment editor. The translations
281 were checked using Blastp (<https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE=Proteins>). These
282 predictions were aligned with the prediction of the reference genome NC_005881.2 and the two
283 microvariant genomes KY242788 and KY271630 using Mega 10.
284

285

286

287 **3. Results**

288 3.1. DNA sequencing analysis

289

290 3.1.1. Description of the variants

291 All nucleotide sequences are available on the NCBI database and accession numbers are
292 listed in Table 3. Results and metadata for each of the 100 samples is provided in Supplementary
293 data 1. Sequence alignments are illustrated in Supplementary data 2.
294

294

295 For Region 1, located at ORF 4, nucleotide sequence was obtained from 87 of 100
296 samples. None of the sequences were identical to the OsHV-1 reference sequence but they were
297 partially similar to the microvariants, with the presence of most of the mutations except the two
298 substitutions of adenine instead of guanine in the downstream region (base 178,696 and
299 178,702). However, they all presented a novel substitution of an adenine instead of a guanine
300 in a different position (base 178,705), also in the downstream region. Several mutations were
301 observed allowing the identification of five different nucleotide sequences: G1, G2, G4, G6 and
302 G7 (Figure 1, Supplementary data 2). Eight of the sequences that grouped in G4 (Supplementary
303 data 2) had 15 deletions (base 178,558 - 178,572) in the microsatellite zone instead of the 12

304 deletions seen in microvariants (Burioli et al., 2017; Segarra et al., 2010), as was observed in
305 OsHV-1 μ Var Δ 15 reported in Martenot et al. (2012).

306 For Region 2, situated at ORF 43, nucleotide sequence was obtained from 77 of 100
307 samples. They were all different from the reference sequence and the microvariants. They were
308 separated into two groups, classified by one deletion of a thymine in the downstream region
309 (base 60,504) (G8) for 36 samples and two deletions of two adenine in the upstream region
310 (base 60,059 - 60,060) for 41 samples (G9) (Figure 2, Supplementary data 2).

311 In the third region coding for ORF 35 to ORF 38, nucleotide sequences were obtained
312 from 96 of 100 samples and all were identical to the two microvariant sequences. The features
313 included the large deletion zone of 606bp (base 52,251 - 52,856) common to microvariants and
314 the substitution of a cytosine instead of a thymine (base 52,885). They were classified into G10
315 (Figure 3, Supplementary data 2).

316 Eighty-seven of 100 samples presented an amplification of the fourth region located on
317 the non-coding site between ORF 49 and ORF 50. All the nucleotide sequences presented a
318 common substitution of an adenine instead of a guanine in the upstream zone (base 72,489) and
319 two deletions of two thymine next to the middle zone (base 75,592 - 75,593). However there
320 were many differences compared to the reference genotype and microvariants consisting of 19
321 mutations over 408 nucleotides which were not observed in the microvariant genotypes. These
322 mutations allowed the characterization of 14 groups (Figure 4, Supplementary data 2),
323 confirming the prior observations of high variability in this region (Batista et al., 2015). Four
324 samples of a nucleotide sequence named G12 presented a thymine as well as a cytosine (Y) at
325 the same position (base 72,536). The chromatograms of this and all the other regions of these
326 four samples comprising G12 were unambiguous. All samples which contained the G12 variant
327 were pools of 5 spat (Supplementary data 1).

328 For Region 5 sitting on ORF 11, nucleotide sequence was obtained from 42 of 100
329 samples. None had the large deletion observed in the microvariants. Even though the nucleotide
330 sequences were close to the reference sequence, a few mutations were present allowing their
331 classification into four groups (G31, G32, G33, G34) (Figure 5, Supplementary data 2). All had
332 a common substitution of a thymine instead of a guanine (base 18,621).

333 Of the 91 samples which amplified from Region 6 in the first part of ORF 88, 54
334 nucleotide sequences were similar to the reference sequence (G37). The others clustered into
335 two groups, comprising the substitution of a cytosine instead of a guanine (base 133,574) in 36
336 samples (G36) and a substitution of a guanine instead of an adenine (base 133,574) in one
337 sample (G35) (Figure 6, Supplementary data 2).

338 Nucleotide sequences for Regions 1, 2, 3, 4 and 6 were obtained from 61 samples, which
339 when concatenated led to the identification of 12 DNA variants (Table 4). V1 and V8 were the
340 most numerous of these variants.

341

342 3.1.2. Phylogenetic analyses

343 Firstly, a phylogenetic tree using Regions 1, 2, 3, 4 and 6 was constructed with the
344 concatenated nucleotide sequences of the 12 variants and the 29 whole genomes present in
345 NCBI. All Australian genotypes clustered in the same clade, distinct from isolates from Europe,
346 China and the reference genotype and with a distinctly shorter branch length (Figure 2).
347 Secondly, a phylogenetic tree was constructed based only on the C region (Region 1), which
348 has been well-studied internationally, in order to incorporate a wider range of isolates (Figure
349 3). A single cluster distinguished all Australian genotypes from those from Europe, New
350 Zealand, Japan and the Americas.

351

352 3.1.3 Diversity

353 The number of genotypes available for analysis had almost doubled from 48 to 95 since
354 the study of Mineur et al in 2015 (Table 5). The Australian isolates had the lowest genetic
355 diversity overall (Pi 0.99×10^{-3}) when compared to viruses from other regions of the world
356 where diversity ranged from 5.31×10^{-3} to 7.61×10^{-3} (Table 5).

357

358

359 3.1.3. Influencing factors

360 The distribution among the sites over time of the 12 DNA variants identified from
361 concatenated nucleotide sequences is illustrated in Table 6 while similar data for all genomic
362 regions is presented in Table 1, Supplementary data 2.

363 Considering the concatenated nucleotide sequences, up to four variants were found in
364 one summer season in the Georges River, compared to only one or two variants in the
365 Hawkesbury River. Nine of the 12 variants occurred at only one site and in only one summer
366 season. The variants V1 and V6 were present only in the Hawkesbury River; V1 was present at
367 most sites and in each of the four summers, while V6 was found only at Kimmerikong Bay and
368 Mooney Mooney in summer 2013-2014. The other variants were confined to the Georges River
369 where V8 was identified at various sites within Woollooware Bay in three summer seasons. The
370 other variants in the Georges River were detected only at a single site and a single sampling
371 date.

372 More than one variant was detected over time at some sites; for example, five were seen
373 at Site B in the Georges River while two were seen at both Kimmerikong Bay and Mooney
374 Mooney in the Hawkesbury river (Table 7). More than one variant was found in the same
375 summer season at some sites; for example, in 2014-2015 V8, V11 and V12 were found at Site
376 B, while in 2013-2014 both V1 and V6 were found at Kimmerikong Bay and at Mooney
377 Mooney. Furthermore, there were also several instances where different variants were detected

378 in the same population of oysters at the same time: V11 and V12 were both present at Site B
379 on 27/1/15; V1 and V6 were both present at Kimmerong Bay and Mooney Mooney on 13/12/13
380 and also on 25/1/14.

381 There were similar observations for Region 4 (Figure 4, Supplementary data 2), which
382 had the most variability; 12 of the 14 variants were found in only one season and at only one
383 site. Three of five variants in Region 1 and two of four variants in Region 5 were also seen in
384 only one season and at only one site.

385 Two isolates from Tasmania were not incorporated into the phylogenetic analysis
386 because of incomplete sequencing of regions from those samples. However one of these
387 samples was the only isolate classified as G6 in Region 1 (Table 1 and Figure 1, Supplementary
388 data 2); there were no sequencing data from this region of the other sample from Tasmania. The
389 oysters sampled in New South Wales had been supplied from at least nine different production
390 batches (i.e. different broodstock spawnings) from one hatchery in Tasmania. Oysters grown
391 from five of these batches yielded more than one variant, and in one case, five different OsHV-
392 1 variants, which has important implications when assessing the sources of the virus (see
393 discussion).

394

395 3.2. Amino acid sequence prediction of variants

396 3.2.1. Description of the predicted amino acid sequences

397 Predicted amino acid sequences were aligned with the reference and microvariant
398 sequences. All alignments are illustrated in Supplementary data 2. Regions 3 and 4 were not
399 included in this analysis because Region 3 had a large deletion of ORF 36 and 37, inducing
400 overlap of ORF 35 and 38 and creating uncertainty about whether a longer ORF could be
401 translated, while Region 4 was a non-coding area.

402 For Region 1, two amino acid sequences were identified for ORF 4 among the five
403 DNA sequences described above in part 3.1.1. They were both different to the reference
404 genotype and the microvariants. The first group was named <<A>> and contained G1, G2, G4
405 and G7; it had an asparagine (N) instead of an aspartic acid (D). The second group named
406 <> contained G6 and had in addition a leucine (L) instead of a serine (S) (Figure 7,
407 Supplementary data 2).

408 For Region 2, the two DNA groups characterised for ORF 43 presented two different
409 amino acid sequences. The first was named <<C>> and contained G8. It had a similar amino
410 acid sequence to the reference genotype and the microvariant. The second was named <<D>>.
411 It contained G9 and was characterised by three successive mutations of a serine (S), lysine (K)
412 and a deletion instead of a phenylalanine (F), a glutamic acid (E) and an arginine (R) (Figure
413 8, Supplementary data 2).

414 Region 5 located on ORF 11 had four DNA groups which translated into 3 amino acid
415 sequences. One was similar to OsHV-1 reference genotype, and was named « E ». The second
416 was named « F » and had a mutation of an asparagine (N) instead of a lysine (K). The third was
417 named « G » and had 4 substitutions of a lysine (K) (Figure 9, Supplementary data 2).

418 Region 6 coding for the first part of ORF 88 presented two amino acid sequences named
419 « H » containing G36 and G37 and « I » containing G35. <<H>> had a similar sequence to the
420 reference genotype and one of the microvariants, KY242785, while <<I>> had one mutation of
421 a glutamic acid (E) instead of lysine (K) (Figure 10, Supplementary data 2).

422 The amino acid prediction of the 12 DNA variants identified among 61 samples resulted
423 in two profiles based on ORF 4, ORF 43 and ORF88: ACH (n= 34) and ADH (n=27) (Table
424 8). Region 5 (ORF11) was not included because there were only 42 sequences available for it.

425

426 3.2.2. Influencing factors

427 The amino acid profile of isolates was different between the two estuaries in NSW
428 (Tables 6 and 7). The ACH profile was found only in the Hawkesbury River, an estuary located
429 40 km north of Sydney while the ADH profile was found only in the Georges River, an estuary
430 located 20 km south of Sydney.

431

432 **4. Discussion**

433

434 *Australian OsHV-1 isolates have lower diversity than OsHV-1 isolates globally*

435

436 It is difficult to compare genomic diversity between published studies of OsHV-1 due
437 to differences in the genomic regions sequenced, sample sizes, time span, geographic extent,
438 host species and disease state (healthy, sick, or dead). However, in this study we report diversity
439 of *Ostreid herpes virus-1* in estuaries near Sydney Australia that may be lower than in other
440 geographic regions of the world.

441 Regions of the OsHV-1 genome under selection pressure and other regions which
442 accumulate a high number of substitutions have been reported (Morga et al., 2021). Selection
443 pressure is less on a non-coding area and mutations acquired over time are less likely to be
444 eliminated. The diversity revealed in this study is mainly explained by inclusion of non-coding
445 Region 4 between ORF 49 and ORF 50, which is a polymorphic area (Batista et al., 2015). In
446 this region alone, 14 groups were identified among the 87 samples (one variant for every 6.2
447 isolates). Similarly, Batista et al. (2015) identified one variant for every 6.0 isolates examined
448 in a study of 18 isolates from mostly diseased *C. gigas* in Portugal.

449 C region, corresponding to Region 1 in the present study, is the most commonly
450 analysed part of the OsHV-1 genome. With the typical primers C2/C6 used for amplification
451 there is representation of part of a non-coding region and part of ORF 4 (Batista et al., 2015).

452 Nucleotide diversity in the C region was calculated formally and compared to data published in
453 2015 (Mineur et al., 2015). There were more sequences in public databases, but the nucleotide
454 diversity had not changed markedly except for Europe where it appeared to have decreased
455 (from 15.3×10^{-3} to 7.6×10^{-3}). The Australian isolates had the lowest genetic diversity (P_i 0.99
456 $\times 10^{-3}$) overall. Globally, 48 different genotypes had been identified in C region (including the
457 microsatellite region) by 2015 (Mineur et al., 2015). Expressed in simple terms of the frequency
458 of genotypes among isolates examined, in Australia seven genotypes have been identified
459 among 89 samples in diseased *C. gigas* (five from 87 isolates in the present study and two
460 existing in NCBI), which equates to one variant for every 12.7 isolates. In contrast, in a study
461 covering a wide area of coastal Japan, 23 genotypes were found among 123 samples from
462 healthy *Crassostrea* species (one variant for every 5.4 isolates) (Shimahara et al., 2012); in both
463 countries samples were collected over about 5 years. The genetic diversity among diseased
464 oysters in Australia was clearly much less than that in healthy oysters in Japan. The data from
465 Japan in the north west Pacific region likely represent the diverse environmental reservoir of
466 viral types detectable in healthy shellfish, while the low diversity in Australia may represent
467 selected environmental strains of OsHV-1 capable of causing disease in farmed oysters under
468 certain environmental conditions, that then became locally dominant. Relatively low diversity
469 was seen in Europe after 2008 when OsHV-1 *uVar* became the most common strain in diseased
470 oysters (Table 5). In Europe there had been spread of the virus with movements of live oysters
471 for farming, for example from France to Ireland and Jersey (Peeler et al., 2012). Within France,
472 OsHV-1 *uVar* was dispersed widely with commercial movements of live oysters from
473 Marennes-Oléron to farming areas on both the Atlantic and Mediterranean coasts (Delmotte et
474 al., 2022). Thus, in addition to the health/disease condition of oysters and their geographic
475 locations, the observed regional differences in the diversity of OsHV-1 genotypes depends on

476 epidemiological differences between the sources and modes of spread of the viruses. All these
477 factors will need to be considered in future studies of the diversity of OsHV-1.

478

479 In Region 4 we observed a double nucleotide (thymine and cytosine) in four individual
480 samples for which remaining nucleotide sequences were unambiguous (G12 – Figure 4, and
481 Supplementary data 2) and otherwise matched the nucleotide sequence of G18 which had a
482 cytosine at this position (Figure 4, Supplementary data 2). Our first hypothesis is that there has
483 been a recent mutation at this position to explain why we found two different copies of this part
484 of the genome, one being the ancestral form and the other being the mutated one. This is feasible
485 due to the persistence of the herpesvirus in surviving oyster hosts (Evans et al., 2017b), because
486 gene exchange in herpesviruses can occur by recombination (Thiry et al., 2005) and because
487 Rosani and Venier (2017) showed the presence of DNA recombination-initiating promoter
488 binding in *Malacoherpesviridae*. The four samples with this mutation were from oysters in
489 close proximity to one another and they could have been infected almost simultaneously from
490 a host in which the mutation first occurred, resulting in a pool of two variants in each of the four
491 samples. This is consistent with Martenot et al (2011) who reported various genotypes in the
492 same batch of oysters and Morga et al (2021) who reported various genotypes in the same
493 individual. A second possibility would be that there were two otherwise identical but long-
494 standing OsHV-1 variants in each of the four samples, due to different individuals containing a
495 different genotype of OsHV-1, because each sample came from a pool of five individual
496 oysters. While this does increase the chance of mixing several variants into one DNA extract,
497 in no other sample did we find a group with a thymine at this position. The least likely
498 explanation because the non-coding area between ORF 49 and ORF 50 is present as a single
499 copy in the complete OsHV-1 genomes available on NCBI, is the existence of a double copy
500 of this region.

501

502 The microsatellite zone of the C region of OsHV-1 is polymorphic because of repeated
503 trinucleotides which allow replication mistakes, as observed by many authors (Martenot et al.,
504 2012; Martenot et al., 2011; Mineur et al., 2015; Renault et al., 2014). The range in the number
505 of ACT trinucleotide repeats across all genotypes globally was reported to be 3 - 13 (Mineur et
506 al., 2015), corresponding to a range of 3 - 10 trinucleotide deletions (9 - 30 base deletions).
507 However, we observed only two profiles among a large number of isolates, either 12 or 15 base
508 deletions. This is consistent with the only other Australian sequences deposited in NCBI; these
509 came from the first episode of mortality in 2010 and had 12 base deletions.

510

511 *Australian isolates differed from OsHV-1 isolates globally*

512 Concatenated nucleotide sequences showed a distinct cluster including all the Australian
513 isolates, different from all other genotypes reported worldwide and supported by the distinct
514 distribution with a shorter branch size for the Australian genotypes (Figure 2). Although the
515 bootstrap value was only 63, this could be explained by the presence of Region 3 located
516 between ORF 35 and 38 including a large deletion area and region 5 located on ORF 11, with
517 a 462bp deletion only in the microvariant. These differences are calculated as multiple mutation
518 events when it is more likely that each corresponds to a single mutation event. Burioli et al.
519 (2016) studied the same disturbed region and discussed the need for “sophisticated algorithms
520 for alignment and phylogenetical analysis, adapted to take into account DNA-virus evolution
521 analysis, in order to obtain accurate results with the correct phylogenetic distance”. The
522 Australian clade was also confirmed by the inclusion of the only two NCBI sequences from
523 Australia, KC685525.1 and MT157286.1 (Burge et al., 2020; Jenkins et al., 2013) in the C
524 region tree (Figure 3). This is consistent with the phylogeny reported in Mineur et al. (2015),
525 which was performed on the same locus with another algorithm, median joining genotype, and

526 showed three closely related genotypes from temperate Australasia. In contrast, the
527 phylogenetic tree presented in Burge et al. (2021) with the neighbour joining method showed
528 that the Australian genotype was close to the Japanese, American, Korean and French
529 microvariants. However, unlike the current phylogeny, the tree described in Burge et al. (2021)
530 is not based on evolutionary distance and contains too few samples - a single genotype from
531 Australia - which is not sufficient to construct a cluster. The Australian viruses appear to be
532 genetically different across the genomic regions analyzed and this raises the question about
533 their true phylogenic origins. Sanger sequencing of a limited number of genomic regions may
534 bias analyses compared to whole genome analysis, which will be indispensable for determining
535 clades of OsHV-1 and should be the subject of future studies.

536

537 *Australian isolates were not OsHV-1 reference strain, OsHV-1 μ Var or microvariants*

538 Neither OsHV-1 reference genotype nor OsHV-1 μ Var were identified among the
539 Australian samples. Currently OsHV-1 “microvariants” are described as being close to OsHV-
540 1 μ Var as defined by Segarra et al. (2010) but not identical to it (Burge et al., 2021; Morga et
541 al., 2021). In all of the Australian isolates the two regions which characterise OsHV-1 μ Var
542 according to Segarra et al. (2010), ORF 4 and ORF 43, had dissimilar nucleotide sequences.
543 None of these sequences had the large deletion in ORF 11 reported in the microvariant
544 sequences (Burioli et al., 2017; Martenot et al., 2013; Morga et al., 2021) and presented a similar
545 nucleotide sequence to the OsHV-1 reference virus on ORF 11 except for one point mutation.
546 Moreover, half the samples from Australia did not present an amplification product from this
547 region. Since there was enough biological material in the same samples to perform sequencing
548 of the other regions, we suggest that there were changes at the primer binding sites, which others
549 have also mentioned to be a likely reason for this problem (Arzul et al., 2001b; Martenot et al.,

550 2011); PCR-sequencing of the other regions was not uniformly successful either, consistent
551 with many other studies of OsHV-1.

552

553 The variant of OsHV-1 known as μ Var is thought to be responsible for the sudden
554 escalation of mass mortality events in *C. gigas* in France from 2008, and then later in other
555 European countries. However, μ Var and the more recently described “microvariants” are
556 clearly not the only variants of OsHV-1 associated with mass mortality events in *C. gigas*. The
557 disease outbreaks in both Australia since 2010 have not been caused by these types of OsHV-1
558 but have been just as sudden in onset, recurrent and economically devastating for the oyster
559 industry as those in France (Fuhrmann et al., 2019). Therefore further research is required to
560 understand the virulence determinants of OsHV-1 and how these are coded in the genome.

561

562 *Internal comparison of Australian isolates*

563

564 The two estuaries studied in New South Wales had different OsHV-1 profiles. At the
565 Georges River sites, ten variants were identified over five summer seasons while at the
566 Hawkesbury River sites only two variants were found. Delmotte et al. (2022) showed a great
567 diversity of variants in areas with high oyster farming activity. However, the causes of the
568 different patterns of distribution of variants here are unknown since these two estuaries in
569 Australia had equivalent, intense oyster farming activity using similar genetic stock of *C. gigas*
570 from the same OsHV-1 free hatchery in Tasmania. As a consequence of the two distributions
571 of DNA variants, the amino acid sequences of the viruses present in the Hawkesbury and
572 Georges river estuaries were different: the ACH profile in the Hawkesbury River and ADH in
573 the Georges River. This distinct pattern has a precedent in Europe, but with far greater
574 geographic separation than the 60 km that exists between these two estuaries. Based on RNA-

575 Seq SNP analysis, Delmotte et al. (2020) reported two genetically different OsHV-1
576 populations in POMS outbreaks in *C. gigas* on the Atlantic and Mediterranean coasts of France;
577 they had amino acid sequence differences in viral membrane proteins, which may affect
578 virulence (Delmotte et al., 2020).

579

580 Due to the selection of samples based on high viral load, which in general is correlated
581 with mortality (Oden et al., 2011), it was not appropriate to compare the OsHV-1 isolates in
582 this study according to their degree of replication fitness or observed mortalities. However, in
583 the oyster populations from which the samples were collected there were differences in the
584 patterns of mortality: spat in the Georges River estuary were 6.4 times more likely to have
585 mortality due to OsHV-1 than those in the Hawkesbury River estuary (Whittington et al., 2019).
586 Subclinical infection of oysters with OsHV-1 was also more likely in the Georges River (Evans
587 et al., 2017b; Whittington et al., 2019). It is possible that the mutually exclusive genotypic
588 variants of OsHV-1 identified between the two estuaries differed in virulence. However, other
589 factors could also be involved. While oyster genotype was similar between the two rivers
590 because of the common origin of spat, there could be a wide range of environmental differences.
591 These could include differences in the co-occurrence of reservoirs/hosts, particulate vectors,
592 filter feeding species and seaweeds that could all influence the severity of POMS (Dugeny et
593 al., 2022; Evans et al., 2017c; Pernet et al., 2021; Whittington et al., 2018).

594

595 To our knowledge there were no transfers of commercial oysters between the Georges
596 and Hawkesbury Rivers after first detection of OsHV-1 in the Georges River late in 2010 due
597 to government biosecurity directives (Paul-Pont et al., 2014). The only common factor between
598 the two oyster production locations was the recruitment site of all batches of farmed *C. gigas*
599 in this study: Tasmania. However, the current data indicate that the virus contamination did not

600 come from this recruitment step but from the culture environment. There are three reasons for
601 this : i) consistent pre-shipment PCR testing of all batches of spat from the hatchery by the
602 government laboratory in Tasmania and again upon arrival in NSW by the researchers (all
603 negative results) means that spat were specific-pathogen free (SPF); ii) different genotypes of
604 OsHV-1 were detected in the same batch of spat after deployment in NSW; iii) a unique
605 genotype (G6) of OsHV-1 was detected in Tasmania; G6 was not found among 85 samples
606 from the Hawkesbury River or Georges River. Epidemiological observations alone had
607 suggested that recurrent POMS outbreaks in Australia were derived from the environment and
608 were not due to disease transfer with oysters during farming (Whittington et al., 2018), a feature
609 which is now strongly supported by DNA sequence evidence. This scenario is in direct contrast
610 to France where evidence points to a radiation of disease associated with commercial spat
611 distribution (Delmotte et al., 2022). The time of first appearance of OsHV-1 in a putative
612 environmental host in Australia cannot be determined, but it could have been a point source
613 introduction to a naive population, then adaptation to local conditions. In fact there are historical
614 records of a virus like Ostreid herpesvirus from *Ostrea angasi* in Western Australia (Hine and
615 Thorne, 1997). These authors demonstrated herpes-like viral particles by electron microscopy,
616 but did not have the molecular tools to identify the virus. In any case, this observation raises
617 the possibility that OsHV-1 could be either indigenous to Australia or could have been
618 introduced many decades ago and become endemic, and it could have mutated to form its own
619 lineage over time, as is supported by the current phylogenetic analysis. Knowing that OsHV-1
620 μ Var emerged around 2008 in Europe and then dispersed rapidly, the absence of OsHV-1 μ Var
621 in Australia suggests that there has been effective compliance with the strict quarantine laws
622 that aim to prevent introduction of exotic pathogens into Australia.

623 Arzul et al. (2001a) speculated 20 years ago that pathogenic forms of OsHV-1 arose
624 from possibly benign bivalve herpesviruses which could be transmitted between bivalve species

625 and which would be favoured by modern shellfish farming practices (Arzul et al., 2001a; Arzul
626 et al., 2001b). In Australia, there is evidence that invertebrate species in the Georges River
627 besides *C. gigas* are exposed to OsHV-1 and may become infected, but their role in the
628 epidemiology of POMS remains to be determined (Evans et al., 2017c). The environment is an
629 important factor in the transmission cycle because OsHV-1 can be found in seawater and
630 sediments (Evans et al., 2014 ;2017) and is likely transported on suspended particles over 5µm
631 (Evans et al., 2017a; Evans et al., 2017c; Paul-Pont et al., 2013b; Whittington et al., 2015b).
632 Depending on conditions such as temperature, the virus is expected to persist in seawater and
633 remain infectious for several days according to *in vitro* experimentation (Hick et al., 2016;
634 Martenot et al., 2015). This gives the virus time to transit between hosts through the
635 environment. *Ostreid herpesvirus 1* can infect several different host species (Arzul et al., 2001a;
636 Bai et al., 2016; Batista et al., 2015; Bookelaar et al., 2018; Comps and Cochenec, 1993; Evans
637 et al., 2017c; Farley et al., 1972; Hine and Thorne, 1997; Hine et al., 1998; Prado-Alvarez et
638 al., 2021; Renault and Arzul, 2001; Renault et al., 2001; Xia et al., 2015) and may persist in
639 survivors of outbreaks or remain latent (Arzul et al., 2002; Evans et al., 2017b), which may
640 explain why each environment has its own assemblage of viruses. While the molecular evidence
641 strongly suggested that the virus contamination came from reservoirs in the local culture
642 environment rather than a point source such as an oyster hatchery, the distribution of the
643 variants across sites and time suggests that some reservoirs were persistent and widespread,
644 while others were highly site and time specific and therefore possibly short-lived, or that
645 transmission was infrequent. The reservoirs may be subject to local environmental influences
646 which affect their distribution and abundance. Further field research is required to discover the
647 environmental reservoir of OsHV-1 and the specific environmental factors that facilitate
648 emergence of the virus. Climate change, short term climatic and water quality variations and
649 intensification of farming are among the many factors to be considered.

650

651

652 **5. Conclusion**

653

654 Sequencing analyses of OsHV-1 isolates from estuaries near Sydney, Australia,
655 revealed a low diversity of genotypes relative to other geographic regions, reflecting differences
656 in sources and modes of spread of the viruses and the types of oyster populations (healthy, or
657 diseased) that have been examined so far. Importantly, Australian POMS outbreaks were
658 associated with a type of OsHV-1 distinct from others globally and were not a “microvariant”
659 genotype. Elsewhere, since 2008 OsHV-1 microvariants have been the only variants associated
660 with mass mortality events in *C. gigas*. Future whole genome sequencing investigations will
661 allow more detailed classification of Australian isolates of OsHV-1. Based on both DNA and
662 amino acid sequences, OsHV-1 variants were absolutely correlated to estuary. The *C. gigas*
663 populations in the two estuaries near Sydney had different prevalence and severity of POMS,
664 which suggests that the geographically mutually exclusive genotypes differed in virulence. The
665 molecular evidence supports field epidemiological evidence that the disease-causing virus came
666 from reservoirs in the local environment. The sequencing findings in this study help inform
667 understanding of the various sources and modes of spread of OsHV-1 that have led to mass
668 mortality events in farmed *C. gigas* globally.

669

670 **Author contributions**

671 Suzanne Trancart: Investigation, Data curation, Methodology, Visualization , Formal analysis,
672 Writing - Original Draft, Writing - Review & Editing. Alison Tweedie: Investigation, Data
673 curation, Writing - Review & Editing. Olivia Liu: Investigation, Writing - Review & Editing.
674 Ika Paul-Pont: Conceptualization, Investigation, Writing - Review & Editing. Paul Hick:
675 Investigation, Writing - Review & Editing. Maryline Houssin: Resources, Investigation,
676 Supervision, Writing - Review & Editing. Richard Whittington: Funding acquisition, Project

677 administration, Supervision, Resources, Conceptualization, Investigation, Data curation,
678 Formal analysis, Writing - Original Draft, Writing - Review & Editing.

679

680

681 **Acknowledgements**

682 Sample collection and sequencing was funded by Fisheries Research and Development
683 Corporation Australia under projects 2011/053, 2012/032 and 2014/040 and The University of
684 Sydney and LABÉO France. Sanger sequencing was undertaken by Australian Genome
685 Research Facility and Monash Health Translation Precinct Medical Genomics Facility
686 Australia. Bruce Allford, Len Drake, Bob Drake and Robert Hill are thanked for their invaluable
687 assistance with field work. Shellfish Culture Tasmania kindly provided all batches of spat that
688 were deployed and sampled in cited research trials. Vickie Patten and Ann-Michele Whittington
689 provided skilled laboratory assistance with oyster processing.

690

691

692

693

694

695

696 **Captions to figures**

697 Figure 1 : Sampling sites. George river estuary sites are located on the bottom map
698 corresponding to the south of Sydney and Hawkesbury river estuary sites are located on the
699 upper map corresponding to the north of Sydney.

700

701 Figure 2: Distribution of the DNA variants according to the NCBI deposited nucleotide
702 sequences. The unrooted phylogenetic tree was performed using the neighbour joining
703 method and 1000 bootstrap replications were applied based on maximum composite
704 likelihood. Regions 1,2,3,4 and 6 were concatenated on 61 Australian isolates showing 12
705 variants (V1, V2, V3, V4, V5, V6, V7, V8, V9, V10, V11, V12). Yellow rectangle is the
706 reference nucleotide sequence of OsHV-1. Metadata for the nucleotide sequences of the
707 current study that are included in the tree are in Supplementary data 1 while metadata for the
708 nucleotide sequences that are already available in NCBI are in Table 1, Supplementary data
709 3.

710 Figure 3: Distribution of the DNA variants according to nucleotide sequences deposited in
711 NCBI. The unrooted phylogenetic tree was performed using the neighbour joining method
712 and 1000 bootstrap replications were applied based on maximum composite likelihood.
713 Region 1 from 87 Australian isolates was used, showing 5 groups (G1, G2, G4, G6, G7).
714 Crosses indicate Australian nucleotide sequences extant in NCBI; asterisks indicate the New
715 Zealand nucleotide sequences. Yellow rectangle is the reference nucleotide sequence of
716 OsHV-1. Metadata for the nucleotide sequences of the current study that are included in the
717 tree are in Supplementary data 1 while metadata for the nucleotide sequences that are already
718 available in NCBI are in Table 2, Supplementary data 3.

719

720 **List of supplemental data**

721 *Supplemental data 1*

722 Table 1 Metadata for all samples

723 Table 2 Citations of scientific publications describing the populations of oysters from which
724 samples were collected

725

726 *Supplemental data 2*

727 Table 1 Variants by genomic region and geographic location

728 Figures 1 to 10 DNA and amino acid alignments

729

730 *Supplemental data 3*

731 Genbank accession numbers of isolates included in phylogenetic trees

732
733
734

Table 1 : Number of samples of OsHV-1 PCR-positive oysters from which nucleotide sequences were analysed according to the estuary and the season of collection. The number of sites sampled are in parentheses; sites differed between seasons.

Estuary	2011-2012	2012-2013	2013-2014	2014-2015	2015-2016	Total
Georges River NSW	19 (3)	9 (3)	14 (4)	13 (3)	5 (3)	60
Hawkesbury River NSW		11 (3)	15 (2)	9 (2)	3 (2)	38
Pitt Water TAS					2 (1)	2
Total	19	20	29	22	10	100

735
736

737
738
739

Table 2 : Regions of the OsHV-1 genome targeted for PCR amplification.

Sequencing region	Amplification target				Amplicon size (base pairs)	Reference	
	Position ^a	Gene	Primer name and purpose ^b	Primer sequence (5'→3')			
1 (C region)	178211 178919	ORF 4	C2 (A, S)	CTCTTTACCATGAAGATACCCACC	709	(Renault and Arzul, 2001)	
			C6 (A, S)	GTGCACGGCTTACCATTTTT			
			C1585 (S)	GTATAAATAGGCGCG			This paper
			C1586 (S)	CGCGCCTATTTATAC			This paper
			C1587 (S)	CAGACGAGGTTAAC			This paper
			C1588 (S)	GTAAACCTCGTCTG			This paper
			C1593 (S)	GGAGCTGCGGCGCTATG			This paper
2 (A region)	59950 60557	ORF 42-43	IA1 (A, S)	CGCGGTTTCATATCCAAAGTT	607	Segarra et al. (2010)	
			IA2 (A, S)	AATCCCCATGTTTCTTGCTG			
			C1589 (S)	CTTGCTCATCGTATTC			This paper
3 (605bpDel)	51979 52968	ORF 35 -36 - 37 and 38	Del 36-37F2 (A, S)	ATACGATGCGTCGGTAGAGC	989, 384 or no amplification	(Renault et al., 2012)	
			Del 36-37R (A, S)	CGAGAACCCCATTCCTGTAA			
			C1590 (S)	CATGGTGATGAATGAAG			This paper
			C1591 (S)	CTTCATTCATCACCATG			This paper
			C1592 (S)	CATTCCTGTAAACAC			This paper
4 (NC1NC2 region)	72414 72919	Between ORF49 and 50	NC1 (A, S)	ACACCTAATGACCCCAAAGG	506	(Batista et al., 2015)	
			NC2 (A, S)	GACCAATCACCAGCTCAACA			
5	17402 19518	ORF 11	ORF11For (A, S)	ACCACCGCGCCAAAATCTG	2116 or 731	(Martenot et al., 2013)	
			ORF11Rev (A, S)	CGCTTCCTATCACCTTGTGG			
			C1872 (S)	CTAGTCGTGCTCGTTCCTCTGC			This paper
			C1873 (S)	GGCAGAGATAGAACAATG			This paper
			C1874 (S)	CATTGTGTTCTATCTCTGCC			This paper
6	133088 134107	ORF 88a	ORF88aFor (A,S)	CCCAGTCTATTATCCAGGTAC	1020	(Martenot et al., 2013)	
			ORF88aRev (A,S)	ACCGTTCCTCAATCAGTCCC			

740

741 ^a Position on the OsHV-1 genome relative to the OsHV-1 reference genome, GenBank: AY509253

742 ^b Purpose: A amplification, S sequencing

743

744
745
746

Table 3: GenBank accession numbers of the 29 DNA groups of the 6 regions analysed.

Region	Gene	Sequence Group	GenBank Accession No.				
1	C region	ORF4	G1	ON953753			
			G2	ON953754			
			G4	ON953755			
			G6	ON953752			
			G7	ON953756			
2	A region	ORF 42-43	G8	ON953757			
			G9	ON953758			
3	605 bpDel	ORF 35 to 38	G10	ON953759			
			G11	ON954001			
			G12	ON954014			
			G13	ON954007			
			G14	ON954004			
			G15	ON954003			
			G17	ON954002			
			4	NC2 region	Non-coding, between ORF 49 and ORF 50	G18	ON954013
						G19	ON954012
						G20	ON954006
						G21	ON954010
						G22	ON954009
						G23	ON954008
G24	ON954005						
5	ORF 11 gene	ORF 11	G29	ON954011			
			G31	ON953762			
			G32	ON953760			
			G33	ON953763			
			G34	ON953761			
6		ORF 88	G35	ON953765			
			G36	ON953764			
			G37	ON953766			

747

748
749

Table 4: Description of the 12 genotypic variants based on concatenated nucleotide sequences of Regions 1,2,3,4 and 6 for 61 isolates according to their group identification numbers for these regions.

Variant	n	R1	R2	R3	R4	R6
		ORF 4	ORF 43	ORFs 35, 36, 37, 38	uncoded site between ORF49 and 50	ORF 88
V1	32	G7	G8	G10	G11	G36
V2	3	G1	G9	G10	G11	G37
V3	1	G7	G9	G10	G20	G37
V4	1	G7	G9	G10	G21	G37
V5	1	G7	G9	G10	G29	G37
V6	2	G7	G8	G10	G11	G37
V7	5	G7	G9	G10	G13	G37
V8	10	G7	G9	G10	G11	G37
V9	2	G4	G9	G10	G12	G37
V10	2	G7	G9	G10	G14	G37
V11	1	G7	G9	G10	G15	G37
V12	1	G7	G9	G10	G17	G37

750

751

752

753

754 *Table 5: Nucleotide diversity in the C region (including the microsatellite region) of OsHV-1 from different geographical*
755 *regions. The data are based on sequences listed in Table 2 of Supplementary data 3 as well as sequences from the current*
756 *study. Six sequences were added to be consistent with Mineur et al (2015) (see materials and methods). NW Pacific includes*
757 *isolates from China, Japan and Korea; NE Pacific includes Mexico and USA; Europe includes France, Ireland, Italy,*
758 *Portugal and Spain; Australasia includes Australia and New Zealand.*

Geographic region	Current study		Mineur et al (2015)	
	No. of genotypes	Nucleotide diversity (P_i)	No. of genotypes	Nucleotide diversity (P_i)**
All	95	7.46×10^{-3}	48	14.97×10^{-3}
NW Pacific	45	6.45×10^{-3}	27	9.34×10^{-3}
Australasia	10	1.49×10^{-3}	3	2.09×10^{-3}
Australia	7	0.99×10^{-3}	nr	nr
NE Pacific	5	6.24×10^{-3}	2	6.27×10^{-3}
Europe all years	41	7.61×10^{-3}	16	15.30×10^{-3}
*Europe < 2008	9	3.43×10^{-3}	10	4.67×10^{-3}
Europe \geq 2008	32	2.07×10^{-3}	6	3.45×10^{-3}
France	27	5.31×10^{-3}	nr	nr

759

760 *One sequence (AY459363) with a high frequency of undetermined nucleotides (“N”) was
761 excluded from the current study

762 **corrected for $H_d=1$, see materials and methods

763 nr, no result

764

771
772

Table 7: Distribution of 12 DNA variants and protein prediction profiles among 61 samples from ten different sites in the Georges and Hawkesbury river estuaries between 2011 and 2016

Estuary	Site	No. samples	DNA variant	Predicted protein
Hawkesbury river (n=34)	Coba Bay	1	V1	ACH
	Kimmerikong Bay	4	V1 (n=3) V6 (n=1)	ACH
	Mooney Mooney	15	V1 (n=14) V6 (n=1)	ACH
	Mullet Creek	8	V1	ACH
	Porto Bay	6	V1	ACH
George river (n=27)	Neverfail	2	V10	ADH
	site A	2	V4 V8	ADH
	site B	5	V3 V5 V8 V11 V12	ADH
	site C	13	V8 (n=8) V9 (n=2) V2 (n=3)	ADH
	The shed	5	V7	ADH

773
774

775
776
777

Table 8: Protein profiles of the 12 DNA variants identified among 61 samples from ten different sites in the Georges and Hawkesbury river estuaries between 2011 and 2016

Variant	n	ORF 4	ORF 43	ORF 88
V1	32	A	C	H
V2	3	A	D	H
V3	1	A	D	H
V4	1	A	D	H
V5	1	A	D	H
V6	2	A	C	H
V7	5	A	D	H
V8	10	A	D	H
V9	2	A	D	H
V10	2	A	D	H
V11	1	A	D	H
V12	1	A	D	H

778
779
780

781

782 **References**

783

784

785 Abbadi, M., Zamperin, G., Gastaldelli, M., Pascoli, F., Rosani, U., Milani, A., Schivo, A.,
786 Rossetti, E., Turolla, E., Gennari, L., Toffan, A., Arcangeli, G., Venier, P., 2018.
787 Identification of a newly described OsHV-1 mu var from the North Adriatic Sea
788 (Italy). *Journal of General Virology* 99(5), 693-703.

789 <https://doi.org/10.1099/jgv.0.001042>

790 Arzul, I., Nicolas, J.L., Davison, A.J., Renault, T., 2001a. French scallops: A new host for
791 ostreid herpesvirus-1. *Virology* 290(2), 342-349.

792 <https://doi.org/10.1006/viro.2001.1186>

793 Arzul, I., Renault, T., Lipart, C., Davison, A.J., 2001b. Evidence for interspecies transmission
794 of oyster herpesvirus in marine bivalves. *Journal of General Virology* 82, 865-870.

795 <https://doi.org/10.1099/0022-1317-82-4-865>

796 Arzul, I., Renault, T., Thébault, A., Gérard, A., 2002. Detection of oyster herpesvirus DNA
797 and proteins in asymptomatic *Crassostrea gigas* adults. *Virus Res* 84(1-2), 151-160.

798 [https://doi.org/10.1016/s0168-1702\(02\)00007-2](https://doi.org/10.1016/s0168-1702(02)00007-2)

799 Bai, C., Gao, W., Wang, C., Yu, T., Zhang, T., Qiu, Z., Wang, Q., Huang, J., 2016.

800 Identification and characterization of *Ostreid herpesvirus 1* associated with massive
801 mortalities of *Scapharca broughtonii* broodstocks in China. *Dis Aquat Org* 118(1), 65-

802 75. <https://doi.org/10.3354/dao02958>

803 Bai, C.M., Morga, B., Rosani, U., Shi, J., Li, C., Xin, L.S., Wang, C.M., 2019. Long-range
804 PCR and high-throughput sequencing of *Ostreid herpesvirus 1* indicate high genetic
805 diversity and complex evolution process. *Virology* 526, 81-90.

806 <https://doi.org/10.1016/j.virol.2018.09.026>

807 Batista, F.M., López-Sanmartín, M., Grade, A., Morgado, I., Valente, M., Navas, J.I., Power,
808 D.M., Ruano, F., 2015. Sequence variation in *Ostreid herpesvirus 1* microvar isolates
809 detected in dying and asymptomatic *Crassostrea angulata* adults in the Iberian
810 Peninsula: Insights into viral origin and spread. *Aquaculture* 435(0), 43-51.

811 <http://dx.doi.org/10.1016/j.aquaculture.2014.09.016>

812 Bookelaar, B.E., O'Reilly, A.J., Lynch, S.A., Culloty, S.C., 2018. Role of the intertidal
813 predatory shore crab *Carcinus maenas* in transmission dynamics of *Ostreid*

814 *herpesvirus-1* microvariant. Dis Aquat Org 130(3), 221-233.
815 <https://doi.org/10.3354/dao03264>

816 Burge, C.A., Friedman, C.S., Kachmar, M.L., Humphrey, K.L., Moore, J.D., Elston, R.A.,
817 2021. The first detection of a novel OsHV-1 microvariant in San Diego, California,
818 USA. J Invert Pathol 184. <https://doi.org/10.1016/j.jip.2021.107636>

819 Burge, C.A., Reece, K.S., Dhar, A.K., Kirkland, P., Morga, B., Degremont, L., Faury, N.,
820 Wippel, B.J.T., MacIntyre, A., Friedman, C.S., 2020. First comparison of French and
821 Australian OsHV-1 mu vars by bath exposure. Dis Aquat Org 138, 137-144.
822 <https://doi.org/10.3354/dao03452>

823 Burioli, E.A.V., Prearo, M., Houssin, M., 2017. Complete genome sequence of *Ostreid*
824 *herpesvirus* type 1 microVar isolated during mortality events in the Pacific oyster
825 *Crassostrea gigas* in France and Ireland. Virology 509, 239-251.
826 <https://doi.org/10.1016/j.virol.2017.06.027>

827 Burioli, E.A.V., Varello, K., Lavazza, A., Bozzetta, E., Prearo, M., Houssin, M., 2018. A
828 novel divergent group of *Ostreid herpesvirus 1* mu Var variants associated with a
829 mortality event in Pacific oyster spat in Normandy (France) in 2016. J Fish Dis
830 41(11), 1759-1769. <https://doi.org/10.1111/jfd.12883>

831 Cochenec-Laureau, N., Baud, J., Bedier, E., Boudry, P., Huvet, A., Nicolas, J., Pepin, J.,
832 Petton, B., 2010. Bilan des « Journées Surmortalité des huîtres creuses, *Crassostrea*
833 *gigas* » du Programme P7 « Aquaculture Durable » des 8 et 9 décembre 2009 (Review
834 of "Oysters *Crassostrea gigas* mortality days" Program P7 "Sustainable Aquaculture"
835 December 8 - 9, 2009), <https://archimer.ifremer.fr/doc/00000/7393/>.

836 Comps, M., Cochenec, N., 1993. A Herpes-like virus from the European oyster *Ostrea*
837 *edulis* L. J Invert Pathol 62(2), 201-203. <http://dx.doi.org/10.1006/jipa.1993.1098>

838 Davison, A.J., Eberle, R., Ehlers, B., Hayward, G.S., McGeoch, D.J., Minson, A.C., Pellett,
839 P.E., Roizman, B., Studdert, M.J., Thiry, E., 2009. The order Herpesvirales. Archives
840 of Virology 154(1), 171-177. <https://doi.org/10.1007/s00705-008-0278-4>

841 Davison, A.J., Trus, B.L., Cheng, N.Q., Steven, A.C., Watson, M.S., Cunningham, C., Le
842 Deuff, R.M., Renault, T., 2005. A novel class of herpesvirus with bivalve hosts.
843 Journal of General Virology 86, 41-53. <https://doi.org/10.1099/vir.0.80382-0>

844 de Kantzow, M., Hick, P., Becker, J., Whittington, R., 2016. Effect of water temperature on
845 mortality of Pacific oysters *Crassostrea gigas* associated with microvariant *Ostreid*
846 *herpesvirus 1* (OsHV-1 μ Var). Aquaculture Environ Interact 8, 419-428.

847 de Kantzow, M.C., Hick, P.M., Dhand, N.K., Whittington, R.J., 2017. Risk factors for
848 mortality during the first occurrence of Pacific Oyster Mortality Syndrome due to
849 *Ostreid herpesvirus - I* in Tasmania, 2016. Aquaculture 468, 328-336.
850 <https://doi.org/10.1016/j.aquaculture.2016.10.025>

851 de Lorgeril, J., Lucasson, A., Petton, B., Toulza, E., Montagnani, C., Clerissi, C., Vidal-
852 Dupiol, J., Chaparro, C., Galinier, R., Escoubas, J.M., Haffner, P., Degremont, L.,
853 Charriere, G.M., Lafont, M., Delort, A., Vergnes, A., Chiarello, M., Faury, N., Rubio,
854 T., Leroy, M.A., Perignon, A., Regler, D., Morga, B., Alunno-Bruscia, M., Boudry, P.,
855 Le Roux, F., Destoumieux-Garzon, D., Gueguen, Y., Mitta, G., 2018. Immune-
856 suppression by OsHV-1 viral infection causes fatal bacteraemia in Pacific oysters. Nat
857 Commun 9. ARTN 4215
858 10.1038/s41467-018-06659-3

859 Delmotte, J., Chaparro, C., Galinier, R., de Lorgeril, J., Petton, B., Stenger, P.L., Vidal-
860 Dupiol, J., Destoumieux-Garzon, D., Gueguen, Y., Montagnani, C., Escoubas, J.M.,
861 Mitta, G., 2020. Contribution of viral genomic diversity to oyster susceptibility in the
862 Pacific Oyster Mortality Syndrome. Front Microbiol 11, 1579.
863 <https://doi.org/10.3389/fmicb.2020.01579>

864 Delmotte, J., Pelletier, C., Morga, B., Galinier, R., Petton, B., Lamy, J.B., Kaltz, O., Avarre,
865 J.C., Jacquot, M., Montagnani, C., Escoubas, J.M., 2022. Genetic diversity and
866 connectivity of the *Ostreid herpesvirus I* populations in France: A first attempt to
867 phylogeographic inference for a marine mollusc disease. Virus Evolution 8(1).
868 <https://doi.org/10.1093/ve/veac039>

869 Dugeny, E., de Lorgeril, J., Petton, B., Toulza, E., Gueguen, Y., Pernet, F., 2022. Seaweeds
870 influence oyster microbiota and disease susceptibility. J. Anim. Ecol. 91(4), 805-818.
871 10.1111/1365-2656.13662

872 Evans, O., Hick, P., Alford, B., Whittington, R.J., 2017a. Transmission of *Ostreid*
873 *herpesvirus-1* microvariant in seawater: Detection of viral DNA in seawater, filter
874 retentates, filter membranes and sentinel *Crassostrea gigas* spat in upwellers.
875 Aquaculture 473, 456-467. <https://doi.org/10.1016/j.aquaculture.2017.03.021>

876 Evans, O., Hick, P., Whittington, R.J., 2017b. Detection of *Ostreid herpesvirus-1*
877 microvariants in healthy *Crassostrea gigas* following disease events and their possible
878 role as reservoirs of infection. J Invert Pathol 148, 20-33.
879 <https://doi.org/10.1016/j.jip.2017.05.004>

880 Evans, O., Paul-Pont, I., Hick, P., Whittington, R.J., 2014. A simple centrifugation method for
881 improving the detection of *Ostreid herpesvirus-1* (OsHV-1) in natural seawater
882 samples with an assessment of the potential for particulate attachment. Journal of
883 Virological Methods 210(0), 59-66. <http://dx.doi.org/10.1016/j.jviromet.2014.09.023>

884 Evans, O., Paul-Pont, I., Whittington, R.J., 2017c. Detection of *Ostreid herpesvirus 1*
885 microvariant DNA in aquatic invertebrate species, sediment and other samples
886 collected from the Georges River estuary, New South Wales, Australia. Dis Aquat Org
887 122(3), 247-255. <https://doi.org/10.3354/dao03078>

888 Farley, C.A., Banfield, W.G., Kasnic, G., Jr., Foster, W.S., 1972. Oyster Herpes-Type Virus.
889 Science 178(4062), 759-760. <https://doi.org/10.2307/1735848>

890 Friedman, C., Estes, R., Stokes, N., Burge, C., Hargove, J., Barber, B., Elston, R., Burreson,
891 E., Reece, K., 2005. Herpes virus in juvenile Pacific oysters *Crassostrea gigas* from
892 Tomales Bay, California, coincides with summer mortality episodes. Dis Aquat Org
893 63(1), 33-41. 10.3354/dao063033

894 Fuhrmann, M., Castinel, A., Cheslett, D., Nozal, D.F., Whittington, R.J., 2019. The impacts
895 of *Ostreid herpesvirus 1* microvariants on Pacific oyster aquaculture in the Northern
896 and Southern Hemispheres since 2008. Revue Scientifique Et Technique-Office
897 International Des Epizooties 38(2), 491-509. <https://doi.org/10.20506/rst.38.2.3000>

898 Gittenberger, A., Voorbergen-Laarman, M.A., Engelsma, M.Y., 2016. *Ostreid herpesvirus*
899 OsHV-1 muVar in Pacific oysters *Crassostrea gigas* (Thunberg 1793) of the Wadden
900 Sea, a UNESCO world heritage site. J Fish Dis 39(1), 105-109.
901 <https://doi.org/10.1111/jfd.12332>

902 Hick, P., Evans, O., Looi, R., English, C., Whittington, R.J., 2016. Stability of *Ostreid*
903 *herpesvirus-1* (OsHV-1) and assessment of disinfection of seawater and oyster tissues
904 using a bioassay. Aquaculture 450, 412-421.
905 <https://doi.org/10.1016/j.aquaculture.2015.08.025>

906 Hick, P.M., Evans, O., Rubio, A., Dhand, N.K., Whittington, R.J., 2018. Both age and size
907 influence susceptibility of Pacific oysters (*Crassostrea gigas*) to disease caused by
908 *Ostreid herpesvirus-1* (OsHV-1) in replicated field and laboratory experiments.
909 Aquaculture 489, 110-120. <https://doi.org/10.1016/j.aquaculture.2018.02.013>

910 Hine, P.M., Thorne, T., 1997. Replication of herpes-like viruses in haemocytes of adult flat
911 oysters *Ostrea angasi*: An ultrastructural study. Disease of Aquatic Organisms 29(3),
912 189-196. [http://www.scopus.com/inward/record.url?eid=2-s2.0-
913 0030953661&partnerID=40&md5=14baec6c1cd641cbd52daa52c1cadf8d](http://www.scopus.com/inward/record.url?eid=2-s2.0-0030953661&partnerID=40&md5=14baec6c1cd641cbd52daa52c1cadf8d)

914 Hine, P.M., Wesney, B., Besant, P., 1998. Replication of a herpes-like virus in larvae of the
915 flat oyster *Tiostrea chilensis* at ambient temperatures. *Dis Aquat Org* 32(3), 161-171.
916 <https://doi.org/10.3354/dao032161>

917 Hwang, J.Y., Park, J.J., Yu, H.J., Hur, Y.B., Arzul, I., Couraleau, Y., Park, M.A., 2013.
918 *Ostreid herpesvirus 1* infection in farmed Pacific oyster larvae *Crassostrea gigas*
919 (Thunberg) in Korea. *J Fish Dis* 36(11), 969-972. <https://doi.org/10.1111/Jfd.12093>

920 Jenkins, C., Hick, P., Gabor, M., Spiers, Z., Fell, S.A., Gu, X., Read, A., Go, J., Dove, M.,
921 O'Connor, W., Kirkland, P.D., Frances, J., 2013. Identification and characterisation of
922 an *Ostreid herpesvirus-1* microvariant (OsHV-1 micro-var) in *Crassostrea gigas*
923 (Pacific oysters) in Australia. *Dis Aquat Org* 105(2), 109-126.
924 <https://doi.org/10.3354/dao02623>

925 Keeling, S., Brosnahan, C., Williams, R., Gias, E., Hannah, M., Bueno, R., McDonald, W.,
926 Johnston, C., 2014. New Zealand juvenile oyster mortality associated with *Ostreid*
927 *herpesvirus 1*—an opportunistic longitudinal study. *Dis Aquat Org* 109(3), 231-239.
928 <https://doi.org/10.3354/dao02735>

929 Kumar, S., Stecher, G., Li, M., Knyaz, C., Tamura, K., 2018. MEGA X: molecular
930 evolutionary genetics analysis across computing platforms. *Molecular biology and*
931 *evolution* 35(6), 1547. <https://doi.org/10.1093/molbev/msy096>

932 Martenot, C., Denechere, L., Hubert, P., Metayer, L., Oden, E., Trancart, S., Travaille, E.,
933 Houssin, M., 2015. Virulence of *Ostreid herpesvirus 1* mu Var in sea water at 16
934 degrees C and 25 degrees C. *Aquaculture* 439, 1-6.
935 <https://doi.org/10.1016/j.aquaculture.2015.01.012>

936 Martenot, C., Fourour, S., Oden, E., Jouaux, A., Travaillé, E., Malas, J.P., Houssin, M., 2012.
937 Detection of the OsHV-1 μ Var in the Pacific oyster *Crassostrea gigas* before 2008 in
938 France and description of two new microvariants of the *Ostreid Herpesvirus 1* (OsHV-
939 1). *Aquaculture* 338–341(0), 293-296.
940 <http://dx.doi.org/10.1016/j.aquaculture.2011.12.030>

941 Martenot, C., Oden, E., Travaillé, E., Malas, J.-P., Houssin, M., 2011. Detection of different
942 variants of *Ostreid Herpesvirus 1* in the Pacific oyster, *Crassostrea gigas* between
943 2008 and 2010. *Virus Res* 160(1–2), 25-31.
944 <http://dx.doi.org/10.1016/j.virusres.2011.04.012>

945 Martenot, C., Oden, E., Travaillé, E., Malas, J.P., Houssin, M., 2010. Comparison of two real-
946 time PCR methods for detection of *ostreid herpesvirus 1* in the Pacific oyster

947 *Crassostrea gigas*. Journal of Virological Methods 170(1-2), 86-89.
948 <https://doi.org/10.1016/j.jviromet.2010.09.003>

949 Martenot, C., Travaille, E., Lethuillier, O., Lelong, C., Houssin, M., 2013. Genome
950 exploration of six variants of the *Ostreid Herpesvirus 1* and characterization of large
951 deletion in OsHV-1muVar specimens. Virus Res 178(2), 462-470.
952 <https://doi.org/10.1016/j.virusres.2013.08.006>

953 Mineur, F., Provan, J., Arnott, G., 2015. Phylogeographical analyses of shellfish viruses:
954 inferring a geographical origin for ostreid herpesviruses OsHV-1
955 (*Malacoherpesviridae*). Marine Biology 162(1), 181-192.
956 <https://doi.org/10.1007/s00227-014-2566-8>

957 Morga, B., Jacquot, M., Pelletier, C., Chevignon, G., Degremont, L., Bietry, A., Pepin, J.F.,
958 Heurtebise, S., Escoubas, J.M., Bean, T.P., Rosani, U., Bai, C.M., Renault, T., Lamy,
959 J.B., 2021. Genomic Diversity of the *Ostreid Herpesvirus* Type 1 Across Time and
960 Location and Among Host Species. Frontiers in Microbiology 12.
961 <https://doi.org/10.3389/fmicb.2021.711377>

962 Mortensen, S., Strand, A., Bodvin, T., Alfjorden, A., Skar, C.K., Jelmert, A., Aspan, A.,
963 Saelemyr, L., Naustvoll, L.J., Albretsen, J., 2016. Summer mortalities and detection of
964 *Ostreid herpesvirus* microvariant in Pacific oyster *Crassostrea gigas* in Sweden and
965 Norway. Dis Aquat Org 117(3), 171-176. <https://doi.org/10.3354/dao02944>

966 Nicolas, J., Comps, M., Cochenec, N., 1992. Herpes-like virus infecting Pacific oyster
967 larvae, *C.gigas*. Bulletin of the European Association of Fish Pathologists 12(1), 11-
968 13. <http://eafp.squarespace.com/bulletin-archive/1992-volume-12/issue-1/>

969 Oden, E., Martenot, C., Berthaux, M., Travaille, E., Malas, J.P., Houssin, M., 2011.
970 Quantification of *ostreid herpesvirus 1* (OsHV-1) in *Crassostrea gigas* by real-time
971 PCR: Determination of a viral load threshold to prevent summer mortalities.
972 Aquaculture 317(1-4), 27-31. <https://doi.org/10.1016/j.aquaculture.2011.04.001>

973 Paul-Pont, I., Dhand, N.K., Whittington, R.J., 2013a. Influence of husbandry practices on
974 OsHV-1 associated mortality of Pacific oysters *Crassostrea gigas*. Aquaculture 412,
975 202-214. <https://doi.org/10.1016/j.aquaculture.2013.07.038>

976 Paul-Pont, I., Dhand, N.K., Whittington, R.J., 2013b. Spatial distribution of mortality in
977 Pacific oysters *Crassostrea gigas*: reflection on mechanisms of OsHV-1 transmission.
978 Dis Aquat Org 105(2), 127-138. <https://doi.org/10.3354/dao02615>

979 Paul-Pont, I., Evans, O., Dhand, N.K., Rubio, A., Coad, P., Whittington, R.J., 2014.
980 Descriptive epidemiology of mass mortality due to *Ostreid herpesvirus-1* (OsHV-1) in

981 commercially farmed Pacific oysters (*Crassostrea gigas*) in the Hawkesbury River
982 estuary, Australia. *Aquaculture* 422–423, 146-159.
983 <http://dx.doi.org/10.1016/j.aquaculture.2013.12.009>

984 Paul-Pont, I., Evans, O., Dhand, N.K., Whittington, R.J., 2015. Experimental infections of
985 Pacific oyster *Crassostrea gigas* using the Australian *Ostreid herpesvirus-1* (OsHV-1)
986 μ Var strain. *Dis Aquat Org* 113(2), 137-147. [http://www.int-](http://www.int-res.com/abstracts/dao/v113/n2/p137-147/)
987 [res.com/abstracts/dao/v113/n2/p137-147/](http://www.int-res.com/abstracts/dao/v113/n2/p137-147/)

988 Peck, K.M., Lauring, A.S., 2018. Complexities of viral mutation rates. *Journal of Virology*
989 92(14). <https://doi.org/10.1128/jvi.01031-17>

990 Peeler, E.J., Allan Reese, R., Cheslett, D.L., Geoghegan, F., Power, A., Thrush, M.A., 2012.
991 Investigation of mortality in Pacific oysters associated with *Ostreid herpesvirus-1*
992 μ Var in the Republic of Ireland in 2009. *Prev Vet Med* 105(1-2), 136-143.
993 <https://doi.org/10.1016/j.prevetmed.2012.02.001>

994 Pernet, F., Lague, K., Petton, B., 2021. Competition for food reduces disease susceptibility in
995 a marine invertebrate. *Ecosphere* 12(4). 10.1002/ecs2.3435

996 Petton, B., Destoumieux-Garzón, D., Pernet, F., Toulza, E., de Lorgeril, J., Degremont, L.,
997 Mitta, G., 2021. The Pacific oyster mortality syndrome, a polymicrobial and
998 multifactorial disease: state of knowledge and future directions. *Frontiers in*
999 *Immunology* 12. <https://doi.org/10.3389/fimmu.2021.630343>

1000 Petton, B., Pernet, F., Robert, R., Boudry, P., 2013. Temperature influence on pathogen
1001 transmission and subsequent mortalities in juvenile Pacific oysters *Crassostrea gigas*.
1002 *Aquaculture Environ Interact* 3(3), 257-273. Doi 10.3354/Aei00070

1003 Prado-Alvarez, M., Garcia-Fernandez, P., Faury, N., Azevedo, C., Morga, B., Gestal, C.,
1004 2021. First detection of OsHV-1 in the cephalopod *Octopus vulgaris*. Is the octopus a
1005 dead-end for OsHV-1? *J Invert Pathol* 183. <https://doi.org/10.1016/j.jip.2021.107553>

1006 Ren, W., Chen, H., Renault, T., Cai, Y., Bai, C., Wang, C., Huang, J., 2013. Complete
1007 genome sequence of acute viral necrosis virus associated with massive mortality
1008 outbreaks in the Chinese scallop, *Chlamys farreri*. *Virology* 450, 10-19.
1009 <https://doi.org/10.1186/1743-422x-10-110>

1010 Renault, T., Arzul, I., 2001. Herpes-like virus infections in hatchery-reared bivalve larvae in
1011 Europe: Specific viral DNA detection by PCR. *J Fish Dis* 24(3), 161-167.
1012 <https://doi.org/10.1046/j.1365-2761.2001.00282.x>

1013 Renault, T., Le Deuff, R.M., Chollet, B., Cochenne, N., Gerard, A., 2000. Concomitant
1014 herpes-like virus infections in hatchery-reared larvae and nursery-cultured spat

1015 *Crassostrea gigas* and *Ostrea edulis*. Dis Aquat Org 42(3), 173-183.
1016 <https://doi.org/10.3354/dao042173>

1017 Renault, T., Le Deuff, R.M., Cochenec, N., Maffart, P., 1994. Herpesviruses associated with
1018 mortalities among Pacific oyster, *Crassostrea gigas*, in France - comparative study.
1019 Revue de Medecine Veterinaire 145, 735-742.

1020 Renault, T., Lipart, C., Arzul, I., 2001. A herpes-like virus infecting *Crassostrea gigas* and
1021 *Ruditapes philippinarum* larvae in France. J Fish Dis 24(6), 369-376.
1022 <https://doi.org/10.1046/j.1365-2761.2001.00300.x>

1023 Renault, T., Moreau, P., Faury, N., Pepin, J.-F., Segarra, A., Webb, S., 2012. Analysis of
1024 clinical *Ostreid herpesvirus 1* (Malacoherpesviridae) specimens by sequencing
1025 amplified fragments from three Vvirus genome areas. Journal of Virology 86(10),
1026 5942-5947. <https://doi.org/10.1128/jvi.06534-11>

1027 Renault, T., Tchaleu, G., Faury, N., Moreau, P., Segarra, A., Barbosa-Solomieu, V., Lapegue,
1028 S., 2014. Genotyping of a microsatellite locus to differentiate clinical *Ostreid*
1029 *herpesvirus 1* specimens. Vet Res 45(3), 3. <https://doi.org/10.1186/1297-9716-45-3>

1030 Renzette, N., Gibson, L., Bhattacharjee, B., Fisher, D., Schleiss, M.R., Jensen, J.D., Kowalik,
1031 T.F., 2013. Rapid intrahost evolution of human cytomegalovirus is shaped by
1032 demography and positive selection. Plos Genetics 9(9).
1033 <https://doi.org/10.1371/journal.pgen.1003735>

1034 Roque, A., Carrasco, N., Andree, K.B., Lacuesta, B., Elandalousi, L., Gairin, I., Rodgers,
1035 C.J., Furones, M.D., 2012. First report of OsHV-1 microvar in Pacific oyster
1036 (*Crassostrea gigas*) cultured in Spain. Aquaculture 324, 303-306.
1037 <https://doi.org/10.1016/j.aquaculture.2011.10.018>

1038 Rosani, U., Venier, P., 2017. Oyster RNA-seq data support the development of
1039 *Malacoherpesviridae* genomics. Frontiers in Microbiology 8.
1040 <https://doi.org/10.3389/fmicb.2017.01515>

1041 Rozas, J., Ferrer-Mata, A., Sanchez-DelBarrio, J.C., Guirao-Rico, S., Librado, P., Ramos-
1042 Onsins, S.E., Sanchez-Gracia, A., 2017. DnaSP v6: DNA Sequence Polymorphism
1043 Analysis of Large Datasets. Mol. Biol. Evol. 34, 3299-3302.

1044 Samain, J.F., Degremont, L., Soletchnik, P., Haure, J., Bedier, E., Ropert, M., Moal, J.,
1045 Huvet, A., Bacca, H., Van Wormhoudt, A., Delaporte, M., Costil, K., Pouvreau, S.,
1046 Lambert, C., Boulo, V., Soudant, P., Nicolas, J.L., Le Roux, F., Renault, T., Gagnaire,
1047 B., Geret, F., Boutet, I., Burgeot, T., Boudry, P., 2007. Genetically based resistance to
1048 summer mortality in the Pacific oyster (*Crassostrea gigas*) and its relationship with

1049 physiological, immunological characteristics and infection processes. *Aquaculture*
1050 268(1-4), 227-243. <https://doi.org/10.1016/J.Aquaculture.2007.04.044>

1051 Segarra, A., Pépin, J.F., Arzul, I., Morga, B., Faury, N., Renault, T., 2010. Detection and
1052 description of a particular *Ostreid herpesvirus 1* genotype associated with massive
1053 mortality outbreaks of Pacific oysters, *Crassostrea gigas*, in France in 2008. *Virus Res*
1054 153(1), 92-99. <https://doi.org/10.1016/j.virusres.2010.07.011>

1055 Shimahara, Y., Kurita, J., Kiryu, I., Nishioka, T., Yuasa, K., Kawana, M., Kamaishi, T.,
1056 Oseko, N., 2012. Surveillance of Type 1 *Ostreid herpesvirus* (OsHV-1) variants in
1057 Japan. *Fish Pathol* 47(4), 129-136. <https://doi.org/10.3147/jsfp.47.129>

1058 Thiry, E., Meurens, F., Muylkens, B., McVoy, M., Gogev, S., Thiry, J., Vanderplasschen, A.,
1059 Epstein, A., Keil, G., Schynts, F., 2005. Recombination in alphaherpesviruses.
1060 *Reviews in Medical Virology* 15(2), 89-103. <https://doi.org/10.1002/rmv.451>

1061 Whittington, R., Dhand, N., Evans, O., Paul-Pont, I., 2015a. Further observations on the
1062 influence of husbandry practices on OsHV-1 μ Var mortality in Pacific oysters
1063 *Crassostrea gigas*: age, cultivation structures and growing height. *Aquaculture* 438,
1064 82-97. <https://doi.org/10.1016/j.aquaculture.2014.12.040>

1065 Whittington, R., Hick, P., Evans, O., Rubio, A., Alford, B., Dhand, N., Paul-Pont, I., 2015b.
1066 Protection of Pacific oyster (*Crassostrea gigas*) spat from mortality due to *Ostreid*
1067 *herpesvirus-1* OsHV-1 μ Var) using simple treatments of incoming seawater in land-
1068 based upwellers. *Aquaculture* 437, 10-20.
1069 <https://doi.org/10.1016/j.aquaculture.2014.11.016>

1070 Whittington, R.J., Liu, O., Hick, P.M., Dhand, N., Rubio, A., 2019. Long-term temporal and
1071 spatial patterns of *Ostreid herpesvirus 1* (OsHV-1) infection and mortality in sentinel
1072 Pacific oyster spat (*Crassostrea gigas*) inform farm management. *Aquaculture* 513,
1073 734395. <https://doi.org/10.1016/j.aquaculture.2019.734395>

1074 Whittington, R.J., Paul-Pont, I., Evans, O., Hick, P., Dhand, N.K., 2018. Counting the dead to
1075 determine the source and transmission of the marine herpesvirus OsHV-1 in
1076 *Crassostrea gigas*. *Vet Res* 49(1), 34. <https://doi.org/10.1186/s13567-018-0529-7>

1077 Xia, J., Bai, C., Wang, C., Song, X., Huang, J., 2015. Complete genome sequence of *Ostreid*
1078 *herpesvirus-1* associated with mortalities of *Scapharca broughtonii* broodstocks. *Virology*
1079 J 12. <https://doi.org/10.1186/s12985-015-0334-0>

1080

1081