



HAL
open science

The open ocean kinetic energy cascade is strongest in late winter and spring

René Schubert, Oscar Vergara, Jonathan Gula

► **To cite this version:**

René Schubert, Oscar Vergara, Jonathan Gula. The open ocean kinetic energy cascade is strongest in late winter and spring. *Communications Earth & Environment*, 2023, 4, pp.1-8. 10.1038/s43247-023-01111-x . hal-04360959

HAL Id: hal-04360959

<https://hal.univ-brest.fr/hal-04360959>

Submitted on 22 Dec 2023


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

The open ocean kinetic energy cascade is strongest in late winter and spring

René Schubert ¹✉, Oscar Vergara² & Jonathan Gula ^{1,3}

The oceanic kinetic energy cascade, the flux of kinetic energy between currents of different horizontal scales, shapes the structure of the global ocean circulation and the associated heat, salt, nutrient, and oxygen fluxes. Here, we show with a numerical ocean simulation that the surface geostrophic cascade can be estimated from satellite altimetry observations and present its regional distribution and seasonal cycle at scales of 40 to 150 km for large parts of the global ocean based on observations. The time-mean cascade is inverse (towards larger scales), strongest in large-scale current systems, and decreases with distance from these systems. In the open ocean, the inverse cascade is associated with a maximum in late winter at the smallest scales studied, which transitions to scales larger than 100 km within two to three months, consistent with the widespread absorption of mixed-layer eddies by mesoscale eddies in spring.

¹ Univ Brest, CNRS, IRD, Ifremer, Laboratoire d'Océanographie Physique et Spatiale (LOPS), IUEM, Plouzané, France. ² Collecte Localisation Satellites (CLS), Ramonville Saint-Agne, France. ³ Institut Universitaire de France (IUF), Paris, France. ✉email: rene.schubert@univ-brest.fr

Ocean motions can be decomposed into movements at different horizontal scales ranging from large-scales ($\mathcal{O}(1000 \text{ km})$) through mesoscales ($\mathcal{O}(100 \text{ km})$) and sub-mesoscales ($\mathcal{O}(100 \text{ m}) - \mathcal{O}(10 \text{ km})$) to microscales ($<\mathcal{O}(100 \text{ m})$). The flux of kinetic energy (KE) between ocean currents at different scales, the KE cascade, plays a key role in several aspects of global climate. First, in combination with the transformation of KE into available potential energy (and vice versa), the cascade mediates the balance between the oceans forcing by the atmosphere, primarily by wind at large scales, and the dissipation of KE into heat at molecular scales¹. Second, the KE cascade controls the scale-distribution of KE including the strength of the subtropical gyre², the strength and position of large-scale currents such as the Gulf Stream^{3,4}, the strength of inter-ocean exchanges such as Agulhas leakage⁵, and the strength and distribution of ocean mesoscale eddies^{6–8} and thus the associated transports of heat, salt, nutrients, and oxygen. Third, through ocean-atmosphere interactions, the ocean and atmosphere KE cascades determine their response times to external forcing⁹. This induces a modulation of the atmospheric circulation by the oceanic KE cascade. In addition to being a key component of the climate system, understanding the KE cascade is critical for the validation and development of climate models, particularly regarding the parameterization of the sub-grid-scale energetic fluxes and dissipation.

For the large mesoscales, the KE flux was computed from gridded sea-surface height data that were interpolated from along-track satellite altimetry measurements on a regular 0.25° and daily grid (AVISO) assuming geostrophy and an f-plane^{6,10–12}. The results showed an inverse cascade (from smaller to larger scales) at scales larger than about 75 km and a forward cascade (from larger to smaller scales) at smaller scales. While the forward cascade was later found to be an artifact of the filtering and interpolation technique onto the regular grid, the inverse cascade at larger scales was found to be robust to filtering^{11,12}. At scales of the maximum inverse flux (about 200 km), the cascade has been shown to be orders of magnitude smaller in the open ocean compared to regions of strong current systems¹³. This is not surprising, as the inverse cascade is stopped at smaller scales in the open ocean, for example due to the Rhines effect¹⁴. Substantial inverse fluxes occur at these scales only in the large-scale current systems, where very large mesoscale eddies form as a consequence of large-scale instabilities and interact with the large-scale currents. Regional submesoscale-permitting simulations and regional observations have shown that the inverse cascade extends into the submesoscales to scales of the mixed-layer Rossby radius of deformation (about 15 km)^{12,15–20}. Applying a filtering approach for computing the KE flux^{21–24} to submesoscale-permitting model data, it was found that the underlying process of the submesoscale inverse cascade is primarily the absorption of submesoscale mixed layer eddies by mesoscale eddies¹⁷. Consistently, the maximum of the submesoscale inverse flux has been found to occur immediately after the submesoscale season in late winter and to shift to larger scales in spring¹⁷. Based on observations, the existence of the submesoscale inverse KE cascade and its seasonality could only be demonstrated for small ocean regions^{16,18–20}. Indirect evidence that this submesoscale inverse cascade is active in large parts of the global ocean has recently been provided by high-resolution satellite products²⁵. Recent model studies have applied Helmholtz decomposition and principal strain coordinate transformation prior to the computation of the scale KE flux, and have shown that the inverse cascade is primarily driven by geostrophically balanced flows^{26,27}. This confirms that it should be possible to analyze the near-surface inverse KE cascade on the basis of sea-surface height.

In this study, we provide an observation-based estimate of the surface geostrophic KE cascade and its seasonal cycle at scales of

40–150 km in large parts of the global ocean. We use information on the dynamics in the respective scale-band from satellite along-track altimetry data measured by JASON-3 as well as from a submesoscale-permitting simulation of the Atlantic. The KE flux can not be computed directly from along-track sea-surface height. However, using the simulation, we prove that it is possible to estimate the KE flux from the simulated sea-surface height along the tracks of JASON-3 and apply the proposed estimation technique to the actual measurements.

Results

If the measured along-track absolute dynamic topography (η) is dominated by geostrophic flows, the across-track f-plane geostrophic flow component can be computed as $u = -\frac{g}{f}\eta_y$, where $g = 9.81 \text{ m s}^{-2}$ is the gravitational acceleration, $f = 2\Omega \sin(2\pi \frac{\theta}{360^\circ})$ is the Coriolis parameter with the Earth's angular speed $\Omega = \frac{2\pi}{86400} \text{ s}^{-1}$ and the latitude θ , and y is the along-track direction with increasing latitude. Partial derivatives of a with respect to b are written here and in the following as $\frac{\partial a}{\partial b} = a_b$. For this study, we use five years of tide-corrected and filtered measurements of η taken by the JASON-3 satellite along the tracks shown in Fig. 1a. From these, we estimate the surface geostrophic $5^\circ \times 5^\circ$ domain-averaged scale-KE flux through a specific scale L as

$$\langle \Pi_{est} \rangle = -C(L)\rho_0 \langle \overline{u^2} - \bar{u}^2 \rangle \langle |\bar{u}_y| \rangle, \quad (1)$$

where angle brackets indicate averages over $5^\circ \times 5^\circ$ subdomains for each measurement cycle, C is an estimation coefficient identified from a numerical ocean simulation, and $\rho_0 = 1024 \text{ kg m}^{-3}$ is the standard density. The overlines denote fields convoluted with a tophat kernel whose length is equal to that of the respective scale L and which is normalized so that it integrates to 1. The computation of C and the proof that the proposed estimation works is given in the “Methods” section.




Fig. 1 The time-mean inverse scale kinetic energy flux. The time-mean inverse scale kinetic energy flux $-\Pi_{est}$ from JASON-3 SSH at scales of 200 km (a), 140 km (b), and 60 km (c). In (a), thin black lines show the JASON-3 tracks. Hatches mark regions where the time-mean transition scale from balanced to unbalanced flows T is larger than the scale, $-\Pi_{est}$ is shown for, and thus, where the results have to be interpreted with care.




Fig. 2 The seasonal cycle of the inverse scale kinetic energy flux. Monthly climatological anomalies of the scale kinetic energy flux $-\Pi_{est}$ from JASON-3 with respect to the 2017–2021 mean. Red (blue) colors mark enhanced (reduced) inverse cascade. Shown are the relative anomalies for the FMA-mean (a) and ASO-mean (b) flux at 60 km, as well as the anomalies for the area-mean flux between 15°N and 65°N (c), and between 15°S and 65°S (d). Contour lines and diagonal hatches mark where the maximum transition scale from balanced to unbalanced flows T is in the respective season larger than 60 km and thus, where the results are potentially erroneous due to non-geostrophic gradients of η . Regions of strong large-scale currents are excluded before the area-averaging for the bottom panels. They are identified by the temporal standard deviations of 2017–2021 AVISO-SSH averaged over $5^\circ \times 5^\circ$ domains larger than 20 cm (dotted) and by anomalies of the KE flux that are larger than $15 \text{ mW km}^{-1} \text{ m}^{-3}$ at 60 km (horizontal hatches). The respective area-mean T (solid line) plus and minus one standard deviation (dashed lines) are shown in (c) and (d).

If η is not dominated by geostrophic flows, the computation of geostrophic currents from the measurements fails. A lower bound for the scales at which the respective gradients of η can be used to derive geostrophic flows is provided by the transition scale from balanced to unbalanced flows T . T is estimated from the JASON-3 along-track data with a previously published method²⁸ and is averaged over the same $5^\circ \times 5^\circ$ grid on which we estimate the scale KE flux. T is associated with lower values in regions with strong balanced flows (large-scale currents and surface eddy pathways) and higher values in regions with weak balanced flows (weak eddy activity)²⁸. Furthermore, T is associated with a seasonal cycle, with lower values in spring when there are more balanced submesoscale vortices, and higher values in summer and autumn when the surface signature of internal waves is amplified²⁹. In Figs. 1 and 2 we highlight with black lines the regions, scales, and seasons where the results are partially corrupted by too strong imprints of unbalanced flows into the absolute dynamic topography measurements. Furthermore, we exclude the tropics between 15°S and 15°N from our analysis, because the baroclinic mode-1 and mode-2 diurnal tides significantly affect the SSH-spectrum³⁰ there, and consequently T exceeds 75 km throughout the year³¹.

The time-mean scale KE flux. The time-mean estimated fluxes at scales of 60, 140, and 200 km are shown in Fig. 1. The largest (inverse) fluxes occur at all scales in regions of very strong near-surface current systems, such as the Gulf-Stream - North-Atlantic-Current system, the Agulhas system, the Brazil-Malvinas-Confluence or the Kuroshio. The further away from the large-scale currents, the smaller the fluxes. Fluxes at 140 km and 200 km scales are very close to each other and have substantial amplitudes almost only in regions of large-scale current systems. At 200 km, the pattern is very similar to the one presented in a previous study based on gridded AVISO data¹³. At this scale, almost no time-mean fluxes are found in the open ocean. At 60 km, fluxes of about $10\text{--}20 \text{ mW km}^{-1} \text{ m}^{-3}$ occur in large parts of the mid-latitudes. This is consistent with wide-spread mixed-layer instabilities leading

to the formation of submesoscale eddies, which are subsequently absorbed by the mesoscales¹⁷.

The seasonal cycle of the scale KE flux. The seasonal cycle of Π_{est} and its scale distribution are shown in Fig. 2. At 60 km, the anomalies of the FMA-mean (Fig. 2a) and the ASO-mean (Fig. 2b) relative to the 2017–2021 mean flux show enhanced inverse fluxes in spring and reduced inverse fluxes in autumn for most open ocean regions. This is consistent with a previously published map of the seasonal difference in geostrophic KE computed from along-track altimetry²⁵. In the open ocean, enhanced fluxes occur in winter and spring with a maximum in early spring (February in the Northern Hemisphere and July in the Southern Hemisphere) at 40 km scale (Fig. 2c, d). During the following three to four months, enhanced fluxes occur at increasingly larger scales. Deeper mixed-layer depths in winter allow for the accumulation of more available potential energy at mixed-layer fronts. Mixed-layer baroclinic instability of the fronts releases this potential energy into KE in the form of mixed-layer eddies²⁵. These eddies have a diameter on the order of the mixed-layer Rossby deformation radius (about 15 km) and are stronger and more frequent in winter. Subsequently, these eddies grow and are absorbed by mesoscale eddies. The absorption process takes about 2–3 months, resulting in a shift of the maximum scale KE flux from late winter at 30 km to late spring at 140 km. This climatological scale-time pattern of the observation-based estimated inverse flux is consistent with that of the power-spectral density of KE computed from AVISO³² and along-track altimetry²⁵. The same consistency was also found in a simulation of the open ocean near the Agulhas-system¹⁷. In regions of strong large-scale flows, in particular the Gulf Stream and Kuroshio extensions, the Brazil Malvinas Confluence, and the Agulhas region and Return Current, the estimated fluxes at 60 km mainly show an opposite seasonal cycle to that of the open ocean. This is consistent with results from a recent model study of the Gulf Stream, which found a stronger seasonal cycle of the balanced cascade outside of the Gulf Stream core, including a scale-shift of the maximum

inverse flux and a reduced seasonality without a scale-shift of the maximum in a region that partially includes the Gulf Stream²⁶. Furthermore, this is consistent with previously published results on along-track altimetry-based geostrophic KE levels, which revealed an opposite seasonal cycle within the strong current systems²⁵. We speculate that the reason for this phenomenon is that other energetic submesoscale and mesoscale processes overcome the seasonal cycle of mixed-layer instabilities and the subsequent inverse cascade of the resulting mixed-layer eddies. In the future, these other processes need to be identified.

Particularly in summer, in regions of weak mesoscale activity at scales of 70 km and smaller, the measured absolute dynamic topography is not dominated by balanced flows and $-\Pi_{est}$ gives spuriously enhanced values. This manifests itself in a spurious reduction or reversal of the seasonal cycle at 60 km in regions such as the North East Pacific or the Central Atlantic, where the maximum T of the respective season is larger than 60 km (contours and hatches in Fig. 2a, b), as well as small secondary maxima of $-\Pi_{est}$ in summer (August in the Northern Hemisphere and February in the Southern Hemisphere) mainly at scales smaller than 70 km (Fig. 2c, d).

Methods

JASON-3 satellite along-track altimetry data. JASON-3 measures the absolute dynamic topography η , the sum of the sea-level anomaly and the mean dynamic topography, between 66°S and 66°N with an along-track resolution of about 6 km at 1 Hz and a repeated measurement of each track every 10th day. Below scales of 25–35 km, JASON-class data are associated with noise^{33,34}. This noise is associated with a seasonal cycle that is similar, but not identical, to that of the submesoscale KE level³⁴ (at least north of 40°S³⁵): higher noise and submesoscale KE in winter and lower in summer. The sea-level anomaly is available in an original (unfiltered) and a low-pass “filtered” version. Fourier analysis shows that the seasonal cycle of power spectral density of the unfiltered η has deviations from 20 km on all scales larger than 30 km (Fig. 3e), indicating that the seasonal cycle at these scales is not corrupted by measurement noise. Furthermore, for this study, we use the tide-corrected data, where the barotropic tides and the coherent part of the baroclinic tides for modes M2, K1, O1, and S2 have been removed from the data. Note, however, that there is a residual effect of tides that cannot be corrected. Note also that η is the sea-surface height (SSH) corrected by the geoid. Since the geoid is associated with very large scales, this correction is not relevant for the analysis presented in this study. Therefore, η is also referred to as SSH in the following.

GIGATL1 simulation. The GIGATL1 simulation³⁶ used here has been performed with the Coastal and Regional Ocean COmmunity model (CROCO, 10.5281/zenodo.7415055), which is based on the Regional Ocean Modeling System (ROMS³⁷). The integration is performed on an Arakawa C-grid³⁸ using 100 vertical terrain-following sigma-levels where the bathymetry is taken from the SRTM30plus dataset. Vertical mixing is parameterized with the k-epsilon turbulence closure scheme with applied Canuto A stability function. No explicit lateral diffusivities and viscosities are used. The effect of bottom friction is parameterized with a logarithmic law using a roughness length of 0.01 m. The GIGATL1 grid is orthogonal based on an oblique Mercator projection and designed to have nearly uniform spacing in both horizontal dimensions. The grid spacing varies from 1 km at the central longitude of the grid to 735 m at the west and east extremes of the grid. A simulation with a grid-spacing three times as large as the one of GIGATL1, “GIGATL3”, has been initialized in January 2004. Initialization fields and boundary conditions are

from Simple Ocean Data Assimilation (SODA³⁹). GIGATL1 is initialized from the restarts of GIGATL3 in July 2007 and integrated twice, one experiment with simulated tides and one experiment without simulated tides. All simulations are forced by hourly atmospheric forcing data from the Climate Forecast System Reanalysis (CFSR) with a bulk formulation and a stress correction approach to parameterize the surface ocean current feedback to the atmosphere⁴⁰. Barotropic tidal forcing at the boundaries and tidal potential and self attraction from TPX07.2 and GOT99.2b are applied to GIGATL3 and GIGATL1 with tides. In this study, we analyze snapshots every fifth day over the period April 2008 to March 2009 for both simulations.

Model validation. The temporal standard deviation of SSH, computed from daily averages during Apr 2008–Mar 2009 averaged onto the AVISO grid, shows similar large-scale patterns for GIGATL1 without simulated tides and AVISO (Fig. 3a, b). The largest values occur in the region of the Gulf Stream, North Atlantic Current, and Brazil Malvinas Confluence systems, moderate values in the surrounding ocean, as well as in the Agulhas ring path, the Azores Current, and the North Equatorial Counter Current, and low values above the shelves and in the central subtropical gyres. The simulated SSH variabilities are smaller than in the observations in the Gulf Stream extension, the Brazil Current extension, and the Agulhas Ring Path, and larger north of 50°N. The standard deviation of SSH in the Northwest-Corner and the North Atlantic Current are in remarkably good agreement. The too low simulated SSH variability in the Agulhas ring path is a result of the monthly boundary conditions applied at the southern boundary of the domain, which are not frequent enough to resolve the full variability of the energetic eddies drifting out of the Agulhas region into the Atlantic.

Time-mean horizontal wavenumber spectra computed from SSH at JASON-3 tracks in a 20° × 20° domain in the open ocean show similar results for JASON-3 and GIGATL1 without simulated tides (Fig. 3c). For comparison, the GIGATL1 SSH is extracted at the measurement locations of JASON-3 from model snapshots every fifth day. The SSH-spectra from GIGATL1 without tides and the unfiltered, tide-corrected JASON-3 data show a very good agreement at scales larger than 60 km. At scales smaller than 60 km, the spectrum of the observations shows higher values compared to the simulation, which could be due to observational noise or unresolved processes. The spectrum of the filtered JASON-3 SSH follows that of the unfiltered down to 45 km, but with a shallower slope between 60 km and 45 km than in the simulation, which shows a k^{-4} slope down to 30 km.

Hovmöller plots of the monthly climatology of the SSH-spectrum for both the simulation and the observations (Fig. 3d and e) show maxima that shift from winter at scales of $\mathcal{O}(10\text{ km})$ to summer at scales of $\mathcal{O}(100\text{ km})$ similar to the maximum inverse cascade estimated from JASON-3 (Fig. 2c). This shift is consistent with previous results from AVISO and simulations^{17,32}, as well as with a maximum of submesoscale activity in winter and spring and a subsequent inverse cascade involving the growth of submesoscale eddies and their absorption by mesoscale eddies¹⁷. For the simulation, the Hovmöller plot is more patchy and shows the shift to larger scales one to two months later than in the observations. This can be due to the specific conditions in the single simulated year, which is compared here to a five-year average from the observations.

The scale KE flux. With the filtering approach, the scale KE flux can be computed as

$$\Pi = -\rho_0 \left[(\overline{u^2} - \bar{u}^2) \bar{u}_x + (\overline{uv} - \bar{u}\bar{v})(\bar{u}_y + \bar{v}_x) + (\overline{v^2} - \bar{v}^2) \bar{v}_y \right], \quad (2)$$




Fig. 3 Model validation. The standard deviation of SSH computed from Apr 2008–Mar 2009 daily mean fields of GIGATL1 without simulated tides (a) and AVISO (b). Thick black lines show the GIGATL1 domain, and the white box the region for which the Fourier transform is performed for the middle and bottom panels. c The time-mean SSH spectra and a gray straight line for the k^{-4} slope. d, e Hovmöller plots of the monthly climatological anomalies of the SSH spectra with respect to the mean over the period Apr 2008–Mar 2009 for (d) and 2017–2021 for (e). The respective area-mean T (solid line) plus and minus one standard deviation (dashed lines) are shown in (d) and (e).

where u and v are the total horizontal velocity components in perpendicular directions x and y ^{21–24}. The overlies denote fields convoluted with a two-dimensional top-hat kernel whose diameter is equal to that of the respective scale L and which is normalized by the respective enclosed area so that it integrates to 1. Π is only computed if the full kernel area is filled with data. To reduce the memory requirement for convolutions in the computation of Π , which increases exponentially with the respective scale, u and v are first averaged within 5×5 grid-cell boxes. This has almost no effect on the results, as the variations of the simulated values on the near-grid-scale are very small. Due to limits in computational time and storage, the computations were only feasible for snapshots every fifth day. While this does not

affect the results for area averages, the results for time averages may contain aliasing effects.

Large amplitudes of surface time-mean Π occur mainly in regions of strong current systems. While the Gulf Stream - North Atlantic Current and Brazil Malvinas Confluence systems show time-mean fluxes of both signs at 60 km scales, the equatorial current system is associated with a time-mean forward cascade (Fig. 4a). The area-mean flux for a large region in the North Atlantic, shows a dominant forward cascade in summer and autumn (JJASON) and a dominant inverse cascade in winter and spring (DJFMAM) (lower panel in Fig. 4a). The maximum inverse cascade shifts by a few months with increasing scale. This is consistent with the absorption of the winter-time submesoscale




Fig. 4 The simulated scale kinetic energy flux. The scale kinetic energy flux Π computed from GIGATL1 without simulated tides in the period Apr 2008–Mar 2009 from the total velocity (a), the geostrophic velocity (b), the geostrophic velocity and subsequently averaged over $5^\circ \times 5^\circ$ domains (c), and estimated with equation (1) from SSH cut from the simulation along JASON-3 tracks (d). The time-mean flux at 60 km scales is shown in (a) to (d). 15°S and 15°N are shown with black horizontal lines. Hatches mark regions that have been excluded for the computation of the estimation coefficient C . In (e) to (h), Hovmöller plots of the respective area-averaged flux in the North Atlantic are shown (domain marked with a black box in (a) to (d)). The scale of 60 km is marked with a black line.

vortices by the mesoscales which takes this time¹⁷, and with the seasonal cycle of the SSH spectrum in the North Atlantic Current (Fig. 3d).

The geostrophic scale KE flux can be computed by using (2), with the geostrophic velocity. Noisy large-amplitude time-mean geostrophic fluxes occur on the shelf, as well as in the tropics (Fig. 4b). Away from these regions, the time-mean geostrophic flux is inverse almost everywhere. When averaged over $5^\circ \times 5^\circ$ regions, the instantaneous geostrophic flux is only very rarely forward. In the time-mean, only a few regions near the coast or near the boundary of the domain show time-mean forward fluxes (Fig. 4c). Geostrophic fluxes averaged over the North Atlantic box show a dominant inverse cascade at all times and scales as well (bottom panels in Fig. 4b and c). The inverse cascade is maximum at the smallest investigated scales in February, March, and April, which shifts to April and May at scales of 100 km and larger. The similar timing and amplitude of the inverse cascade from the total velocity (Fig. 4a) support that balanced flows are primarily responsible for the inverse cascade, as shown in previous studies^{26,27}. Furthermore, this confirms that the near-surface winter and spring inverse cascade can be studied using SSH. However, for the interpretation of the geostrophic flux, one needs to be aware that the ageostrophic (net forward) cascade reduces the geostrophic (net inverse) cascade mainly at scales smaller than 40 km throughout the year (Fig. 4a). Note that the results presented here for the surface cascade from total velocities correspond to an extreme case, as forward fluxes are surface intensified and restricted to the very upper ocean, while the inverse cascade driven by the balanced flows extends deep into the ocean^{17,27}. The estimate of the geostrophic flux presented in the “Results” section is therefore more representative of the cascade over the depth-range of surface intensified balanced flows than of the cascade directly at the surface.

Estimating the scale KE flux from along-track data. In order to access the smallest scales observed by satellite altimetry, we must use the non-gridded along-track SSH product. However, it is not

possible to compute the geostrophic Π directly from along-track SSH, as one can only compute the across-track geostrophic velocity component. This implies that, of all the terms in (2), only the Leonard stress $\overline{u^2} - \overline{u}^2$, and the along-track derivative \overline{u}_y , are computable. For this study, we propose to estimate the flux with equation (1), which includes the product of these two terms giving the correct unit of a scale KE flux. Since $\overline{u^2} - \overline{u}^2$ is always positive, using the absolute value of \overline{u}_y , along with the minus sign means that the estimated geostrophic flux is globally inverse, consistent with what we found for the time- and area-averaged geostrophic fluxes in the simulations (Fig. 4c), with AVISO based observational studies at larger scales^{6,11}, and with theoretical studies that showed that the balanced flows are associated with inverse fluxes^{26,27}. However, in some shelf regions, $5^\circ \times 5^\circ$ averaged forward geostrophic fluxes do occur in our simulations, which implies that the estimated fluxes from JASON-3 need to be interpreted with care on the shelf. Trying to estimate the sign of the geostrophic flux directly, for example by using \overline{u}_y , without taking its absolute value and removing the minus sign in equation (1), has not been successful.

We average both subterms in equation (1) over $5^\circ \times 5^\circ$ boxes, as the estimation works better the larger this region is, and as they are large enough to cover reasonable amounts of along-track data (which have a track-to-track distance of several degrees for ascending and descending tracks (Fig. 1a)). This technique is associated with the inclusion of information from both horizontal directions, as the angle between ascending and descending tracks is large (Fig. 1a), and thus reduces the effect of the isotropy assumption that is necessarily made when one-dimensional data is cut from a two-dimensional field and then used to make a statement about that field.

The estimation coefficient C is identified from the simulations for each chosen scale by area- and time-averaging the term $\langle \Pi \rangle / (-\rho_0 \langle \tau_{uu} \rangle \langle |\overline{u}_y| \rangle)$, where $\langle \Pi \rangle$ is computed from the 5×5 grid-cell averaged SSH field and the divisor terms from the SSH extracted from the simulation along the JASON-3 tracks. For the final area-averaging of the term, regions near the coast and

boundary of the simulation, where the respective $5^\circ \times 5^\circ$ domain includes only a small ocean area are excluded (shown hatched in Fig. 4c), as well as regions and times where $L < T$. For scales of [30,40,50,60,70,80,90,100,120,140,160,180,200] km, C is computed to be [0.32,0.26,0.23,0.22,0.20,0.19,0.18,0.17,0.15,0.14,0.13,0.12,0.11] and thus shows an exponential decay with larger scales. Computing C from a parallel simulation with simulated tides, gives similar values (see Supplementary Note 1 and Supplementary Fig. 1).

Comparisons of the original and estimated time-mean fluxes at a scale of 60 km and their area-mean in the North Atlantic box show that the estimation reproduces the horizontal and temporal patterns and their amplitudes reasonably well (Fig. 4c, d). Large differences only show up in the Gulf Stream extension, where the flux is overestimated, and in a few shelf regions where the original flux is forward and thus cannot be estimated correctly with equation (1). The former may be due to the extreme non-isotropy of the core Gulf Stream extension and indicates that the fluxes are overestimated in similar energetic non-isotropic flows, such as the western Kuroshio or the Agulhas retroflection. Outside of these regions and small parts of the shelf, the estimation works very well. The pattern and amplitude of the estimated fluxes from GIGATL1 without simulated tides are very close to those estimated from JASON-3 data (Fig. 1). However, for the observations, higher fluxes are estimated in the Gulf Stream extension, the northern Brazil Malvinas Confluence, and the Agulhas ring path, which is consistent with higher mesoscale eddy activity in the observations (Fig. 3a, b). The largest differences between the area-averaged original and estimated geostrophic Π occur in April, where the maximum flux event is a bit overestimated (bottom panels Fig. 4c and d). Furthermore, the weak summer inverse cascade is slightly overestimated. This contributes to a slightly reduced amplitude of the seasonal cycle of the estimated flux compared to the original flux.

The horizontal scale of a SSH feature in the two-dimensional field may differ from that identified from the SSH along a track that cuts through the same feature. For example, if waves do not propagate in the direction of the track, they will appear to be associated with a larger wavelength along the track. Or if a track cuts through a circular eddy and does not cross its center, the eddy will appear to be smaller from the track information. The fact that the flux estimated from the one-dimensional SSH data agrees with that from the two-dimensional original shows that the estimation procedure corrects for this issue. We hypothesize that, on average, shorter scale features erroneously imprint at larger scales, and that the decreasing C with increasing scale corrects for this by reducing the pre-estimate (Eq. (1) without C) more at larger scales. Future research needs to address the question, why the estimation works.

Data availability

The data to reproduce the figures of the present paper can be accessed at <https://doi.org/10.17882/96675>. Furthermore, with the same link, the transition scale from balanced to unbalanced flows as well as the estimated scale kinetic energy flux can be accessed. The ocean model output used for this study is too large for an online upload. The full JASON-3 data used for this study is available at https://data.marine.copernicus.eu/product/SEALEVEL_GLO_PHY_L3_MY_008_062/services. The coastline shown in the Figures has been taken from www.naturalearthdata.com/downloads/110m-physical-vectors/.

Code availability

All the code to reproduce the study is available at <https://doi.org/10.17882/96675>. The scale kinetic energy flux computations based on the two-dimensional model data has thereby been performed with a modified version of the code published at <https://doi.org/10.5281/zenodo.4486265>.

Received: 26 May 2023; Accepted: 14 November 2023;

Published online: 30 November 2023

References

- Ferrari, R. & Wunsch, C. Ocean circulation kinetic energy: reservoirs, sources, and sinks. *Annu. Rev. Fluid Mechanics* **41**, 253–282 (2009).
- Lévy, M. et al. Modifications of gyre circulation by sub-mesoscale physics. *Ocean Modelling* **34**, 1–15 (2010).
- Hurlburt, H. E. & Hogan, P. J. Impact of $1/8^\circ$ to $1/64^\circ$ resolution on Gulf Stream model–data comparisons in basin-scale subtropical Atlantic Ocean models. *Dynamics Atmos. Oceans* **32**, 283–329 (2000).
- Chassignet, E. P. & Xu, X. Impact of horizontal resolution ($1/12^\circ$ to $1/50^\circ$) on gulf stream separation, penetration, and variability. *J. Phys. Oceanogr.* **47**, 1999–2021 (2017).
- Schubert, R., Gula, J. & Biastoch, A. Submesoscale flows impact Agulhas leakage in ocean simulations. *Commun. Earth Environ.* **2**, 1–8 (2021).
- Scott, R. B. & Wang, F. Direct evidence of an oceanic inverse kinetic energy cascade from satellite altimetry. *J. Phys. Oceanogr.* **35**, 1650–1666 (2005).
- Molemaker, M. J., McWilliams, J. C. & Dewar, W. K. Submesoscale instability and generation of mesoscale anticyclones near a separation of the California Undercurrent. *J. Phys. Oceanogr.* **45**, 613–629 (2015).
- Schubert, R., Schwarzkopf, F. U., Baschek, B. & Biastoch, A. Submesoscale impacts on mesoscale Agulhas dynamics. *J. Adv. Model. Earth Syst.* **11**, 2745–2767 (2019).
- Jansen, M. & Ferrari, R. Macroturbulent equilibration in a thermally forced primitive equation system. *J. Atmos. Sci.* **69**, 695–713 (2012).
- Tulloch, R., Marshall, J., Hill, C. & Smith, K. S. Scales, growth rates, and spectral fluxes of baroclinic instability in the ocean. *J. Phys. Oceanogr.* **41**, 1057–1076 (2011).
- Arbic, B. K., Polzin, K. L., Scott, R. B., Richman, J. G. & Shriver, J. F. On eddy viscosity, energy cascades, and the horizontal resolution of gridded satellite altimeter products. *J. Phys. Oceanogr.* **43**, 283–300 (2013).
- Qiu, B., Chen, S., Klein, P., Sasaki, H. & Sasai, Y. Seasonal mesoscale and submesoscale eddy variability along the North Pacific Subtropical Countercurrent. *J. Phys. Oceanogr.* **44**, 3079–3098 (2014).
- Wang, S., Liu, Z. & Pang, C. Geographical distribution and anisotropy of the inverse kinetic energy cascade, and its role in the eddy equilibrium processes. *J. Geophys. Res. Oceans* **120**, 4891–4906 (2015).
- Rhines, P. B. Waves and turbulence on a beta-plane. *J. Fluid Mechanics* **69**, 417–443 (1975).
- Capet, X., McWilliams, J. C., Molemaker, M. J. & Shchepetkin, A. Mesoscale to submesoscale transition in the California Current System. Part I: Flow structure, eddy flux, and observational tests. *J. Phys. Oceanogr.* **38**, 29–43 (2008).
- Soh, H. S. & Kim, S. Y. Diagnostic characteristics of submesoscale coastal surface currents. *J. Geophys. Res. Oceans* **123**, 1838–1859 (2018).
- Schubert, R., Gula, J., Greatbatch, R. J., Baschek, B. & Biastoch, A. The submesoscale kinetic energy cascade: Mesoscale absorption of submesoscale mixed layer eddies and frontal downscale fluxes. *J. Phys. Oceanogr.* **50**, 2573–2589 (2020).
- Balwada, D., Xie, J.-H., Marino, R. & Feraco, F. Direct observational evidence of an oceanic dual kinetic energy cascade and its seasonality. *Sci. Adv.* **8**, eabq2566 (2022).
- Garabato, A. C. N. et al. Kinetic energy transfers between mesoscale and submesoscale motions in the open ocean's upper layers. *J. Phys. Oceanogr.* **52**, 75–97 (2022).
- Qiu, B., Nakano, T., Chen, S. & Klein, P. Bi-Directional Energy Cascades in the Pacific Ocean from Equator to Subarctic Gyre. *Geophys. Res. Lett.* **49**, e2022GL097713 (2022).
- Leonard, A. Energy cascade in large-eddy simulations of turbulent fluid flows. *Adv. Geophys.* **18**, 237–248 (1975).
- Germano, M. Turbulence: the filtering approach. *J. Fluid Mechanics* **238**, 325–336 (1992).
- Eyink, G. L. Locality of turbulent cascades. *Phys. D: Nonlinear Phenomena* **207**, 91–116 (2005).
- Aluie, H., Hecht, M. & Vallis, G. K. Mapping the energy cascade in the North Atlantic Ocean: the coarse-graining approach. *J. Phys. Oceanogr.* **48**, 225–244 (2018).
- Steinberg, J. M., Cole, S. T., Drushka, K. & Abernathy, R. P. Seasonality of the mesoscale inverse cascade as inferred from global scale-dependent eddy energy observations. *J. Phys. Oceanogr.* **52**, 1677–1691 (2022).
- Contreras, M., Renault, L. & Marchesiello, P. Understanding energy pathways in the gulf stream. *J. Phys. Oceanogr.* **53**, 719–736 (2022).
- Srinivasan, K., Barkan, R. & McWilliams, J. C. A forward energy flux at submesoscales driven by frontogenesis. *J. Phys. Oceanogr.* **53**, 287–305 (2023).

28. Vergara, O., Morrow, R., Pujol, M.-I., Dibarboure, G. & Ubelmann, C. Global submesoscale diagnosis using along-track satellite altimetry. *Ocean Sci.* **19**, 363–379 (2023).
29. Lahaye, N., Gula, J. & Rouillet, G. Sea surface signature of internal tides. *Geophys. Res. Lett.* **46**, 3880–3890 (2019).
30. Lawrence, A. & Callies, J. Seasonality and spatial dependence of meso- and submesoscale ocean currents from along-track satellite altimetry. *J. Phys. Oceanogr.* **52**, 2069–2089 (2022).
31. Qiu, B. et al. Seasonality in transition scale from balanced to unbalanced motions in the world ocean. *J. Phys. Oceanogr.* **48**, 591–605 (2018).
32. Storer, B. A., Buzzicotti, M., Khatri, H., Griffies, S. M. & Aluie, H. Global energy spectrum of the general oceanic circulation. *Nat. Commun.* **13**, 1–9 (2022).
33. Xu, Y. & Fu, L.-L. The effects of altimeter instrument noise on the estimation of the wavenumber spectrum of sea surface height. *J. Phys. Oceanogr.* **42**, 2229–2233 (2012).
34. Dufau, C., Orszynowicz, M., Dibarboure, G., Morrow, R. & Le Traon, P.-Y. Mesoscale resolution capability of altimetry: present and future. *J. Geophys. Res. Oceans* **121**, 4910–4927 (2016).
35. Klein, P. et al. Ocean-scale interactions from space. *Earth Space Sci.* **6**, 795–817 (2019).
36. Gula, J., Theetten, S., Cambon, G. & Rouillet, G. Description of the GIGATL simulations <https://doi.org/10.5281/zenodo.4948523> (2021).
37. Shchepetkin, A. & McWilliams, J. C. The Regional Oceanic Modeling System (ROMS): a split-explicit, free-surface, topography-following-coordinate ocean model. *Oceanogr. Meteorol.* **9**, 347–404 (2005).
38. Arakawa, A. & Lamb, V. R. Computational design of the basic dynamical processes of the UCLA general circulation model. *General Circulation Models Atmos.* **17**, 173–265 (1977).
39. Carton, J. & Giese, B. A reanalysis of ocean climate using Simple Ocean Data Assimilation (SODA). *Mon. Weather Rev.* **136**, 2999–3017 (2008).
40. Renault, L., Masson, S., Arsouze, T., Madec, G. & McWilliams, J. C. Recipes for how to force oceanic model dynamics. *J. Adv. Model. Earth Syst.* **12**, e2019MS001715 (2020).

Acknowledgements

The authors gratefully acknowledge support from the French National Agency for Research (ANR) through the project DEEPER (ANR-19-CE01-0002-01). J.G. further gratefully acknowledges support from ISblue “Interdisciplinary graduate school for the blue planet” (ANR-17-EURE-0015) and O.V. from the French Space Agency (CNES) via the French SWOT TOSCA program. This work was performed using HPC/AI resources from GENCI-TGCC (Grant 2023-A0090112051) and from DATARMOR at Ifremer, Brest, France.

Author contributions

R.S. and J.G. designed the study. J.G. performed the numerical model simulations. O.V. computed the monthly climatology of the transition scale from balanced to unbalanced flows. R.S. developed and executed all other analysis, produced the figures and wrote the text. J.G. and O.V. contributed to the discussion of the results and the writing of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s43247-023-01111-x>.

Correspondence and requests for materials should be addressed to René Schubert.

Peer review information *Communications Earth & Environment* thanks the anonymous reviewers for their contribution to the peer review of this work. Primary Handling Editors: Rachael Rhodes, Heike Langenberg. Peer reviewer reports are available.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023