



HAL
open science

Orf (contagious ecthyma) transmitted by a cat

Victoire Laumondais, Muriel Delhomme, Laurent Misery

► **To cite this version:**

Victoire Laumondais, Muriel Delhomme, Laurent Misery. Orf (contagious ecthyma) transmitted by a cat. *JEADV Clinical Practice*, 2024, 10.1002/jvc2.481 . hal-04731271

HAL Id: hal-04731271

<https://hal.univ-brest.fr/hal-04731271v1>

Submitted on 10 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Orf (contagious ecthyma) transmitted by a cat

Victoire Laumondais¹ | Muriel Delhomme² | Laurent Misery¹ 

¹Service de dermatologie et vénéréologie CHU Brest, Brest, France

²Médecine de prévention et de santé au travail, secrétariat général pour l'administration de la zone de défense Sud-Est, Lyon, France

Correspondence

Laurent Misery, Service de dermatologie et vénéréologie CHU Brest 2, Ave Foch 29200 Brest, France.

Email: laurent.misery@chu-brest.fr

KEYWORDS

cat, contagious ecthyma, ecthyma contagiosum, orf, vector

Funding information

None

IMAGING DIAGNOSIS

A 65-year-old woman presented with a 6-cm umbilicated lesion on the forearm (Figure 1), which was evolving for a few days. Orf was clinically diagnosed. There was no viral sample. The patient had not been in contact with any goats or sheep. However, she reported having been in contact with a cat presenting with scabbed sores around the right ear (Figure 2) 4 weeks earlier. This cat was leaving in a house next to a farm with ewes and lambs, that could have been in contact with this cat. Lesions spontaneously disappeared in the cat and in our

patient in 4 weeks, with a very discrete scar. We concluded that orf was transmitted by this cat.

Orf is usually transmitted by sheep and goats.¹ We found only three cases of orf in cats² in the literature, all involving close contact of cats with sheep or goats. Only one case of cat-to-human transmission³ has been noted, linked to a cat scratch.

The main differential diagnosis would be cowpox/catpox'.^{4,5} In this disease, the human patients present with large painful papules which progresses to ulceration and necrosis with thick, hard, black crusts and flu-like symptoms, which was not the case in our patient.

V. Laumondais wrote the paper and performed the literature review, M. Delhomme provided the case and L. Misery made the diagnosis and supervised the work.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Author(s). JEADV Clinical Practice published by John Wiley & Sons Ltd on behalf of European Academy of Dermatology and Venereology.



FIGURE 1 Orf lesion on the forearm of the patient.




FIGURE 2 Orf lesion on the ear of the cat.

AUTHOR CONTRIBUTIONS

Victoire Laumondais and Laurent Misery have written the paper. Muriel Delhomme and Laurent Misery performed the diagnosis.

ACKNOWLEDGEMENTS

The authors have no funding to report.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

ETHICS STATEMENT

The patient has given written informed consent for participation in the study and the use of her deidentified, anonymized, aggregated data and her case details (including photographs) for publication.

ORCID

Laurent Misery  <http://orcid.org/0000-0001-8088-7059>

REFERENCES

1. Rossi L, Tiecco G, Venturini M, Castelli F, Quiros-Roldan E. Human Orf with Immune-mediated reactions: a systematic review. *Microorganisms*. 2023;11:1138.
2. Fairley RA, Whelam EM, Pesavento PA, Mercer AA. Recurrent localised cutaneous parapoxvirus infection in three cats. *NZ Vet J*. 2008;56:196–201.
3. Frandsen J, Enslow M, Bowen AR. Orf parapoxvirus infection from a cat scratch. *Dermatol Online J*. 2011;17:9.
4. Vestey JP, Yirrell DL, Aldridge RD. Cowpox/catpox infection. *Br J Dermatol*. 1991;124:74–8.
5. Tack DM, Reynolds MG. Zoonotic poxviruses associated with companion animals. *Animals*. 2011;1:377–95.

How to cite this article: Laumondais V, Delhomme M, Misery L. Orf (contagious ecthyma) transmitted by a cat. *JEADV Clin Pract*. 2024;1–2. <https://doi.org/10.1002/jvc2.481>