



HAL
open science

Analysis of the structural characteristics and spatial organization of macrobenthic fauna in Oualidia lagoon, Morocco

Fatima El Asri, Ahmed Errhif, Mohamed-Naoufal Tamsouri, Daniel Martin, Mohamed Maanan, Hakima Zidane

► **To cite this version:**

Fatima El Asri, Ahmed Errhif, Mohamed-Naoufal Tamsouri, Daniel Martin, Mohamed Maanan, et al.. Analysis of the structural characteristics and spatial organization of macrobenthic fauna in Oualidia lagoon, Morocco. *Applied Water Science*, 2022, 12 (5), pp.96. 10.1007/s13201-022-01629-z . hal-04767247

HAL Id: hal-04767247

<https://hal.univ-brest.fr/hal-04767247v1>

Submitted on 5 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Analysis of the structural characteristics and spatial organization of macrobenthic fauna in Oualidia lagoon, Morocco

Fatima El Asri^{1,5} · Ahmed Errhif¹ · Mohamed-Naoufal Tamsouri² · Daniel Martin³ · Mohamed Maanan⁴ · Hakima Zidane⁵

Received: 13 September 2021 / Accepted: 8 March 2022 / Published online: 4 April 2022
© The Author(s) 2022

Abstract

Morocco is a very rich country in terms of wetlands, including bays, estuaries, lagoons, lakes, and rivers, among others. However, many of them need to be well managed and better exploited, thus requiring baseline descriptive studies to assess faunal patterns and trends. Oualidia lagoon is not an exception. In fact, the studies conducted to date in the lagoon mainly focused on its oceanographic features and aquaculture capacities. The present paper presents the first study on the whole macrofaunal assemblages of Oualidia lagoon, having as main objectives analyze the composition and structure of the assemblages, as well as their relationships with the main environmental variables explaining their spatial distribution variability in the enclosed ecosystem of the Oualidia coastal lagoon. The benthic macroinvertebrates of Oualidia lagoon were studied in summer 2013 based on 43 stations spread over the whole lagoon by using a Van Veen grab. Two replicate samples were collected for macrobenthos and one additional replicate for sediment analyses at each station. Also, single water salinity and temperature measurements were recorded at each station. Fifty-one species from seven different taxa were recorded, among which mollusks (37%), polychaetes (25%) and crustaceans (25%) dominated in terms of species number. *Peringia ulvae*, *Abra alba*, *Sphaeroma serratum*, *Cerastoderma edule* and *Corophium* sp. accounted for 88% of total abundance. The benthos was structured along a downstream–upstream gradient, showing three assemblages, named according to the IndVal index: (1) *Diopatra* cf. *marocensis*, (2) *Peringia ulvae*, and (3) *Cerastoderma edule*, while the main factors driving their distribution were granulometry and salinity. The Biotic and Environmental analysis showed that the combination of granulometry and salinity was the major factors controlling the spatial distribution of the macrozoobenthos in the Oualidia Lagoon.

Keywords Benthic macrofauna · Biodiversity · Distribution · Community structure · Environmental factors · Oualidia lagoon · Morocco

✉ Fatima El Asri
fatimaelasri25@gmail.com

- ¹ Health and Environment Laboratory, Faculty of Sciences Ain Chock, University Hassan II, B.P. 5366 Maârif, 20100 Casablanca, Morocco
- ² Specialized Center in Zootechny and Marine Aquaculture Engineering, National Institute of Fisheries Research, BP. 31, 93200 M'diq, Morocco
- ³ Centre for Advanced Studies of Blanes (CEAB-CSIC), Accés a la Cala St. Francesc, 14 Blanes 17300, Girona, Spain
- ⁴ UMR 6554 LETG-Nantes, University of Nantes, BP 81227, 44312 Nantes, France
- ⁵ Prospection of Littoral Resources Laboratory, National Institute for Fisheries Research (INRH), Road Sidi Abderrahmane Club Equestre Ould Jmel, Casablanca, Morocco

Introduction

Coastal lagoons are highly productive ecosystems acting as feeding, nursery, reproduction, and protection spots for invertebrates, fishes, and birds, as well as rich fishery grounds (Newton et al. 2018). However, they are fragile ecosystems, highly susceptible to both natural and anthropogenic disturbances (Blanchet et al. 2008; Rossi et al. 2006; Affian et al. 2009).

Many interactions occurring in coastal lagoons are mediated through macrobenthic organisms which: (1) contribute to mineralize, promote, and mix the oxygen flux into the sediment, (2) recycle the organic matter and nutrients, (3) decompose detrital matter, (4) stabilize sediments, and (5) contribute to the trophic web by linking producers to higher consumers such as fishes, birds and epibenthic crustaceans (Pearson and

Rosenberg 1978; Nybakken 1993; Heilskov and Holmer 2001; Newton et al. 2018). Such a relevant role, together with their sensitivity to human disturbance, leads these organisms to be often employed as indicators of coastal ecosystems health, and coastal lagoons are not an exception (Dauer 1993; Borja et al. 2000; Salas et al. 2004). The relatively limited capacity of dispersion and sedentary habits of adult benthic organisms does not allow them to easily avoid unfavorable environmental conditions. Consequently, they are considered as sensitive indicators of both naturally and anthropogenically driven changes in coastal ecosystems (Salas et al. 2004; Borja et al. 2000). Moreover, their relatively long lifespan allows them to integrate water and sediment quality conditions through time, as well as to show responses to temporal and chronic disturbances. Particularly, both stress tolerant and sensitive benthic species may show changes in diversity, biomass and abundance, while their assemblages may show changes in trophic and functional structure (Pearson and Rosenberg 1978; Reiss and Kröncke 2005).

The above-mentioned relationships between benthic organisms and environmental conditions in coastal lagoons have been seldom analyzed along Moroccan coasts (El Asri et al. 2018, 2021). Nevertheless, the country has a rich network of lagoons, particularly in the Atlantic littoral where Oualidia lagoon is located. Oualidia is the most important wintering area for migratory birds in Morocco (El Hamoumi et al. 2003) and a Wetland of International Importance (under the RAMSAR Convention). It also provides a valuable environment for a rich variety of plants, fish, and other wildlife (El Hamoumi et al. 2003) and is currently considered a Natural Park. Moreover, the lagoon watershed supports intensive agricultural, cattle rearing, shell-fishing and tourism activities being thus an on-going economical pole (Maanan et al. 2017). Despite its intrinsic interest, most previous studies in Oualidia focused on the effects of pollution (e.g., Zourarah et al. 2007; Hassou et al. 2014; Maanan et al. 2014), while only one described its hydrographic functioning (Hilmi et al. 2005) and four analyzed specific macrobenthic taxa (Chbicheb 1996; El Asri et al. 2015, 2018, 2021).

The present paper presents the first study on the whole macrofaunal assemblages of Oualidia lagoon, having as main objectives: (1) to describe the species composition and abundance, (2) to analyze the assemblage structure and distribution, and (3) to identify the main environmental variables driving the assemblage structure and distribution.

Materials and methods

Study area


Oualidia lagoon, located in the Moroccan Atlantic Ocean coast (32°44'42" N, 9°02'50" W, Fig. 1), measures 7 km long and about 1 km wide. Characteristically, it shows numerous side channels connected to a meandering main channel with 2 m of average depth and 5 m of maximum depth during flood tides (Bidet and Carruesco 1982). Flood tides cover more than 75% (2.25 km²) of the lagoon surface, bringing salt water to its upstream reaches and into a saline marsh beyond its second dam (Maanan et al. 2014; Hilmi et al. 2017). Salinity ranges from 30 to 36‰ in the inner areas with very low mixing and very weak stratification of the water column (Makaoui et al. 2018). The rainfall reached c.a. 250 mm per year, and the prevailing winds are W–NW and E, almost parallel to the major lagoon axis (Zourarah et al. 2007).

Sampling and data analysis

Forty-three stations were sampled in July 2013 (Fig. 1) using a Van Veen grab (0.125 m² surface area). Two replicate samples were collected for macrobenthos and one additional replicate for sediment analyses at each station. Also, single water salinity and temperature measurements were recorded at each station with the help of a thermo-salinometer. The macrobenthos replicates were sieved in situ through a 1 mm pore-sized mesh, and the retained materials were transferred to plastic containers and fixed in a 10% formalin solution. The sediment replicates were transferred to plastic contained and kept cold until analyzed.

In the laboratory, the macrofauna was sorted under a binocular microscope, identified at the lowest taxonomic level possible, counted and classified into trophic groups (e.g., Fauchald and Jumars 1979; Hily and Bouteille 1999; Pranovi et al. 2000; Affi et al. 2008; Khedhri et al. 2015) as follows: (1) herbivores (H), algae-feeding organisms (e.g., some echinoids), (2) scavengers, feeding on carrions deposited on the bottom (essentially gastropods and decapods), (3) detritus feeders, feeding on particulate organic matter, essentially vegetable detritus (mainly amphipods and tanaids), (4) carnivores, predators like errant polychaetes or sea-anemones, (5) suspension feeders, feeding on food suspended in the water column (e.g., most bivalves and some polychaetes), (6) surface deposit feeders, feeding on organic particles settling on the sediment (most

Fig. 1 Map showing the location of the study area and the sampling sites in Oualidia lagoon



sedentary polychaetes), and (7) subsurface deposit feeders, burrowers that ingest the sediment from which they extract their food.

Sediment replicates were used to measure: (1) chlorophyll ‘a’ contents (mg/m^2 , Lorenzen method, Holm-Hansen et al. 1965), (2) percentages of the total organic matter (TOM, % of weight loss on ignition 4 h at $450\text{ }^\circ\text{C}$), and (3) grain size (μm , laser granulometer Malvern, Mastersizer).

Macrobenthic spatial distribution and biodiversity were described by the species richness (S), abundance (A), Shannon diversity (H' , as \log_2) (Shannon 1948) and evenness (J') (Pielou 1966), as estimated with the Paleontological Statistics program (PAST v2.14) (Hammer et al. 2001). The structure of the assemblages was analyzed by a Hierarchical Ascending Classification (HAC) on a density [transformed to $\log_{10}(x + 1)$ to limit the influence of the most dominant taxa] per station matrix using the Bray–Curtis distance and the Ward’s method. Each identified assemblage was classified according to the IndVal index (Duf rene and Legendre 1997). Visually observed patterns and correlations of species density with abiotic parameters were quantitatively explored with the BIOENV (Biotic and Environmental linking) routine, based on Euclidean distances (for the abiotic dataset) and Spearman’s rank correlations. All multivariate

analyses were executed on an R environment (www.r-project.org). The results were plotted on maps using ARCGIS 10.0 (Figs. 2, 3).

Results

Macrofaunal descriptors

A total of 51 species of macrobenthic invertebrates were identified (Table 1), of which mollusks were the most diverse group (19 species, 37%), followed by polychaetes and crustaceans (13, 25.5% each) and echinoderms (three, 5.9%). Cnidarians, insects and tunicates (one species each) constituted a 6.0%. Mollusks clearly dominate in abundance (77% of all specimens recorded), followed by far by crustaceans (15%), polychaetes (5%), and echinoderms, cnidarians, insects and tunicates (3% altogether).

Density ranged from 0 to 6336 individuals per m^2 (1274 ± 1469 ind. m^{-2} , mean \pm SD). The stations showing the highest densities were roughly situated in the center of the lagoon (Fig. 4a). The number of species per station ranged from 0 (station 7) to 18 (station 40), with 7.4 ± 3.7 species (mean \pm SD). The stations showing the highest




Fig. 2 Temperature (a) and salinity (b) trends in Oualidia lagoon




Fig. 3 Spatial distribution of organic matter (a), chlorophyll 'a' (b), and granulometry (c) in Oualidia lagoon

richness were roughly situated close to the inlet of the lagoon (Fig. 4b). Diversity ($0\text{--}3.8$ bits, 1.6 ± 0.7 bits, mean \pm SD) and evenness (0.9 , 0.6 ± 0.2 , mean \pm SD) showed both a heterogeneous distribution (Fig. 5a and b).

The “best combination” of abiotic parameters explaining the observed density patterns was obtained by combining granulometry and salinity (BIOENV, $R=0.49$) (Table 2).

Assemblage structure

Three groups of benthic organisms were identified in the lagoon (HAC, Figs. 6, 7). Group 1 included six stations from the outer part, close to the inlet. It was characterized by sediments with median-coarser sand with low organic matter content, which were subject to strong tidal currents. The total number of species was 43, mainly distributed among mollusks (37%), crustaceans (28%) and polychaetes (28%) (Fig. 8a). The average density was low (529 ± 475 ind. m^{-2}) comparing to other groups, and the dominant taxa were mollusks (58%) and crustaceans (25%) (Fig. 8b), while the

average evenness was 0.7 ± 0.2 (Table 3). The assemblage was dominated in density by both surface deposit and suspension feeders (76.1%) (Fig. 9a), while carnivores (22%), detritus feeders (20%), surface deposit feeders (18%) and suspension feeders (18%) dominate in species richness (Fig. 9b). The most dominant species were the gastropod *Peringia ulvae* (31.7%), the crustacean *Corophium* sp. (24%), followed by far by the bivalve *Abra alba* (18%).

Group 2 included 21 stations mainly located inwards, along the middle area of the lagoon, and characterized by having mixed silty-sand and sandy-silt sediments. The average density was 1712 ± 1644 ind. m^{-2} , and mollusks were largely the most dominant, with more than 90% of the total (Fig. 8b). There were 25 species, most of which belonging to crustaceans (11.44%) and mollusks (9.36%) (Fig. 8a) and the evenness was low (0.5 ± 0.2) (Table 3). The assemblage was dominated in density by surface deposit feeders (ca. 80% of the total density), while other trophic groups were present with very low densities (Fig. 9b). Surface deposit feeders also dominate in number of species (24%), but were

Table 1 List of macrofauna established in the Oualidia lagoon

Taxonomic groups	Family	Species	
Mollusca	Cardiidae	<i>Cerastoderma edule</i>	
	Donacidae	<i>Donax trunculus</i> Linnaeus, 1758	
	Semelidae		<i>Abra alba</i> (W. Wood, 1802)
			<i>Scrobicularia plana</i> (da Costa, 1778)
	Lucinidae		<i>Loripes orbiculatus</i> Poli, 1791
			<i>Lucinoma borealis</i> (Linnaeus, 1767)
	Mactridae		<i>Spisula solida</i> (Linnaeus, 1758)
			<i>Spisula</i> sp.
	Veneridae	<i>Ruditapes decussatus</i> (Linnaeus, 1758)	
	Mytilidae	<i>Mytilus galloprovincialis</i> Lamarck, 1819	
	Haminoeidae	<i>Haminoea cf. japonica</i> Pilsbry, 1895	
	Aplysiidae	<i>Aplysia punctata</i> (Cuvier, 1803)	
	Onchidiidae	<i>Onchidella celtica</i> (Cuvier, 1817)	
	Naticidae	<i>Cochlis vittata</i> (Gmelin, 1791)	
	Hydrobiidae	<i>Peringia ulvae</i> (Pennant, 1777)	
	Nassariidae		<i>Tritia pfeifferi</i> (Philippi, 1844)
			<i>Tritia reticulata</i> (Linnaeus, 1758)
	Trochidae		<i>Gibbula umbilicalis</i> (da Costa, 1778)
			<i>Cymbula safana</i> (Lamarck, 1819)
	Polychaeta	Nereididae	<i>Hediste diversicolor</i> (O.F. Müller, 1776)
Polynoidae		<i>Harmothoe</i> sp.	
Phyllodoceidae		<i>Phyllodoce</i> sp.	
Glyceridae			<i>Glycera cf. tridactyla</i> Schmarda, 1861
			<i>Glycera alba</i> (O.F. Müller, 1776)
Nephtyidae			<i>Nephtys kersivalensis</i> McIntosh, 1908
			<i>Nephtys hombergii</i> Savigny in Lamarck, 1818
Lumbrineridae		<i>Lumbrineris coccinea</i> (Renier, 1804)	
Onuphidae		<i>Diopatra cf. marocensis</i> Paxton, Fadlaoui & Lechapt, 1995	
Ampharetidae		<i>Alkmaria romijni</i> Horst, 1919	
Pectinariidae		<i>Lagis cf. koreni</i> Malmgren, 1866	
Capitellidae		<i>Capitella</i> sp.	
Sabellidae	<i>Panousea africana</i> Rullier & Amoureux, 1969		
Crustacea	Melitidae	<i>Melita palmata</i> (Montagu, 1804)	
	Corophiidae	<i>Corophium</i> sp.	
	Caprellidae	<i>Caprella liparotensis</i> Haller, 1879	
	Ampithoidae	<i>Ampithoe</i> sp.	
	Anthuridae	<i>Cyathura carinata</i> (Krøyer, 1847)	
	Sphaeromatidae	<i>Sphaeroma serratum</i> (Fabricius, 1787)	
	Idoteidae	<i>Idotea balthica</i> (Pallas, 1772)	
	Tanaididae	<i>Tanais dulongii</i> (Audouin, 1826)	
	Apseudidae	<i>Apseudes</i> sp.	
	Cumacea		
	Paguridae	<i>Pagurus bernhardus</i> (Linnaeus, 1758)	
	Portunidae	<i>Carcinus maenas</i> (Linnaeus, 1758)	
	Balanidae	<i>Balanus</i> sp.	
	Insecta	Chironomidae	<i>Chironomus</i> sp.
Echinodermata	Holothuriidae	<i>Holothuria polii</i> Delle Chiaje, 1824	
	Ophiuridae	<i>Ophiura</i> sp.	
Cnidaria	Parechinidae	<i>Paracentrotus lividus</i> (Lamarck, 1816)	
	Hormathiidae	<i>Calliactis parasitica</i> (Couch, 1842)	
Chordata	Ascidiacea		




Fig. 4 Spatial distribution of macrofauna abundance (a) and species richness (b) in Oualidia lagoon




Fig. 5 Spatial distribution of Shannon (a) and evenness (b) indexes in Oualidia lagoon

Table 2 Summary of results from Bio-Env

Number of variable	Correlation	Variables
1	0.390	Granulometry
2	0.493	Granulometry & salinity
3	0.436	Granulometry, salinity & organic matter
4	0.415	Granulometry, salinity, organic matter & chlorophyll 'a'
5	0.405	Granulometry, salinity, organic matter, chlorophyll 'a' & temperature

The high value is marked in bold

closely followed by detritus feeders (22%) and scavengers (18%) (Fig. 9a). The most dominant species were the gastropod *P. ulvae* (66%), followed by far by the bivalve *A. alba* (22%).

Group 3 included 16 stations mainly located in the most inner area of the lagoon and characterized by variably silty sediment with high organic matter content (in shallower bottoms) and muddy sediments (in deeper bottoms).




Fig. 6 Dendrogram showing the groups of stations derived from the cluster analysis based on macrofaunal abundances in Oualidia lagoon. I, group 1; II, group 2; III, group 3




Fig. 7 Spatial distribution of the macrofaunal communities in Oualidia lagoon

The total number of species was 25, which were distributed mainly among crustaceans, mollusks and polychaetes (Fig. 8a). The mean density was $1,643 \pm 1,641 \text{ ind.m}^{-2}$, with the mollusks being the most dominant (62%), followed by crustaceans (31%). All remaining groups just represented a $\leq 2\%$ (Fig. 8b). The evenness was 0.7 ± 0.1 (Table 3). The assemblage was dominated in density by surface deposit feeders (44.7%), followed by detritus feeders (26.3%) and suspension feeders (25.8%) and the other groups ($\leq 3\%$) (Fig. 9b). Conversely, the number of species detritus feeders, suspension feeders and surface deposit feeders is 24% each (Fig. 9a). The most dominant species were *A. alba* (24.2%),

Sphaeroma serratum (21.3%), *P. ulvae* (21%), *Cerastoderma edule* (16.9%) and *Corophium* sp. (4%).

Discussion

The present study represents a reliable baseline for future ecological research by providing a detail description of the observed spatial variability in summer conditions, as well as the relationships between the environmental factors, the biotic descriptors and the assemblage structure. The water temperature increased from outer to inner parts of the lagoon due to the influence of the cold ocean waters entering through the inlet (Fig. 2a). The highest salinities are observed mainly at the stations located downstream of the lagoon, directly subject to marine influences. The low values recorded upstream of the lagoon are due to the resurgences of fresh water which are responsible for the slight desalination of the lagoon water upstream. The low value observed at station 43 is explained by the dilution by fresh water from nearby water sources. Both granulometry (Fig. 3c) and organic matter (Fig. 3a) patterns mirrored that of temperature, revealing the relevant influence of the hydrodynamic functioning of the lagoon, with the inner area tending to be relatively calm, thus to having more fine sediments and organic matter (Hilmi et al. 2005; Maanan et al. 2014). Conversely, salinity decreased from outer to inner parts (Fig. 2b), where there were the main freshwater arrivals (Hilmi et al. 2005; Damsiri et al. 2014; Hassou et al. 2014). Moreover, also, a minimum of salinity downstream was detected (but in a single station), which was probably caused by the numerous underwater freshwater outbreaks located in this area of the lagoon (Rharbi et al. 2001; Hilmi et al. 2005).

This environmental framework characterizes Oualidia as a typical lagoon environment, where the benthic macrofauna is dominated by gastropods, bivalves, amphipods, isopods and polychaetes (Mistri 2002; Bazairi et al. 2003). However, the presence of cnidarians, tunicates and insects was also recorded. The former two occurred likely in connection with the seaweed coverage, while the later was only present in the most enclosed parts of the lagoon, where the salinity was low due to the numerous underwater freshwater outbreaks.

The biodiversity in Oualidia largely doubled the previous record in the area by Chbicheb (1996), 51 versus 24, and was also higher than those reported in nearby lagoons (Mergaoui et al. 2003; Marchini et al. 2004; Chaouti and Bayed 2005). This relatively higher number of species may be related to the combined influence of the open sea at the entrance, the underground freshwater seepage in the central area and the main freshwater inputs in the most enclosed area (Carruesco 1989; Hilmi et al. 2005; El Asri et al. 2015, 2018). This particular structure seemed to favor a mixed presence of marine,

Fig. 8 Faunal composition structure of the macrofauna community expressed in relative number of species (a) and individuals (b)





Table 3 List of the main species of each macrofauna community classified by descending order using the IndVal index values

Cluster	Species	IndVal
1	<i>Diopatra cf. marocensis</i>	196.8
	<i>Lumbrineris coccinea</i>	92.3
	<i>Loripes orbiculatus</i>	82.8
	<i>Tritia reticulata</i>	70.8
	<i>Ruditapes decussatus</i>	52.9
2	<i>Peringia ulvae</i>	190.5
	<i>Tritia pfeifferi</i>	129.6
	<i>Hediste diversicolor</i>	127.9
	<i>Abra alba</i>	127.7
	<i>Calliactis parasitica</i>	64.4
3	<i>Cerastoderma edule</i>	257.4
	<i>Melita palmata</i>	232.8
	<i>Sphaeroma serratum</i>	163.4
	<i>Cyathura carinata</i>	139.5
	<i>Abra alba</i>	132.9

brackish and freshwater species in a lagoon environment (Zabi and Leloeuf 1993).

Our results based on the whole macrofauna in summer agree with the previously reported structuring based on mollusk (El Asri et al. 2015) and polychaete assemblages in summer and winter in Oualidia (El Asri et al. 2018) when were studied independently and, to some extent, with those of the whole macrofauna in winter (El Asri et al. 2021). In all cases, there were three assemblages clearly organized along the outer–inner axis of the lagoon. As reported for other lagoons, this type of assemblage structuring resulted from combining the particular habitat preferences of all macrobenthic species (Bazairi et al. 2003; Lakhdar Idrissi et al. 2004; Chaouti and Bayed 2008; Rodrigues et al. 2011; Lefrere et al. 2015). However, lagoon environments generally show marked seasonal variability (e.g., Bertrán et al. 2016), and there are no reasons allowing us to suspect that Oualidia could be an exception. Therefore, to assess the possible seasonal differences in the macrofaunal assemblages of Oualidia, there is an on-going research attempting to compare the summer situation analyzed in the present work with the results of a previous study carried out in winter conditions (El Asri et al. 2021).

Fig. 9 Trophic structure of the macrofauna community expressed in relative number of species (a) and individuals (b). Surface deposit feeders (SDF), Subsurface deposit feeders (SSDF), Detritus feeders (Dt), Suspension feeders (S), Carnivores (C), Scavengers (N), Herbivores (H)



In Oualidia, the benthic assemblages conforming the three groups identified in the HAC were named according to the IndVal index (Table 3). The *Diopatra* cf. *marocensis* assemblage (Group 1) was mainly distributed in the outer part of the lagoon and mollusks, crustaceans and polychaetes were almost equally dominant. The species richness was the highest found in the lagoon, due to the presence of both lagoon and marine species, the later entering the lagoon environment thanks to the proximity to the inlet (Cherkaoui 2006; Koaudio et al. 2008). Such a proximity also influenced the assemblage density, with the low numbers being likely caused by the high hydrodynamics favoring sedimentary instability (El Asri et al. 2015). Nevertheless, the assemblage revealed to be well-balanced, as indicated by a relatively high evenness.

The *Peringia ulvae* assemblage (Group 2) was the largest in number of stations and occurred mainly in the middle area of the lagoon. The hydrodynamic regime was moderated and the environment showed a relatively high nutrient enrichment (Rharbi et al. 2001; Hilmi et al. 2005). These conditions favored some of the established species, which led this assemblage to show the maximum average density combines with a poor structuring (as indicated by the very low evenness). Likely, this also explained the extreme dominance (i.e., ca. 90% of the total density) of two mollusk species, *P. ulvae* that gave the name to the assemblage, and the bivalve *A. alba*.

The *Cerastoderma edule* assemblage (Group 3) was mainly distributed in the inner part of the lagoon (Fig. 6) and harbored the same number of species as the *P. ulvae* assemblage (i.e., 25). The average density was also relatively high, with ca. 90% being represented by mollusks and crustaceans. However, contrary to the *P. ulvae* assemblage, the structure was well balanced (as indicated by the relatively high evenness), likely in connection with the relatively high level of nutrient enrichment characterizing this zone (Rharbi et al. 2001).

It can be seen that there is a direct relationship between the nature of the sediment and the dominance of certain trophic groups. In fact, surface deposit feeders and subsurface deposit feeders are better represented on silted substrates than on sandy substrates, deposit feeders are favored by a low water current speed unlike suspensivores which require a stronger current (McLusky and McIntyre 1988). This further explains the very pronounced numerical presence of suspensivores at the level of downstream stations where the speed of the tidal currents is higher, these results were found in several coastal marine ecosystems (e.g., Dauvin 1988; Grall and Glémarec 1997; Bazairi et al. 2003).

The overall structure of the macrobenthic population of the lagoon of Oualidia is similar to that of the lagoon of Smir (Chaouti 2003), but also to that of the lagoon of Valle di Gorino in Italy, for example (Mistri et al. 2002). It is

characterized by a small number of abundant and constant species and by many rare species. This type of structure is found mainly in environments under environmental stress (Blondel 1979).

The dominant species (*Peringia ulvae*, *Abra alba*, *Sphaeroma serratum*, *Corophium* sp., *Cerastoderma edule* and *Hediste diversicolor*) are characteristic of the lagoon littoral environments and correspond to the paralic biocoenosis within the meaning of Guélorget and Perthuisot (1983). Macrofauna seems to be well represented in this type of environment. Quantitative data show that variations in overall density are mainly due to numerically dominant species.

The structure of the macrobenthic community of the Oualidia lagoon was approached through the calculation of several indices. The analysis of the structure of the populations of the Oualidia lagoon reveals a low specific diversity compensated by an increase in the density of species and by the widening of their spatial niche. According to Blondel (1979), this type of stand is found particularly in environments subject to restrictive conditions. Nevertheless, the values of the H' index of less than 3 bits / Ind, encountered in this ecosystem, are characteristic of paucispecific populations present in peclihaline lagoon environments such as the Prévost lagoon (Amanieu et al. 1977, 1978) and the Grau du Piémanson pond (Camargue) (Grimonet 1979).

The values of the structure indices showed a great spatial variation testifying to a great heterogeneity in terms of community structure. The maximum values of these indices are observed at the downstream part of the lagoon which is most influenced by the ocean. The marine influence manifests itself in species enrichment but also in the maintenance of a relatively stable environment allowing the establishment and organization of a more structured community. The central zone, on the other hand, has the lowest values of these structure indices. It is a transition zone toward the interior of the lagoon where the environment is relatively unstable. The macrobenthic community at this level is unstructured and strongly dominated by only two species (*Peringia ulvae* and *Abra alba*) which are described as surface deposits. The upstream zone has the characteristics of a confined zone where the number of species is reduced but with appreciable diversity and above all fairness. The reduction in the specific richness in this area is the consequence of a selection of species by this particular environment resulting in a stable and characteristic community of the area. Similar results have been obtained in other lagoons such as the Ria d'Aveiro lagoon in Portugal (Garnerot et al. 2004), the Sacca di Goro lagoon in northeastern Italy (Mistri et al. 2001), or even in Dypsjø fjord in Denmark (Garnerot et al. 2004). At the level of the Bizerte lagoon in Tunisia, Belkhdja-Mahjoud et al. (2007) distinguished two different zones, one well structured with high diversity and regularity and another less structured where these indices are low reflecting a pronounced

imbalance benthic populations. The work of Chaouti and Bayed (2005) showed weak structure indices over the whole of the Smir lagoon in Morocco, testifying to a weak structure of the benthic fauna throughout this lagoon.

The weak structuring of benthic communities observed in general in paralic environments is the direct cause of the importance of variations in abiotic factors such as salinity (Llanso et al. 2002; Teske and Wooldridge 2003; Giménez et al. 2005; Pasquaud 2006), temperature and hydrodynamics (Stora et al. 1995; Blanchet et al. 2005; Tlig-Zouari et al. 2008). Added to this is the sediment factor which plays an important role in the structuring of endogenous benthic communities (Gray 1981; Mannino and Montagna 1997; Teske and Wooldridje 2001; Ysebaert and Herman 2002). Other biotics such as the trophic environment can also intervene in the structural organization of benthic macrofauna (Solidoro et al. 2004).

The results suggest that the spatial patterns of the macrobenthic assemblages of Oualidia were mainly influenced by the salinity gradient and the sediment composition, with hydrodynamics also playing a relevant role. This agrees with previous results found in different lagoons and enclosed bays, such as the Epe Lagoon (Uwadiae 2013), the Guanabara Bay (Mendes et al. 2007), or the Paranagua Bay (Boehs et al. 2004). However, the environmental factors controlling species distribution in different paralic environments may vary. In the Smir lagoon (Chaouti and Bayed 2008), the vegetation and grain size were the major factors influencing the structure of species. In the Sacca di Goro (Mistri et al. 2001), the main control was attributed to dissolved oxygen, temperature and salinity. In the Monolimni lagoon (Kevrekidis 2004), the most influencing factors were temperature and depth. In the Ria de Aveiro (Rodrigues et al. 2011), the spatial distribution of the fauna was found to be mainly related to the hydrodynamic regime and the salinity gradient, while the sediment grain size was much less important. Conversely, in the Arcachon Bay (Do et al. 2011), the structure of benthic communities was largely influenced by the combination of sediment grain size distribution and seagrass presence. Concerning the vegetation found at the level of the lagoon, it is the halophytic phanerogams characteristic of brackish water "the eelgrass", which predominate the intertidal zone and at the level of the schorres of the lagoon of Oualidia, as well as the green algae "the seaweeds". ulves." The phytoplankton is abundant and represented by five large groups (Bennouna 1999); indeed, surely, the vegetation has an effect on the distribution of the benthic macrofauna; however, it has not been studied and its impact pending to be determined.

All these paralic environments differ considerably from one another in morphology and water depth, which certainly gave rise to differences in hydrodynamic regime, temperature, salinity, granulometry, vegetal coverage, etc. This may

certainly explain the reported changes in the environmental factors controlling species distribution, but also the differences in the composition and abundance of the established macrobenthos (Hernandez-Guevara et al. 2008). Moreover, the existing interactions between the macrobenthic fauna and other members of the ecosystem, including different components of the benthic compartment, but also all organisms inhabiting the lagoon (e.g., fish, birds, etc.) are certainly affecting the reported differences, particularly through predation, competition, etc. (Gambi et al. 1998; Nicolaidou et al. 2006; Schückel et al. 2015; Khedhri et al. 2017). Finally, one cannot discard the anthropogenic influences, which tend to be a major force in driving the functioning of most paralic environments all around the world (e.g., Vezzone et al. 2020; Fang et al. 2020) and, certainly, Oualidia is not an exception (Jayed et al. 2015; Lakhalki et al. 2017).

Conclusion

This study on the analysis of the structural characteristics and spatial organization of macrobenthic fauna in Oualidia lagoon has contributed to 54,407 macrozoobenthic organisms from 51 species in Oualidia, with the gastropod *Peringia ulvae*, the bivalves *Abra alba* and *Cerastoderma edule*, the isopod *Sphaeroma serratum* and the amphipod *Corophium* sp. being the most abundant species. Overall, the macrobenthos was structured in three assemblages distributed along an inner–outer axis, with the observed patterns in assemblage composition and structure being mainly driven by granulometry and salinity, with the hydrodynamics also playing a relevant role. This study must be considered as a baseline for future monitoring programs on the coastal management in the Oualidia lagoon. Therefore, further investigations are needed to see the whole picture of biodiversity status of macrofauna in the Oualidia lagoon, and to monitor the ecosystem health, it would be very interesting to study the seasonal variability of the benthic community in the lagoon and its linkage to the fluctuating physical–chemical parameters. To maintain high biodiversity, it is necessary to develop a specific management plan that will take into account the importance of the shallow water habitats in maintaining the ecological functions of the lagoon including that of nursery for juvenile fishes.

Acknowledgements We are deeply grateful to Mr Brahim Moutaki (INRH Oualidia) and Mr Kaddioui (PhD student) for their help during sampling and to the LETG (UMR 6554, University of Nantes) for kindly allowing us to carry out the granulometric analyses.

Funding FE, AE, MT, MM, HZ were partly funded by a research program on coastal resources of the LERL/DRH-Casablanca (National Fisheries Research Institute). DM received funding from the Consolidated Research Group on Marine Benthic Ecology, Generalitat

de Catalunya (<http://agaur.gencat.cat>), grant 2017SGR378, and the Agencia Estatal de Investigación (AEI) (<http://www.ciencia.gob.es/portal/site/MICINN/aei>), European Regional Development Funds (FEDER) (https://ec.europa.eu/regional_policy/en/funding/erdf/), grant CTM2017–88080 / PopCOMics. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data availability The data that support the findings of this study are available from the corresponding author upon reasonable request.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All applicable international, national, and/or institutional guidelines for animal testing, animal care and use of animals were followed by the authors.

Sampling and field studies All necessary permits for sampling and observational field studies have been obtained by the authors from the competent authorities and are mentioned in the acknowledgements, if applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Affian K, Robin M, Maanan M, Digbehi B, Djaoua EV, Kouamé F (2009) Heavy metal and polycyclic aromatic hydrocarbons in Ebrié lagoon sediments, Côte d'Ivoire. *J Environ Monit* 159:531–541. <https://doi.org/10.1007/s10661-008-0649-z>
- Afi A, Ayari R, Brahim M (2008) Trophic organization of the macrozoobenthic assemblages within coastal areas subjected to anthropogenic activities. *J Mar Biol Assoc UK* 88:663–674. <https://doi.org/10.1017/S0025315408001318>
- Amanieu M, Guelorget O, Michel P (1977) Richesse et diversité de la macrofaune benthique d'une lagune littorale méditerranéenne. *Vie Milieu* 27(85):109
- Amanieu M, Do Chi T, Galzin R, Lasserre G (1978) Structure et importance des populations de crabe vert *Carcinus mediterraneus* dans l'étang de Prévost (Languedoc, France). *Bull Ecol* 9:329–341. <https://doi.org/10.1080/19425120.2017.1310155>
- Bazairi H, Bayed A, Glémarec M, Hily C (2003) Spatial organisation of macrozoobenthic communities in response to environmental factors in a coastal lagoon of the NW African coast (Merja Zerga, Morocco). *Oceanol Acta* 26:457–471. [https://doi.org/10.1016/S0399-1784\(03\)00041-0](https://doi.org/10.1016/S0399-1784(03)00041-0)
- Belkhdja Mahjoub H, Rhomdane MS, Missaoui H (2007) Répartition spatiale de la faune malacologique de la lagune de Bizerte : analyses écologiques. *Rapports De La Commission Internationale Pour L'exploration De La Mer Méditerranée* 38:430
- Bennouna A (1999) Etude du phytoplancton nuisible et de son environnement dans la lagune de Oualidia et Sidi Moussa (Maroc). Université Chouiab Doukkali, El Jadida (Maroc), Thèse de Doctorat, p 153
- Bertrán C, Fierro P, Encalada E, Peña-Cortés F, Tapia J, Hauenstein E, Vargas-Chacoff L (2016) Macrozoobenthos of the coastal Budi Lagoon, southern Chile: changes associated with seasonal environmental variation. *Braz J Oceanogr* 64(3):239–248. <https://doi.org/10.1590/S1679-87592016113306403>
- Bidet JC, Carruesco C (1982) Étude sédimentologique de la lagune Oualidia (Maroc). *Oceanol Acta*. pp 29–37
- Blanchet H, De Montaudouin X, Chardy P, Bachelet G (2005) Structuring factors and recent changes in subtidal macrozoobenthic communities of a coastal lagoon, Arcachon Bay (France). *Estuar Coast Shelf Sci* 64(4):561–576. <https://doi.org/10.1016/j.ecss.2005.03.016>
- Blanchet H, Lavesque N, Ruellet T, Dauvin JC, Sauriau PG, Desroy N, Desclaux C, Leconte M, Bachelet G, Janson A-L, Bessineton C, Duhamel S, Jourde J, Mayot S, Simon S, De MX (2008) Use of biotic indices in semi-enclosed coastal ecosystems and transitional waters habitats - Implications for the implementation of the European Water Framework Directive. *Ecol Indic* 8:360–372. <https://doi.org/10.1016/j.ecolind.2007.04.003>
- Blondel J (1979) *Biogéographie et écologie*. Masson ed., Paris. p 221
- Boehs G, Absher T, Da Cruz-Kaled A (2004) Composition and distribution of benthic molluscs on intertidal flats of Paranaguá Bay (Paraná, Brazil). *Sci Mar* 68(4):537–543. <https://doi.org/10.3989/scimar.2004.68n4537>
- Borja A, Franco J, Perez V (2000) A marine biotic index to establish the ecological quality of soft bottom benthos within European estuarine and coastal environments. *Mar Pollut Bull* 40:1100–1114. [https://doi.org/10.1016/S0025-326X\(00\)00061-8](https://doi.org/10.1016/S0025-326X(00)00061-8)
- Carruesco C (1989) Genèse et évolution de trois lagunes du littoral atlantique depuis l'holocène : Oualidia, Moulay Bou Salham (Maroc) et Arcachon (France). Université de Bordeaux I, Bordeaux (France), Thèse de Doctorat, p 485
- Chaouti A (2003) Biotypologie du macrozoobenthos de la lagune méditerranéenne de Smir: analyses spatiale et spatio-temporelle. Univ Mohammed V, Rabat, Thèse Doctorat, p 302
- Chaouti A, Bayed A (2005) Diversité taxonomique et structure de la macrofaune benthique des substrats meubles de la lagune de Smir. *Trav L'institut Sci Rabat Sér Gén* 4:33–42
- Chaouti A, Bayed A (2008) Spatial patterns of soft-bottom macroinvertebrates and relationships with environmental conditions in a north African coastal lagoon (Smir lagoon, Morocco). *Vie Milieu* 58:25–35
- Chbicheb A (1996) Organisation biogéologique d'un bassin paralique : la lagune d'Oualidia (côte atlantique marocaine). Université de Nantes, Nantes (France), Thèse de Doctorat, p 167
- Cherkaoui E (2006) Structure et organisation des peuplements macrozoobenthiques de l'estuaire du Bou Regreg après la construction du barrage. Université Mohammed V, Faculté des Sciences, Rabat (Maroc), Thèse de Doctorat, p 258p
- Damsiri Z, Natij L, Khalil K, Loudiki M, Rabouille C, Ettahiri O (2014) Spatio-temporal nutrients variability in the Oualidia lagoon (Atlantic Moroccan coast). *Int J Adv Res* 2(8):609–618
- Dauer DM (1993) Biological criteria, environmental health and estuarine macrozoobenthic community structure. *Mar Pollut Bull* 26:249–257. [https://doi.org/10.1016/0025-326X\(93\)90063-P](https://doi.org/10.1016/0025-326X(93)90063-P)
- Dauvin J-C (1988) Structure et organisation trophique du peuplement des sables grossiers à *Amphioxus lanceolatus* – Venus fasciata de la baie de Morlaix (Manche occidentale). *Cah Biol Mar* 29:163–185
- Do VT, de Montaudouin X, Lavesque N, Hugues B, Guyard H (2011) Seagrass colonization: knock-on effects on zoobenthic

- community, populations and individual health. *Estuar Coast Shelf Sci* 95:458–469. <https://doi.org/10.1016/j.ecss.2011.10.022>
- Dufrène M, Legendre P (1997) Species assemblages and indicator species : the need for a flexible asymmetrical approach. *Ecol Monogr* 67:345–366
- El Asri F, Zidane H, Maanan M, Tamsouri M, Errhif A (2015) Taxonomic diversity and structure of the molluscan fauna in Oualidia lagoon (Moroccan Atlantic coast). *Environ Monit Assess* 187(8):545. <https://doi.org/10.1007/s10661-015-4752-7>
- El Asri F, Zidane H, Errhif A, Tamsouri MN, Maanan M, Idrissi MM, Martin D (2018) Polychaete diversity and assemblage structure in the Oualidia Lagoon, Moroccan Atlantic coast. *J Mar Biol Assoc UK* 98(6):1337–1346. <https://doi.org/10.1017/S0025315417000388>
- El Asri F, Errhif A, Martin D, Tamsouri MN, Maanan M, Zidane H (2021) Community structure and spatial patterns of soft-bottom macrozoobenthos in Oualidia lagoon, Moroccan Atlantic. *Thalassas Int J Mar Sci* 37:119–129. <https://doi.org/10.1007/s41208-020-00263-3>
- Fang X, Li X, Xiang Y, Hao C, Zhao Y, Zhang Y (2020) Cumulative impact of anthropogenic nutrient inputs on lagoon ecosystems—A case study of Xincun Lagoon, Hainan. *China Reg Stud Mar Sci* 35:101213. <https://doi.org/10.1016/j.risma.2020.101213>
- Fauchald K, Jumars P-A (1979) The diet of worms: a study of polychaete feeding guilds. *Oceanogr Mar Biol Ann Rev* 17:193–284. <https://doi.org/10.1146/annurev-marine-010814-020007>
- Gambi MC, Conti G, Bremec CS, Zoologica S, Dohrn A (1998) Polychaete distribution, diversity and seasonality related to seagrass cover in shallow soft bottoms of the Tyrrhenian Sea (Italy). *Sci Mar* 62:1–17. <https://doi.org/10.3989/scimar.1998.62n1-21>
- Garnerot F, Bouchereau J-L, Rebelo JE, Guelorget O (2004) L'ichtyofaune dans l'organisation biologique d'un système paralique de type lagunaire, La Ria d'Aveiro (Portugal), en 1987–1988 et 1999–2000. *Cybiurn* 28:63–75
- Giménez L, Borthagaray AI, Rodríguez M, Brazeiro A, Dimitriadis C (2005) Scale-dependent patterns of macrofaunal distribution in soft-bottom sediment intertidal habitats along a large-scale estuarine gradient. *Helgol Mar Res* 59:224–236. <https://doi.org/10.1007/s10152-005-0223-9>
- Grall J, Glémarec M (1997) Using biotic indices to estimate macrobenthic community perturbations in the Bay of Brest. *Est Coast Shelf Sci* 44:43–53. [https://doi.org/10.1016/S0272-7714\(97\)80006-6](https://doi.org/10.1016/S0272-7714(97)80006-6)
- Gray JS (1981) *The ecology of marine sediments*. Cambridge University Press
- Grimonet J (1979) Introduction à l'étude démographique de la macrofaune benthique de l'étang du Grau de Piémanson (Camargue). *Mém. Ingén., Laboratoire d'Hydrobiologie Université des sciences et technique du Languedoc, Montpellier* : p 101
- Guélorget O, Penhaisot J-P (1983) Le domaine paralique: Expressions géologiques biologiques et économiques du confinement. *Trav Lab Geol ENS* 16:136
- Hammer Ø, Harper D, Ryan P (2001) PAST: Paleontological statistics software package for education and data analysis. *Palaeontol Electron* 4:1–9
- El Hamoumi R, Dakki M, Rguibi Idrissi H, Radi M (2003) Fiche descriptive sur les zones humides Ramsar (FDR) : Complexe de Sidi Moussa-Walidia (Maroc)
- Hassou N, Maanan M, Hennani M, Zourarah B, Assobhei O (2014) Original research article spatial and temporal variation of faecal pollution indicators (*Escherichia coli* and faecal streptococci) and physico-chemical parameters at the Oualidia lagoon and its watershed (Morocco). *Int J Curr Microbiol Appl Sci* 3:675–694
- Heilskov AC, Holmer M (2001) Effects of benthic fauna on organic matter mineralization in fish-farm sediments: Importance of size and abundance. *ICES J Mar Sci* 58:427–434. <https://doi.org/10.1006/jmsc.2000.1026>
- Hernandez-Guevara NA, Pech D, Ardisson P-L (2008) Temporal trends in benthic macrofauna composition in response to seasonal variation in a tropical coastal lagoon, Celestun. *Gulf of Mexico Mar Freshwater Res* 59(9):772–779
- Hilmi K, Koutitonsky V, Orbi A, Lakhdar J, Chagdali M (2005) Oualidia lagoon, Morocco: an estuary without a river. *African J Aquat Sci* 30:1–10. <https://doi.org/10.2989/16085910509503828>
- Hilmi K, Orbi A, Makaoui A, Bouksim H, Idrissi M, Ettahiri O, Benyounes A, Ait Chattou M (2017) Circulation marine de la baie de Dakhla (Sud du Maroc) par modèle hydrodynamique 2D. *Eur Sci J* 13:69–82
- Hily C, Bouteille M (1999) Modifications of the specific diversity and feeding guilds in an intertidal sediment colonized by an eelgrass meadow (*Zostera marina*) (Brittany, France). *C r Acad Sci Paris* 322:1121–1131
- Holm-Hansen O, Lorenzen CJ, Holmes RW, Strickland JDH (1965) Fluorometric determination of chlorophyll. *ICES J Mar Sci* 30:3–15. <https://doi.org/10.1093/icesjms/30.1.3>
- Jayed M, Benbrahim S, Bakkas S, Ramdani M, Flower R (2015) Accumulation of organochlorines in the European clam (*Ruditapes decussatus*) and sediment of the Oualidia lagoon (Morocco). *Bull Environ Contam Toxicol* 94:614–621. <https://doi.org/10.1007/s00128-015-1517-5>
- Kevrekidis T (2004) Seasonal variation of the macrozoobenthic community structure at low salinities in a Mediterranean lagoon (Monolimni Lagoon, Northern Aegean). *Int Rev Hydrobiol* 89(4):07–425. <https://doi.org/10.1002/iroh.200310703>
- Khedhri I, Djabou H, Afli A (2015) Trophic and functional organization of the benthic macrofauna in the lagoon of Boughrara – Tunisia (SW Mediterranean Sea). *J Mar Biol Assoc* 4:647–659. <https://doi.org/10.1017/S0025315414001544>
- Khedhri I, Afli A, Aleya L (2017) Structuring factors of the spatio-temporal variability of macrozoobenthos assemblages in a southern Mediterranean lagoon: How useful for bioindication is a multi-biotic indices approach? *Mar Pollut Bull* 114:515–527. <https://doi.org/10.1016/j.marpolbul.2016.10.023>
- Koaudio KN, Diomandé D, Ouattara A, Koné YJM, Gourène G (2008) Taxonomic diversity and structure of benthic macroinvertebrates in Aby Lagoon (Ivory Coast, West Africa). *Pak J Biol Sci* 11:2224–2230. <https://doi.org/10.3923/pjbs.2008.2224.2230>
- Lakhdar Idrissi J, Orbi A, Zidane F, Hilmi K, Sarf F, Massik Z, Makaoui A (2004) Organisation et fonctionnement d'un écosystème côtier du Maroc: la lagune de Khnifiss. *Rev Sci Eau* 174:447–462. <https://doi.org/10.7202/705542ar>
- Lakhlalki H, Jayed M, Benbrahim S, Rharbi N, Bouthir F, Ali Benhra A, Brahim Moutaki B (2017) Accumulation of organochlorine pesticides and polychlorinated biphenyls in the sediments of Oualidia lagoon (Morocco). *Bull Soc Zool Fr* 142(3):101–118
- Lefrere L, Ouassas M, Guillois B, Gillet P, Moukrim A (2015) Macro-benthic community structure of soft-bottom sediments in the Khnifiss lagoon, South of Morocco. *J Mater Environ Sci* 6:3226–3235
- Llanso RJ, Scott LC, Dauer DM, Hyland JL, Russell DE (2002) An estuarine benthic index of biotic integrity for the mid- Atlantic region of the United States. I. Classification of assemblages and habitat definition. *Estuaries* 25:1219–1230. <https://doi.org/10.1007/BF02692219>
- Maanan M, Ruiz-Fernández AC, Maanan M, Fattal P, Zourarah B, Sahabi M (2014) A long-term record of land use change impacts on sediments in Oualidia lagoon, Morocco. *Int J Sediment Res* 29:1–10. [https://doi.org/10.1016/S1001-6279\(14\)60017-2](https://doi.org/10.1016/S1001-6279(14)60017-2)
- Maanan M, El Barjy M, Hassou N, Zidane H, Zourarah B, Maanan M (2017) Origin and potential ecological risk assessment of trace elements in the watershed topsoil and coastal sediment of the Oualidia lagoon. *Morocco Hum Ecol Risk Assess* 24(3):602–614. <https://doi.org/10.1080/10807039.2017.1394176>

- Makaoui A, Idrissi M, Agouzouk A, Larissi J, Baibai T, Ouehabi ZE, Laamel MA, Bessa I, Ettahiri O, Hilmi K (2018) Oceanographic state of the Oualidia Lagoon, Morocco (2011–2012). *Eur Sci J* 14:93
- Mannino A, Montagna PA (1997) Small-scale spatial variation in macrobenthic community structure. *Estuaries* 20:159–173
- Marchini A, Gauzer K, Occhipinti-Ambrogi A (2004) Spatial and temporal variability of hard-bottom macrofauna in a disturbed coastal lagoon (Sacca di Goro, Po River Delta, Northwestern Adriatic Sea). *Mar Poll Bull* 48:1084–1095. <https://doi.org/10.1016/j.marpolbul.2003.12.015>
- Mcluskay DS, McIntyre AD (1988) Characteristics of the fauna. In: Postma H, Zijlstra JJ (eds) *Continental shelves ecosystems of the world*. Elsevier
- Mendes CLT, Tavares M, Soares-Gomes A (2007) Taxonomic sufficiency for soft-bottom sublittoral mollusks assemblages in a tropical estuary, Guanabara Bay, Southeast Brazil. *Mar Poll Bull* 54:377–384. <https://doi.org/10.1016/j.marpolbul.2006.08.026>
- Mergaoui L, Fekhaoui M, Bouya D, Gheit A, Stambouli A (2003) Qualité des eaux et macrofaune benthique d'un milieu estuarien du Maroc : cas de l'estuaire de Sebou. *Bull De L'inst Sci Section Sci De La Vie* 25(67–7):5
- Mistri M, Rossi R, Fano EA (2001) Structure and secondary production of a soft bottom macrobenthic community in a brackish lagoon (Sacca di Goro, north-eastern Italy). *Estuar Coast Shelf Sci* 52:605–616. <https://doi.org/10.1006/ecss.2001.0757>
- Mistri M, Ghion F, Modugno S, Rossi R (2002) Response of macrobenthic communities to an hydraulic intervention in an enclosed lagoon (Valle di Gorino, northern Italy). *J Marine Biol Ass United Kingdom* 82:771–778. <https://doi.org/10.1017/S0025315402006136>
- Newton A, Brito AC, Icely JD, Derolez V, Clara I, Angus S, Schernewski G, Inácio M, Lillebø AI, Sousa AI, Béjaoui B, Solidoro C, Tosic M, Cañedo-Argüelles M, Yamamuro M, Reizopoulou S, Tseng H-C, Donata C, Roselli L, Maanan M, Cristina S, Ruiz-Fernández AC, Pastres R, Snoussi M, Turpie J, Tuchkovenko Y, Dyack B, Brookes J, Povilanskas R, Khokhlov V (2018) Assessing quantifying and valuing the ecosystem services of coastal lagoons. *J Nat Conserv* 44:50–67. <https://doi.org/10.1016/j.jnc.2018.02.009>
- Nicolaidou A, Petrou K, Kormas KA, Reizopoulou S (2006) Inter-annual variability of soft bottom macrofaunal communities in two Ionian Sea lagoons. *Hydrobiologia* 555:89–98. <https://doi.org/10.1007/s10750-005-1108-2>
- Nybakken JW (1993) *Marine biology: an ecological approach*. Harper Collins College, New York, p 445
- Pasquaud S (2006) *Les relations trophiques : éléments de structuration des peuplements ichtyologiques en milieu estuarien Application à l'estuaire de la Gironde*. Dissertation, Université de Bordeaux I
- Pearson TH, Rosenberg R (1978) Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanogr Mar Biol Ann Rev* 16:229–311
- Pielou EC (1966) Shannon's formula as a measure of specific diversity: Its use and measure. *Am Nat* 100:463–465
- Pranovi F, Curiel D, Rismondo A, Marzocchi M, Scattolin M (2000) Determination of food sources for benthic invertebrates in a salt marsh (Aiguillon Bay, France) by carbon and nitrogen stable isotopes: importance of locally produced sources. *Sci Mar* 64(3):303–310. <https://doi.org/10.3989/scimar.2000.64n3303>
- Reiss H, Kröncke I (2005) Seasonal variability of benthic indices: an approach to test the applicability of different indices for ecosystem quality assessment. *Mar Poll Bull* 50:1490–1499. <https://doi.org/10.1016/j.marpolbul.2005.06.017>
- Rharbi N, Ramdani M, Berraho A, Idrissi JL (2001) Caractéristiques hydrologiques et écologiques de la lagune d'Oualidia : milieu parallèle de la côte atlantique marocaine. *Mar Life* 11:3–9
- Rodrigues AM, Quintino V, Sampaio L, Freitas R, Neves R (2011) Benthic biodiversity patterns in Ria de Aveiro, Western Portugal: environmental-biological relationships. *Est Coast Shelf Sci* 95:338–348. <https://doi.org/10.1016/j.ecss.2011.05.019>
- Rossi F, Castelli A, Lardicci C (2006) Distribution of macrobenthic assemblages along a marine gradient in Mediterranean eutrophic coastal lagoons. *Mar Ecol* 27:66–75. <https://doi.org/10.1111/j.1439-0485.2006.00078.x>
- Salas F, Neto JM, Borja A, Marques JC (2004) Evaluation of the applicability of a marine biotic index to characterize the status of estuarine ecosystems: the case of Mondego estuary (Portugal). *Ecol Indic* 4:215–225. <https://doi.org/10.1016/j.ecolind.2004.04.003>
- Schückel U, Beck M, Kröncke I (2015) Macrofauna communities of tidal channels in Jade Bay (German Wadden Sea): spatial patterns, relationships with environmental characteristics, and comparative aspects. *Mar Biodiv* 45:841–855. <https://doi.org/10.1007/s12526-014-0308-2>
- Shannon CE (1948) A mathematical theory of communication. *Bell Syst Tech J* 27:379–423. <https://doi.org/10.1145/584091.584093>
- Solidoro C, Pastres R, Cossarini G, Ciavatta S (2004) Seasonal and spatial variability of water quality parameters in the lagoon of Venice. *J Mar Syst* 51:7–18. <https://doi.org/10.1016/j.jmarsys.2004.05.024>
- Stora G, Arnoux A, Galas M (1995) Time and spatial dynamics of Mediterranean lagoon macrobenthos during an exceptionally prolonged interruption of freshwater inputs. *Hydrobiologia* 300:123–132. <https://doi.org/10.1007/BF00024454>
- Teske P, Wooldridge T (2001) A comparison of the macrobenthic faunas of permanently open and temporarily open/closed South African estuaries. *Hydrobiologia* 464:227–243. <https://doi.org/10.1023/A:1013995302300>
- Teske PR, Wooldridge TH (2003) What limits the distribution of subtidal macrobenthos in permanently open and temporarily open/closed South African estuaries? Salinity vs sediment particle size. *Estuar Coast Shelf Sci* 57:225–238. [https://doi.org/10.1016/S0272-7714\(02\)00347-5](https://doi.org/10.1016/S0272-7714(02)00347-5)
- Tlig-zouari S, Maamouri-mokhtar F (2008) Macrozoobenthic species composition distribution in the Northern lagoon of Tunis. *Transit Waters Bull* 2:1–15. <https://doi.org/10.1285/i18252273v2n2p1>
- Uwadiae RE (2013) Spatial patterns in diversity and distribution of benthic molluscs in a weak tidal tropical lagoon. *Int J Environ Sci* 3:111–120. <https://doi.org/10.5376/ijms.2013.03.0014>
- Vezzone M, Dos Anjos RM, Cesar RG, Muniz M, Cardoso R, Felizardo JP, Vasconcelos D, Polivanov H (2020) Using stable isotopes to discriminate anthropogenic impacts of the sedimentary organic matter pollution in the Rodrigo de Freitas Lagoon (RJ, Brazil). *Environ Sci Poll Res* 28:4515–4530. <https://doi.org/10.1007/s11356-020-10835-8>
- Ysebaert T, Herman PMJ (2002) Spatial and temporal variation in benthic macrofauna and relationships with environmental variables in an estuarine, intertidal soft-sediment environment. *Mar Ecol Prog Ser* 244:105–124. <https://doi.org/10.3354/meps244105>
- Zabi SG, Leloeuf P (1993) Review of knowledge on the benthic fauna of marginal and littoral environment of West Africa 22nd part Settlements biotopes. *Rev Hydrobiol Trop* 26:19–51
- Zourarah B, Maanan M, Carruesco C, Ajjane A, Mehdi K, Conceição Freitas M (2007) Fifty-year sedimentary record of heavy metal pollution in the lagoon of Oualidia (Moroccan Atlantic coast). *Est Coast Shelf Sci* 72:359–369. <https://doi.org/10.1016/j.ecss.2006.11.007>